

# NONLINEAR ELLIPTIC EQUATIONS WITH CONCENTRATING REACTION TERMS AT AN OSCILLATORY BOUNDARY

JOSÉ M. ARRIETA, ARIADNE NOGUEIRA<sup>◊</sup> AND MARCONE C. PEREIRA<sup>†</sup>

ABSTRACT. In this paper we analyze the asymptotic behavior of a family of solutions of a semilinear elliptic equation, with homogeneous Neumann boundary condition, posed in a two-dimensional oscillating region with reaction terms concentrated in a neighborhood of the oscillatory boundary  $\theta_\varepsilon \subset \Omega_\varepsilon \subset \mathbb{R}^2$  when a small parameter  $\varepsilon > 0$  goes to zero. Our main result is concerned with the upper and lower semicontinuity of the set of solutions in  $H^1$ . We show that the solutions of our perturbed equation can be approximated with one defined in a fixed limit domain, which also captures the effects of reaction terms that take place in the original problem as a flux condition on the boundary of the limit domain.

## 1. INTRODUCTION

In this paper we analyze the asymptotic behavior of a family of steady state solutions of a semilinear reaction-diffusion equation with homogeneous Neumann boundary conditions on an oscillating domain  $\Omega_\varepsilon \subset \mathbb{R}^2$ , with reaction terms concentrated in an extremely thin region  $\theta_\varepsilon$  close to the border  $\partial\Omega_\varepsilon$  which can also present oscillatory structure. In Figure 1 we illustrate the oscillating domain  $\Omega_\varepsilon$ , as well as, the narrow oscillating neighborhood  $\theta_\varepsilon$  where some reactions of the model take place.

Under our assumptions, the two-dimensional family of oscillating regions  $\Omega_\varepsilon$  approaches a bounded domain  $\Omega \subset \mathbb{R}^2$ , and the narrow strip  $\theta_\varepsilon$ , that may also have an oscillatory behavior, degenerates into a fixed set  $\Gamma \subset \partial\Omega$  as the positive parameter  $\varepsilon$  goes to zero. We assume that the reactions of our model occur in the interior of  $\Omega_\varepsilon \subset \mathbb{R}^2$  as well as in the narrow strip  $\theta_\varepsilon$ .




FIGURE 1. The oscillatory domain  $\Omega_\varepsilon$  and strip  $\theta_\varepsilon$  where reactions take place.

We show that the family of solutions is upper semicontinuous at  $\varepsilon = 0$ , and under the additional condition on hyperbolicity at the limit problem, we provide the lower semicontinuity. Also, we show that the perturbed equation has one and only one solution nearby to the limit equation as  $\varepsilon \rightarrow 0$ . Indeed, we show that the starting singular equation defined in the perturbed two-dimensional region can be approximated with one which is defined in the limit fixed domain  $\Omega$  and captures all relevant effects of the processes that take

---

*Key words and phrases.* Semilinear elliptic equations, nonlinear boundary conditions, singular elliptic equations, concentrating terms, oscillatory behavior.

<sup>◊</sup>Supported by CNPq 141869/2013-5, Brazil, <sup>†</sup>Partially supported by CNPq 303253/2017-7 and FAPESP 2017/02630-2, Brazil.

place in the original problem. Moreover, we will see that the nonlinear concentrating term of the perturbed equation will become a nonlinear boundary condition in the limit case, according to pioneering works as [11, 21, 22].

Indeed, we consider such kind of domains that satisfy  $\Omega_\varepsilon \rightarrow \Omega$  and  $\partial\Omega_\varepsilon \rightarrow \partial\Omega$  as  $\varepsilon \rightarrow 0$  in the sense of Hausdorff. We take  $\text{dist}(\Omega_\varepsilon, \Omega) + \text{dist}(\partial\Omega_\varepsilon, \partial\Omega) \rightarrow 0$  when  $\varepsilon \rightarrow 0$ , with  $\text{dist}$  being the symmetric Hausdorff distance of two sets. Notice that there is no condition on the intersection of the sets  $\Omega_\varepsilon$  and  $\Omega$  improving some results from [1, 2] with respect to the class of domain perturbation. In Section 2, we set our assumptions and introduce our main result precisely.

As we will see, in order to show our results, we need to estimate and analyze the asymptotic behavior of concentrating integrals such as

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u(x)|^q dx \quad (1.1)$$

for different values of  $q \geq 1$  and open sets  $\theta_\varepsilon \subset \Omega_\varepsilon \subset \mathbb{R}^2$ . Notice the factor  $1/\varepsilon$  in (1.1). The arrangement of this one with the narrow strip can be thought as a model to measure the concentration of  $u$  on  $\theta_\varepsilon$  at  $\varepsilon = 0$ . In fact, a suitable control of this integral is useful to analyze models set in regions of  $\mathbb{R}^2$  which present singular behavior. For instance, we mention our recent work [12] where an oscillating thin domain is studied.

Here, we are in agreement with pioneering papers [11, 21, 22] calling (1.1) as concentrating or concentrated integral. Indeed, this kind of problem was initially proposal in [11] where linear elliptic equations were considered with reaction and potential terms concentrated on the boundary. There, the neighborhood  $\theta_\varepsilon$  has been set as a strip without oscillatory behavior in a fixed domain  $\Omega$ . Later, the dynamical system given by a semilinear parabolic problem in the same situation was analyzed in [21, 22] where the upper semicontinuity of attractors at  $\varepsilon = 0$  has been shown. In [3, 4] the results of [11, 21] were extended to a reaction-diffusion problem with delay. In these works, the boundary of the domain is always assumed to be smooth.

Subsequently some results of [11] were adapted in [5] to be considered in a semilinear elliptic problem posed on a Lipschitz fixed domain  $\Omega$  with the  $\varepsilon$ -neighborhood presenting highly oscillatory behavior. The upper and lower semicontinuity of the attractor to the associated parabolic problem in smooth fixed domains were shown in [6].

Recently, some results from [11, 5] have been adapted in [1, 2] to a class of narrow strips  $\theta_\varepsilon$  and bounded oscillatory domains  $\Omega_\varepsilon$ . Under the restricted assumption  $\Omega \subset \Omega_\varepsilon$  and  $\theta_\varepsilon \subset \Omega_\varepsilon \setminus \Omega$  for all  $\varepsilon > 0$ , the authors have been able to estimate concentrating integrals and analyze the asymptotic behavior of semilinear elliptic equations as  $\Omega_\varepsilon \rightarrow \Omega$  and  $\partial\Omega_\varepsilon \rightarrow \partial\Omega$  when  $\varepsilon \rightarrow 0$  in the sense of Hausdorff.

This paper is organized as follows: in Section 2 we introduce the assumptions, notations and the main result. In Section 3, we show some technical results concerning to extension operators, Lebesgue-Bochner and Sobolev-Bochner generalized spaces needed to get our estimates. Following by Section 4, we prove some properties to concentrating integrals which are used in Section 5 to study the nonlinearities of our problem. Finally, in Section 6, we pass to the limit in a semilinear elliptic problem getting the upper semicontinuity of the solutions. Moreover, assuming hyperbolicity to the solutions of the limit equation, we also obtain the lower semicontinuity at  $\varepsilon = 0$ , and we will exclude the possibility that, near an equilibrium point of the limiting equation, may exist several different equilibrium points of the perturbed problem, and therefore, we will also prove some sort of uniqueness of the equilibrium points.

## 2. ASSUMPTIONS, NOTATIONS AND MAIN RESULT

We study a family of solutions of the following semilinear elliptic equation with homogeneous Neumann boundary condition,

$$\begin{cases} -\Delta u^\varepsilon + u^\varepsilon = \Phi(u^\varepsilon) + \frac{1}{\varepsilon} \chi^{\theta_\varepsilon} f(u^\varepsilon) & \text{in } \Omega_\varepsilon, \\ \frac{\partial u^\varepsilon}{\partial \nu^\varepsilon} = 0 & \text{on } \partial\Omega_\varepsilon, \end{cases} \quad (2.1)$$

where

$$\begin{aligned} \Omega_\varepsilon &= \{(x_1, x_2) \in \mathbb{R}^2; x_1 \in (0, 1), 0 < x_2 < G_\varepsilon(x_1)\} \quad \text{and} \\ \theta_\varepsilon &= \{(x_1, x_2) \in \mathbb{R}^2; x_1 \in (0, 1), G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1) < x_2 < G_\varepsilon(x_1)\} \end{aligned} \quad (2.2)$$

are set by functions  $G_\varepsilon, H_\varepsilon : (0, 1) \rightarrow \mathbb{R}$  satisfying conditions:

- H(i)**  $G_\varepsilon(x_1) = m(x_1) + \varepsilon g(x_1/\varepsilon^\alpha)$  with  $0 < \alpha \leq 1$ , where
- (a)  $m : (0, 1) \rightarrow \mathbb{R}$  is  $C^1$ , bounded, with bounded derivative,
  - (b)  $g : (0, 1) \rightarrow \mathbb{R}$  is a  $C^1$  bounded function,  $L_g$ -periodic with bounded derivative.
  - (c)  $G_\varepsilon \rightarrow m$  as  $\varepsilon \rightarrow 0$  uniformly in  $(0, 1)$ .
  - (d) there are constants  $G_0, G_1 > 0$  such that  $G_0 \leq G_\varepsilon(x) \leq G_1$  for all  $x \in (0, 1)$ .
- H(ii)**  $H_\varepsilon(x_1) = h(x_1/\varepsilon^\beta)$ ,  $\beta > 0$ , where the function  $h$  is bounded, ie there are  $H_0, H_1 \geq 0$  such that  $H_0 \leq H_\varepsilon(x) \leq H_1$  for all  $x \in (0, 1)$ , and  $L_h$ -periodic.

The vector  $\nu^\varepsilon = (\nu_1^\varepsilon, \nu_2^\varepsilon)$  is the unit outward normal vector to the boundary  $\partial\Omega_\varepsilon$ ,  $\partial/\partial\nu^\varepsilon$  is the derivative in the direction of  $\nu^\varepsilon$ , and  $\chi^{\theta_\varepsilon}$  is the characteristic function of the neighborhood  $\theta_\varepsilon$ . The nonlinearities  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  and  $f : \mathbb{R} \rightarrow \mathbb{R}$  are bounded functions of class  $C^2$  with bounded derivatives.

Under assumptions **H**, it is not difficult to associate to (2.2) with the following limit sets

$$\begin{aligned} \Omega &= \{(x_1, x_2) \in \mathbb{R}^2; x_1 \in (0, 1), 0 < x_2 < m(x_1)\} \text{ and} \\ \Gamma &= \{(x_1, x_2) \in \mathbb{R}^2; x_1 \in (0, 1), x_2 = m(x_1)\}. \end{aligned} \quad (2.3)$$

We may pass to the limit in the solutions of the equation (2.1) getting the following limit equation

$$\begin{cases} -\Delta u + u = \Phi(u) & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = \hat{\mu} f(u) & \text{on } \Gamma, \end{cases} \quad (2.4)$$

with  $\hat{\mu} \in L^\infty(\Gamma)$  given by

$$\hat{\mu} = \frac{\mu_h}{\sqrt{1 + m'^2}} \in L^\infty(\Gamma), \quad (2.5)$$

where  $\mu_h \in L^\infty(\Gamma)$  is the weak\* limit of  $H_\varepsilon$ . In fact, due to **H(ii)**, it follows from [17, Teorema 2.6] that

$$H_\varepsilon \rightharpoonup \mu_h = \frac{1}{L_h} \int_0^{L_h} h(s) ds.$$

The coefficient  $\hat{\mu}$  captures the influence of the small neighborhood  $\theta_\varepsilon$ , as well as the geometry of the limit domain  $\Omega$ . It also suggests with nonlinearity  $f$  a flux condition on the boundary, giving a qualitative idea on the effect of the concentrating reaction terms on the original problem.

Notice that, to obtain the convergence results, we have to compare functions defined in different functional spaces as  $\varepsilon \rightarrow 0$ . In order to do that, we consider the following family of operators

$$E_\varepsilon : H^1(\Omega) \rightarrow H^1(\Omega_\varepsilon) : u \mapsto E_\varepsilon u := R_\varepsilon P u \quad (2.6)$$

where  $R_\varepsilon : H^1(\mathbb{R}^2) \rightarrow H^1(\Omega_\varepsilon)$  is the restriction operator to the open set  $\Omega_\varepsilon$  and  $P : H^1(\Omega) \rightarrow H^1(\mathbb{R}^2)$  is a continuous extension operator from functions defined in  $\Omega$  to the whole plane  $\mathbb{R}^2$ . The existence of  $P$  is guaranteed by [20, Theorem 1.4.3.1].

From [7], we have

$$\|E_\varepsilon u\|_{H^1(\Omega_\varepsilon)} \rightarrow \|u\|_{H^1(\Omega)}, \quad \text{as } \varepsilon \rightarrow 0,$$

and then, we can compare solutions from (2.1) and (2.4) using the notion of  $E$ -convergence as in [14].

In general, consider a family of Banach spaces  $H_\varepsilon$  and a limit Banach space  $H_0$ . Besides, let  $E_\varepsilon : H_0 \rightarrow H_\varepsilon$  a family of operators such that  $\|E_\varepsilon u\|_{H_\varepsilon} \rightarrow \|u\|_{H_0}$  when  $\varepsilon \rightarrow 0$ .

**Definition 2.1.** *We say that a sequence of  $u^\varepsilon \in H_\varepsilon$   $E$ -converges to  $u_0 \in H_0$ , if  $\|u^\varepsilon - E_\varepsilon u\|_{H_\varepsilon} \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . We denote this convergence by  $u_\varepsilon \xrightarrow{E} u$ .*

If  $H_\varepsilon$  and  $H_0$  are Hilbert spaces, we can define a weak  $E$ -convergence.

**Definition 2.2.** *A sequence of  $\{u^\varepsilon\}$ , with  $u^\varepsilon \in H_\varepsilon$ ,  $E$ -converges weakly to  $u \in H_0$  if for any sequence  $E$ -convergent to  $w$  we have  $(w^\varepsilon, u^\varepsilon)_{H_\varepsilon} \rightarrow (u, w)_{H_0}$  when  $\varepsilon \rightarrow 0$ . We may denote such convergence by  $u^\varepsilon \xrightarrow{E} u$ .*

We also need a notion of compactness for sequences, and convergence for operators which are defined in different spaces. We recall the exposition from [14]. See also [7] and [12].

**Definition 2.3.** A sequence  $\{u_n\}$ ,  $u_n \in H_{\varepsilon_n}$  with  $\varepsilon_n \rightarrow 0$ , is *E-precompact* if for all subsequence  $\{u_{n'}\}$  there are a subsequence  $\{u_{n''}\}$  and an element  $u \in H_0$  such that  $u_{n''} \xrightarrow{E} u$ . A family is said to be *E-precompact* if all sequence  $\{u_n\}$ ,  $u_n \in H_{\varepsilon_n}$  with  $\varepsilon_n \rightarrow 0$ , is *E-precompact*.

**Definition 2.4.** We say that a family of operators  $\{T_\varepsilon\}$ , with  $T_\varepsilon : H_\varepsilon \rightarrow H_\varepsilon$ , *E-converges* to  $T : H_0 \rightarrow H_0$  when  $\varepsilon \rightarrow 0$  if  $T_\varepsilon u^\varepsilon \xrightarrow{E} Tu$  for any  $u^\varepsilon \xrightarrow{E} u$ . We denote this convergence by  $T_\varepsilon \xrightarrow{EE} T$ .

Furthermore we may define a notion of compact convergence for operators.

**Definition 2.5.** A family of compact operators  $\{T_\varepsilon\}$ , with  $T_\varepsilon : H_\varepsilon \rightarrow H_\varepsilon$ , *converges compactly* to  $T : H_0 \rightarrow H_0$  when  $\varepsilon \rightarrow 0$  if, for any family  $\{u^\varepsilon\}$  with  $\|u^\varepsilon\|_{H_\varepsilon}$  uniformly bounded, we have that  $\{T_\varepsilon u^\varepsilon\}$  is *E-precompact* and  $T_\varepsilon \xrightarrow{EE} T$ . We denote this compact convergence by  $T_\varepsilon \xrightarrow{CC} T$ .

This notion of convergence can be extended to sets in the following manner: let  $J_\varepsilon$  be a family of sets in some Banach spaces  $Z_\varepsilon$ . We say that  $J_\varepsilon$  is

- (i) upper semicontinuous at  $\varepsilon = 0$ , if  $\text{dist}_H(J_\varepsilon, E_\varepsilon J_0) \xrightarrow{\varepsilon \rightarrow 0} 0$ ;
- (ii) lower semicontinuous at  $\varepsilon = 0$ , if  $\text{dist}_H(E_\varepsilon J_0, J_\varepsilon) \xrightarrow{\varepsilon \rightarrow 0} 0$ .

Here,  $\text{dist}_H(A, B)$  denotes the Hausdorff semi-distance given by

$$\text{dist}_H(A, B) = \sup_{x \in A} \inf_{y \in B} \|x - y\|_{Z_\varepsilon}.$$

**Remark 2.6.** In order to show the upper or lower semicontinuity of sets, the following characterizations are useful:

- (i) The family  $\{J_\varepsilon\}$  is upper semicontinuous at  $\varepsilon = 0$ , if every sequence  $\{u_\varepsilon\}$ , with  $u_\varepsilon \in J_\varepsilon$  and  $\varepsilon \rightarrow 0$ , has a subsequence *E-convergent* to an element of  $J_0$ ;
- (ii) The family  $\{J_\varepsilon\}$  is lower semicontinuous at  $\varepsilon = 0$ , if  $J_0$  is compact and for all  $u \in J_0$  exists a sequence  $\{u_\varepsilon\}$ , with  $u_\varepsilon \in J_\varepsilon$  and  $\varepsilon \rightarrow 0$ , such that  $u_\varepsilon \xrightarrow{E} u$ .

Finally, for  $\varepsilon > 0$ , let us consider

$$\mathcal{E}_\varepsilon = \{u^\varepsilon \in H^1(\Omega_\varepsilon); u^\varepsilon \text{ is a solution of (2.1)}\}$$

and

$$\mathcal{E}_0 = \{u \in H^1(\Omega); u \text{ is a solution of (2.4)}\}.$$

The main goal of this work is to prove the upper and lower semicontinuity of the set  $\mathcal{E}_\varepsilon$  at  $\varepsilon = 0$ :

**Theorem 2.7.** If we consider the semilinear elliptic problem (2.1) then:

- (i) for any sequence  $u^\varepsilon \in \mathcal{E}_\varepsilon$ , with  $\varepsilon \rightarrow 0$ , there is a subsequence (also denoted by  $u^\varepsilon$ ) and  $u_0 \in \mathcal{E}_0$  such that  $u^\varepsilon \xrightarrow{E} u_0$ .
- (ii) for any hyperbolic equilibrium point  $u^* \in \mathcal{E}_0$ , there is sequence  $u^\varepsilon \in \mathcal{E}_\varepsilon$  such that  $u^\varepsilon \xrightarrow{E} u^*$  when  $\varepsilon \rightarrow 0$ . Moreover, there are  $\eta > 0$  and  $\varepsilon_0 > 0$  such that exists an unique  $u^\varepsilon \in \mathcal{E}_\varepsilon$  which satisfies

$$\|u^\varepsilon - E_\varepsilon u^*\|_{H^1(\Omega_\varepsilon)} \leq \eta, \quad \text{for all } 0 < \varepsilon < \varepsilon_0.$$

**Remark 2.8.** Recall that  $u^*$  is a hyperbolic equilibrium point of (2.4), if  $\lambda = 0$  is not an eigenvalue of the linearized problem of (2.4) around  $u^*$ . For instance, if  $u^*$  is solution of (2.4) and is hyperbolic, then  $\lambda = 0$  is not an eigenvalue of

$$\begin{cases} -\Delta v + v = \Phi'(u^*)v + \lambda v & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = \hat{\mu} f'(u^*)v & \text{on } \Gamma. \end{cases}$$

Furthermore, we notice that item (ii) of the Theorem 2.7 also give us a kind of uniqueness result to the solutions near a hyperbolic equilibrium point of the limit equation for sufficiently small  $\varepsilon$ .

## 3. FUNCTIONAL SPACES AND TECHNICAL RESULTS

In this section, we introduce the main functional spaces used throughout this paper and work with some of their properties. Then we set some technical results that will be useful in next sections. First, we define fractional Sobolev spaces.

**Definition 3.1.** For  $s > 0$ ,  $1 \leq p < \infty$  and  $O \subset \mathbb{R}^n$ , we denote by  $W^{s,p}(O)$  and call fractional Sobolev space, the functional set given by the space of distributions defined in  $O$  such that

- (i)  $\partial^\alpha u \in L^p(O)$ , for  $|\alpha| \leq m$ , when  $s = m \in \mathbb{N}$ ;
- (ii)  $u \in W^{m,p}(O)$  and

$$\iint_{O \times O} \frac{|\partial^\alpha u(x) - \partial^\alpha u(y)|^p}{|x - y|^{n+\sigma p}} dx dy < \infty,$$

for  $|\alpha| = m$ , when  $s = m + \sigma$  with  $\sigma \in (0, 1)$ .

The norm in  $W^{s,p}(O)$ , that makes it Banach, is:

$$\|u\|_{W^{m,p}(O)}^p = \sum_{|\alpha| \leq m} \int_O |\partial^\alpha u(x)|^p dx \quad \text{in the case (i)}$$

and

$$\|u\|_{W^{s,p}(O)}^p = \|u\|_{W^{m,p}(O)}^p + \sum_{|\alpha|=m} \iint_{O \times O} \frac{|\partial^\alpha u(x) - \partial^\alpha u(y)|^p}{|x - y|^{n+\sigma p}} dx dy \quad \text{in the case (ii)}.$$

Besides if  $p = 2$  we denote it by  $H^s(O)$ , which is a Hilbert space.

Now let us introduce the Lebesgue and Sobolev-Bochner generalized spaces. Here, they are given in a similar way to [24], as a natural generalization to the Lebesgue and Sobolev spaces using Bochner integrals. The usual Lebesgue and Sobolev-Bochner spaces may be found, for instance, in [15, 17].

**Definition 3.2.** Let us consider a function  $G : (0, 1) \rightarrow \mathbb{R}$  satisfying  $0 < G_0 \leq G(x) \leq G_1$ ,  $\forall x \in (0, 1)$ , for some constants  $0 < G_0 \leq G_1$ . Let  $1 \leq p \leq \infty$  e  $1 \leq q < \infty$ . The Lebesgue-Bochner generalized spaces, denoted by  $L^p(0, 1; L^q(0, G(x_1)))$ , are defined by

$$L^p(0, 1; L^q(0, G(x_1))) := \{u : \Omega_\varepsilon \rightarrow \mathbb{R} \text{ measurable ; } u(x_1, \cdot) \in L^q(0, G(x_1)) \text{ for almost every } x_1 \in (0, 1)\}.$$

They are Banach spaces with the norm

$$\|u\|_{L^p(0,1;L^q(0,G(x_1)))} = \begin{cases} \left( \int_0^1 \|u(x_1, \cdot)\|_{L^q(0,G(x_1))}^p dx_1 \right)^{1/p}, & p < \infty, \\ \text{ess sup}_{x \in (0,1)} \|u(x_1, \cdot)\|_{L^q(0,G(x_1))}, & p = \infty. \end{cases}$$

When  $p = q = 2$  such space is Hilbert with the inner product

$$(u, v)_{L^2(0,1;L^2(0,G(x_1)))} = \int_0^1 (u(x_1, \cdot), v(x_1, \cdot))_{L^2(0,G(x_1))} dx_1.$$

**Remark 3.3.** Since  $q < \infty$ , the function  $x_1 \mapsto \|u(x_1, \cdot)\|_{L^q(0,G(x_1))}$  is measurable by Fubini's Theorem. Then the space  $L^p(0, 1; L^q(0, G(x_1)))$  is well defined.

Analogously, the Sobolev-Bochner generalized spaces, denoted by  $L^p(0, 1; W^{s,q}(0, G(x_1)))$  for  $s > 0$ , are defined by

$$L^p(0, 1; W^{s,q}(0, G(x_1))) := \{u \in L^p(0, 1; L^q(0, G(x_1))); u(x_1, \cdot) \in W^{s,q}(0, G(x_1))\}.$$

Such spaces are Banach with the norm

$$\|u\|_{L^p(0,1;W^{s,q}(0,G(x_1)))} = \begin{cases} \left( \int_0^1 \|u(x_1, \cdot)\|_{W^{s,q}(0,G(x_1))}^p dx_1 \right)^{1/p}, & p < \infty, \\ \text{ess sup}_{x_1 \in (0,1)} \|u(x_1, \cdot)\|_{W^{s,q}(0,G(x_1))}, & p = \infty, \end{cases}$$

and, again, they are Hilbert spaces if  $p = q = 2$ .

In general, it follows from [17, Proposition 3.59] that, if  $H$  is a Hilbert space and  $1 \leq p < \infty$ , then the dual space of  $L^p(0, 1; H)$  is given by

$$[L^p(0, 1; H)]' = L^q(0, 1; H'),$$

where  $H'$  is the dual space of  $H$  and  $p, q$  are conjugates.

In our case we will consider the family of Lebesgue and Sobolev-Bochner generalized spaces for the function  $G_\varepsilon(x_1) = m(x_1) + \varepsilon g(x_1/\varepsilon^\alpha)$  defined in hypothesis **H**(i) from (2.2).

Now we set important and nontrivial results that will help us to work with different definitions of Sobolev fractional spaces making their norms equivalent. The proofs are analogous to [12, Proposition 3.4, 3.5 and 3.6].

**Lemma 3.4.** *Fixed  $\varepsilon > 0$  and  $x_1 \in (0, 1)$ , if we call  $I_\varepsilon = (0, G_\varepsilon(x_1))$ , there is a continuous linear extension operator  $P : L^2(I_\varepsilon) \rightarrow L^2(\mathbb{R})$  such that  $Pu = u$  in  $I_\varepsilon$ , with  $\|Pu\|_{L^2(\mathbb{R})} \leq \lambda_0 \|u\|_{L^2(I_\varepsilon)}$ ,  $\|Pu\|_{H^s(\mathbb{R})} \leq \lambda_s \|u\|_{H^s(I_\varepsilon)}$  and  $\|Pu\|_{H^1(\mathbb{R})} \leq \lambda_1 \|u\|_{H^1(I_\varepsilon)}$ , for  $0 < s < 1$ , where the constants  $\lambda_0, \lambda_s, \lambda_1 \geq 1$  are independent of  $\varepsilon > 0$  and  $x_1 \in (0, 1)$ .*

**Theorem 3.5.** *Let  $I_\varepsilon = (0, G_\varepsilon(x_1))$ , with  $\varepsilon > 0$ ,  $x_1 \in (0, 1)$  and  $0 < s < 1$  fixed. Then there are  $C_1, C_2 > 0$  independent of  $\varepsilon$  such that*

$$C_1 \|u\|_{H^s(I_\varepsilon)} \leq \|u\|_{H_{\square}^s(I_\varepsilon)} \leq C_2 \|u\|_{H^s(I_\varepsilon)}, \quad \forall u \in H^s(I_\varepsilon),$$

where  $H_{\square}^s(I_\varepsilon)$  is the complex interpolation space

$$H_{\square}^s(I_\varepsilon) = [L^2(I_\varepsilon), H^1(I_\varepsilon)]_s, \quad \text{for } 0 < s < 1.$$

**Proposition 3.6.** *For each  $\varepsilon > 0$ ,  $H^1(\Omega_\varepsilon) \subseteq L^2(0, 1; H^s(0, G_\varepsilon(x_1)))$  for all  $0 \leq s \leq 1$ , with constant of inclusion independent of  $\varepsilon$ . Besides  $H^1(\Omega_\varepsilon) \subseteq L^2(0, 1; H^s(0, G_\varepsilon(x_1)))$  with compact immersion if  $0 < s < 1$ .*

According to the properties of our domains  $\Omega_\varepsilon$  defined in (2.2), we also have the important result.

**Proposition 3.7.** *The family  $\Omega_\varepsilon$  admits a continuous extension operator  $P_\varepsilon : L^2(\Omega_\varepsilon) \rightarrow L^2(U)$ , where the open set  $U = U_1 \times U_2 \subset \mathbb{R}^2$  is such that the closure of  $\Omega_\varepsilon$  is contained in  $U$  for all  $\varepsilon > 0$ , and*

$$\begin{aligned} \|P_\varepsilon u^\varepsilon\|_{H^1(U)} &\leq C_0 \|u^\varepsilon\|_{H^1(\Omega_\varepsilon)}, \\ \|P_\varepsilon u^\varepsilon\|_{L^2(U_1; H^s(U_2))} &\leq C_s \|u^\varepsilon\|_{L^2(0, 1; H^s(0, G_\varepsilon(x)))}, \\ \|P_\varepsilon u^\varepsilon\|_{L^2(U)} &\leq C_1 \|u^\varepsilon\|_{L^2(\Omega_\varepsilon)}, \end{aligned}$$

where the constants  $C_0, C_s, C_1 > 0$  are independent of  $\varepsilon > 0$  and  $0 \leq s \leq 1$ .

*Proof.* By hypothesis **H**(i), we have  $|G'_\varepsilon(x)| \leq C$  for all  $x \in (0, 1)$ , with  $C > 0$  independent of  $\varepsilon > 0$ . Thus, the proof follows from the extension operator defined in [10, Lemma 3.1].  $\square$

Our next step is to prove some inclusions involving Sobolev fractional spaces and Sobolev-Bochner generalized spaces that will be useful in the further analysis of concentrating integrals.

**Proposition 3.8.** *For  $\varepsilon > 0$  and considering the domains defined in (2.2), the following inclusions hold with immersion constants independent of  $\varepsilon$ .*

- (a)  $H^1(\Omega_\varepsilon) \subset L^\infty(0, 1; L^2(0, G_\varepsilon(x)))$ ;
- (b) if  $q \geq 2$  then  $H^1(\Omega_\varepsilon) \subset L^q(0, 1; H^s(0, G_\varepsilon(x)))$ , where  $s = 2/q$ ;
- (c)  $H^1(\Omega_\varepsilon) \subset L^q(\Omega_\varepsilon)$ , for  $2 \leq q \leq 6$ .

*Proof.* (a) For each  $x_1 \in (0, 1)$ , we can use the extension operator given by Proposition 3.7 to get

$$\|u(x_1, \cdot)\|_{L^2(0, G_\varepsilon(x_1))} \leq \|P_\varepsilon u(x_1, \cdot)\|_{L^2(0, G_1)}.$$

Hence,

$$\|u\|_{L^\infty(0,1;L^2(0,G_\varepsilon(x_1)))} \leq \|P_\varepsilon u\|_{L^\infty(0,1;L^2(0,G_1))}. \quad (3.1)$$

From [15, Corollary 1.4.36] follows that

$$\|P_\varepsilon u\|_{L^\infty(0,1;L^2(0,G_1))} \leq C\|P_\varepsilon u\|_{H^1(0,1;L^2(0,G_1))} \leq C\|P_\varepsilon u\|_{H^1(U)}. \quad (3.2)$$

Thus, using (3.2) in (3.1), and the continuity of  $P_\varepsilon$  with constant  $C_1 = \|P_\varepsilon\|_{\mathcal{L}(H^1(\Omega_\varepsilon), H^1(U))}$  uniformly bounded for each  $\varepsilon$ , we have

$$\|u\|_{L^\infty(0,1;L^2(0,G_\varepsilon(x_1)))} \leq C\|P_\varepsilon u\|_{H^1(U)} \leq CC_1\|u\|_{H^1(\Omega_\varepsilon)},$$

which concludes the proof.

- (b) First of all, let  $q \geq 2$  and define  $s = 2/q$ ,  $0 < s \leq 1$ . For each  $x_1 \in (0, 1)$  fixed, we have by Theorem 3.5 and properties of interpolation spaces that there exists  $C > 0$ , independent of  $\varepsilon$  and  $x_1$ , such that

$$\|u(x_1, \cdot)\|_{H^s(0,G_\varepsilon(x_1))} \leq C\|u(x_1, \cdot)\|_{H^1_0(0,G_\varepsilon(x_1))} \leq C\|u(x_1, \cdot)\|_{L^2(0,G_\varepsilon(x_1))}^{1-s} \|u(x_1, \cdot)\|_{H^1(0,G_\varepsilon(x_1))}^s.$$

Since by the item (a)  $H^1(\Omega_\varepsilon) \subset L^\infty(0, 1; L^2(0, G_\varepsilon(x_1)))$ , we have that

$$\|u(x_1, \cdot)\|_{H^s(0,G_\varepsilon(x_1))} \leq C\|u\|_{H^1(\Omega_\varepsilon)}^{1-s} \|u(x_1, \cdot)\|_{H^1(0,G_\varepsilon(x_1))}^s.$$

On the other hand, Proposition 3.6 implies  $H^1(\Omega_\varepsilon) \subset L^2(0, 1; H^1(0, G_\varepsilon(x_1)))$ , and then,

$$\begin{aligned} \|u\|_{L^q(0,1;H^s(0,G_\varepsilon(x_1)))}^q &= \int_0^1 \|u(x_1, \cdot)\|_{H^s(0,G_\varepsilon(x_1))}^{2/s} dx_1 \\ &\leq \int_0^1 \left( C\|u\|_{H^1(\Omega_\varepsilon)}^{1-s} \|u(x_1, \cdot)\|_{H^1(0,G_\varepsilon(x_1))}^s \right)^{2/s} dx_1 \\ &\leq C^{2/s} \|u\|_{H^1(\Omega_\varepsilon)}^{2(1-s)/s} \int_0^1 \|u(x_1, \cdot)\|_{H^1(0,G_\varepsilon(x_1))}^2 dx_1 \\ &\leq C^{2/s} \|u\|_{H^1(\Omega_\varepsilon)}^{2(1-s)/s} \|u\|_{H^1(\Omega_\varepsilon)}^2 = C^{2/s} \|u\|_{H^1(\Omega_\varepsilon)}^{2/s} = C^q \|u\|_{H^1(\Omega_\varepsilon)}^q. \end{aligned}$$

Thus,  $H^1(\Omega_\varepsilon) \subseteq L^q(0, 1; H^{2/q}(0, G_\varepsilon(x_1)))$  for  $q \geq 2$ .

- (c) Since  $L^q(\Omega_\varepsilon) = L^q(0, 1; L^q(0, G_\varepsilon(x_1)))$  isometrically, we conclude the proof by item (b) if we show

$$H^{2/q}(0, G_\varepsilon(x_1)) \subseteq L^q(0, G_\varepsilon(x_1))$$

with constant of inclusion independent of  $x_1 \in (0, 1)$  and  $\varepsilon > 0$ .

If  $q = 2$ , it follows from the definition of the spaces. If  $2 < q \leq 4$ , then  $1/2 \leq 2/q < 1$ . Hence, by [27, Theorem 1.36] we get

$$H^{2/q}(\mathbb{R}) \subseteq H^{1/2}(\mathbb{R}) \subseteq L^r(\mathbb{R}), \quad \forall r \geq 2.$$

In particular, it holds for  $r = q$  with  $2 \leq q \leq 4$ . Besides, by the operator  $P : H^s(0, G_\varepsilon(x_1)) \rightarrow H^s(\mathbb{R})$  from Lemma 3.4, whose norm is independent of  $\varepsilon > 0$  and  $x_1 \in (0, 1)$  for any  $1/2 < s < 1$ , we have

$$\begin{aligned} \|u(x_1, \cdot)\|_{L^q(0,G_\varepsilon(x_1))} &\leq \|Pu(x_1, \cdot)\|_{L^q(\mathbb{R})} \leq C\|Pu(x_1, \cdot)\|_{H^{1/2}(\mathbb{R})} \\ &\leq C\|Pu(x_1, \cdot)\|_{H^{2/q}(\mathbb{R})} \leq C\|P\| \|u(x_1, \cdot)\|_{H^{2/q}(0,G_\varepsilon(x_1))}. \end{aligned}$$

Finally, if  $4 < q \leq 6$ , then  $1/3 \leq 2/q < 1/2$ . Again by [27, Theorem 1.36], we get

$$H^{2/q}(\mathbb{R}) \subseteq L^r(\mathbb{R}), \quad \forall 2 \leq r \leq \frac{2}{1-2s}.$$

In particular, since  $q = \frac{2}{s} \leq \frac{2}{1-2s}$ , we obtain that

$$H^{2/q}(\mathbb{R}) \subseteq L^q(\mathbb{R}).$$

Hence, we conclude the proof arguing as in the previous case  $2 \leq q \leq 4$ .  $\square$

## 4. CONCENTRATING INTEGRALS AND ITS BEHAVIOR AT THE LIMIT

Our first results are about concentrating integrals. Notice that some estimates are given in different functional spaces. Under the conditions of Proposition 3.8, we may improve [12, Theorem 3.7] estimating the concentrated integrals with the  $H^1(\Omega_\varepsilon)$  norm.

**Theorem 4.1.** *For  $\varepsilon_0 > 0$  sufficiently small, there is a constant  $C > 0$ , independent of  $\varepsilon \in (0, \varepsilon_0)$  and  $u^\varepsilon \in H^1(\Omega_\varepsilon)$ , such that, for all  $1/2 < s \leq 1$ ,  $0 < \varepsilon < \varepsilon_0$ , we have*

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u^\varepsilon|^q \leq C \|u^\varepsilon\|_{L^q(0,1;H^s(0,G_\varepsilon(x_1)))}^q, \quad \forall q \geq 1, \quad (4.1)$$

and

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u^\varepsilon|^2 \leq C \left( \|u^\varepsilon\|_{H^s(\Omega_\varepsilon)}^2 + \left\| \frac{\partial u^\varepsilon}{\partial x_2} \right\|_{L^2(\Omega_\varepsilon)}^2 \right). \quad (4.2)$$

In particular,

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u^\varepsilon|^q \leq C \|u^\varepsilon\|_{H^1(\Omega_\varepsilon)}^q, \quad 2 \leq q < 4. \quad (4.3)$$

*Proof.* Take  $u \in H^1(\Omega_\varepsilon)$ . In a.e.  $x_1 \in (0, 1)$ , we have  $u(x_1, \cdot) \in H^1(0, G_\varepsilon(x_1))$ . Define

$$z^* := G_0 - \varepsilon_0 H_1 \text{ and } z^\varepsilon := G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)$$

for  $\varepsilon_0 > 0$  sufficiently small in such way that, for all  $\varepsilon < \varepsilon_0$ , we have

$$[z^\varepsilon - z^*, z^\varepsilon] \subset [0, G_\varepsilon(x_1)].$$

See Figure 2 for a representation:




FIGURE 2. Fixed  $x_1 \in (0, 1)$  and  $\varepsilon > 0$ , we get this fiber to the oscillatory domain for  $\varepsilon < \varepsilon_0$ .

Since  $(G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)) < x_2 < G_\varepsilon(x_1)$  and  $1/2 < s \leq 1$ , it follows from [20, Theorem 1.5.1.3] for  $n = 1$  that there exists  $K > 0$  independent of  $\varepsilon > 0$  such that

$$|u(x_1, x_2)| \leq K \|u(x_1, \cdot)\|_{H^s(x_2 - z^*, x_2)} \leq K \|u(x_1, \cdot)\|_{H^s(0, G_\varepsilon(x_1))}.$$

Indeed, the interval where we are applying the result is fixed and independent of the parameters  $\varepsilon > 0$  and  $x_1 \in (0, 1)$ .

Hence,

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u|^q &= \int_0^1 \frac{1}{\varepsilon} \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} |u(x_1, x_2)|^q dx_2 dx_1 \\ &\leq \int_0^1 \frac{1}{\varepsilon} \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} K^q \|u(x_1, \cdot)\|_{H^s(0, G_\varepsilon(x_1))}^q dx_2 dx_1 \\ &\leq K^q H_1 \int_0^1 \|u(x_1, \cdot)\|_{H^s(0, G_\varepsilon(x_1))}^q dx_1 = C_1 \|u\|_{L^q(0,1;H^s(0, G_\varepsilon(x_1)))}^q, \end{aligned}$$



where  $C_2$  is independent of  $\varepsilon$ , proving (4.1).

Consequently, taking  $q = 2/s$ , since by Proposition 3.8(b) we have  $H^1(\Omega_\varepsilon) \subset L^q(0, 1; H^s(0, G_\varepsilon(x_1)))$  for  $1/2 < s \leq 1$  with constant independent of  $\varepsilon$ , it follows that

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u|^q &\leq K^q \int_0^1 \|u(x_1, \cdot)\|_{H^s(0, G_\varepsilon(x_1))}^q dx_1 \\ &= K^q \|u\|_{L^q(0, 1; H^s(0, G_\varepsilon(x_1)))}^q \leq C \|u\|_{H^1(\Omega_\varepsilon)}^q, \end{aligned}$$

proving (4.3).

Now, let us prove (4.2). Here we use that  $C^\infty(\Omega_\varepsilon)$  is dense in  $H^1(\Omega_\varepsilon)$  (see [20, Theorem 1.4.2.2]). Let  $u \in C^\infty(\Omega_\varepsilon)$  and fixed  $x_1 \in (0, 1)$ . By Fundamental Theorem of Calculus, we have

$$u(x_1, x_2) = u(x_1, 0) + \int_0^{x_2} \frac{\partial u}{\partial x_2}(x_1, s) ds.$$

Then

$$\begin{aligned} |u(x_1, x_2)|^2 &\leq 2|u(x_1, 0)|^2 + 2 \left[ \left( \int_0^{x_2} \left| \frac{\partial u}{\partial x_2}(x_1, s) \right|^2 ds \right)^{1/2} \left( \int_0^{x_2} 1^2 ds \right)^{1/2} \right]^2 \\ &\leq 2|u(x_1, 0)|^2 + 2G_\varepsilon(x_1) \int_0^{x_2} \left| \frac{\partial u}{\partial x_2}(x_1, s) \right|^2 ds. \end{aligned}$$

Consequently,

$$\begin{aligned} \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} |u(x_1, x_2)|^2 dx_2 &\leq 2 \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} |u(x_1, 0)|^2 dx_2 \\ &\quad + 2G_\varepsilon(x_1) \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} \left( \int_0^{x_2} \left| \frac{\partial u}{\partial x_2}(x_1, s) \right|^2 ds \right) dx_2 \\ &\leq 2\varepsilon H_1 |u(x_1, 0)|^2 + 2G_1 \varepsilon H_1 \int_0^{G_\varepsilon(x_1)} \left| \frac{\partial u}{\partial x_2}(x_1, x_2) \right|^2 dx_2. \end{aligned}$$

Hence, if  $\gamma(u)$  is the trace of  $u$  given by [20, Theorem 1.5.1.3], we get

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u|^2 &= \frac{1}{\varepsilon} \int_0^1 \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} |u(x_1, x_2)|^2 dx_2 dx_1 \\ &\leq 2H_1 \int_0^1 |u(x_1, 0)|^2 dx_1 + 2G_1 H_1 \int_0^1 \int_0^{G_\varepsilon(x_1)} \left| \frac{\partial u}{\partial x_2}(x_1, x_2) \right|^2 dx_2 dx_1 \\ &\leq 2H_1 \left( \|\gamma(u)\|_{L^2(0, 1)}^2 + G_1 \left\| \frac{\partial u}{\partial x_2} \right\|_{L^2(\Omega_\varepsilon)}^2 \right). \end{aligned}$$

On the other hand, if  $\Omega_0 = (0, 1) \times (0, G_0)$ , we have  $\Omega_0 \subset \Omega_\varepsilon$ , and there exists a constant  $c > 0$  such that  $\|\gamma(u)\|_{L^2(0, 1)} \leq c \|u\|_{H^s(\Omega_0)}$  for all  $1/2 < s \leq 1$ . Then, due to the previous inequality,

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u|^2 \leq 2H_1 \left( c \|u\|_{H^s(\Omega_0)}^2 + G_1 \left\| \frac{\partial u}{\partial x_2} \right\|_{L^2(\Omega_\varepsilon)}^2 \right) \leq C_1 \left( \|u\|_{H^s(\Omega_\varepsilon)}^2 + \left\| \frac{\partial u}{\partial x_2} \right\|_{L^2(\Omega_\varepsilon)}^2 \right)$$

with  $C_1$  independent of  $\varepsilon$ .  $\square$

Notice that the above theorem is important because give us a better range of estimates with the  $H^1(\Omega_\varepsilon)$  norm. However, the space may still varies with respect to the parameter  $\varepsilon$ .

Now we may study the behavior of the integrals which set the problem. We start analyzing the terms without concentration.

**Proposition 4.2.** *Let  $U \subset \mathbb{R}^2$  an open set such that  $\Omega_\varepsilon \subset U$  for all  $\varepsilon > 0$ . If  $u, \varphi \in H^1(U)$  then*

$$\int_{\Omega_\varepsilon} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 \longrightarrow \int_{\Omega} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1, \quad \text{as } \varepsilon \rightarrow 0.$$

*Proof.* Using [20, Theorem 1.4.2.1], we know that

$$C_c^\infty(\bar{U}) := \{u \in C^\infty(U); u = v|_U, \text{ com } v \in C_c^\infty(\mathbb{R}^2)\}$$

is dense in  $H^1(U)$ . Hence, we can assume  $u, \varphi \in C_c^\infty(\bar{U})$ . Then, since  $G_\varepsilon(x_1) = m(x_1) + \varepsilon g_\varepsilon(x_1)$ , where  $g_\varepsilon(x_1) = g(x_1/\varepsilon^\alpha)$  with  $0 < \alpha \leq 1$ , performing the change of variables

$$y_1 = x_1, \quad y_2 = \frac{x_2 - m(x_1)}{\varepsilon g_\varepsilon(x_1)},$$

we get

$$\begin{aligned} \int_{\Omega_\varepsilon} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 &= \int_0^1 \int_0^{m(x_1) + \varepsilon g_\varepsilon(x_1)} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 \\ &= \int_0^1 \int_0^{m(x_1)} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 + \int_0^1 \int_{m(x_1)}^{m(x_1) + \varepsilon g_\varepsilon(x_1)} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 \\ &= \int_{\Omega} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 + \varepsilon \int_0^1 \int_0^1 u(y_1, m(y_1) + y_2 \varepsilon g_\varepsilon(y_1)) \varphi(y_1, m(y_1) + y_2 \varepsilon g_\varepsilon(y_1)) g_\varepsilon(y_1) dy_2 dy_1 \\ &\rightarrow \int_{\Omega} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1, \end{aligned}$$

since  $u, \varphi \in C_c^\infty(\bar{U})$  and  $g_\varepsilon(x_1)$  is bounded by Hypothesis **H**(i) from the domain (2.2). Thus the result is valid through density properties.  $\square$

We can also prove results concerning to the behavior of the trace operator at  $\varepsilon = 0$ . Notice that, at the limit, a coefficient term appears capturing the geometry of the oscillating domain  $\Omega_\varepsilon$  and the oscillatory strip  $\theta_\varepsilon$ .

**Proposition 4.3.** *Let  $U \subset \mathbb{R}^2$  an open set such that  $\Omega_\varepsilon \subset U$  for all  $\varepsilon > 0$ . If  $u, \varphi \in H^1(U)$  then*

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 \longrightarrow \int_{\Gamma} \hat{\mu} \gamma(u) \gamma(\varphi) dS, \quad \text{as } \varepsilon \rightarrow 0,$$

where  $\gamma$  is the trace operator given by [20, Theorem 1.5.1.3] and  $\hat{\mu}$  given by (2.5).

*Proof.* Again, due to [20, Theorem 1.4.2.1], we know that

$$C_c^\infty(\bar{U}) := \{u \in C^\infty(U); u = v|_U, \text{ com } v \in C_c^\infty(\mathbb{R}^2)\}$$

is dense in  $H^1(U)$  and we can assume  $u, \varphi \in C_c^\infty(\bar{U})$ . Then, performing the change of variables

$$y_1 = x_1, \quad y_2 = \frac{x_2 - G_\varepsilon(x_1) + \varepsilon H_\varepsilon(x_1)}{\varepsilon H_\varepsilon(x_1)},$$

we get

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\theta_\varepsilon} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 &= \frac{1}{\varepsilon} \int_0^1 \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} u(x_1, x_2) \varphi(x_1, x_2) dx_2 dx_1 \\ &= \int_0^1 \int_0^1 u(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2)) \varphi(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2)) H_\varepsilon(y_1) dy_2 dy_1 \\ &= \int_0^1 \int_0^1 (u(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2)) - u(y_1, m(y_1))) \\ &\quad \varphi(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2)) H_\varepsilon(y_1) dy_2 dy_1 \end{aligned}$$

$$\begin{aligned}
& + \int_0^1 \int_0^1 u(y_1, m(y_1))(\varphi(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2)) - \varphi(y_1, m(y_1)))H_\varepsilon(y_1)dy_2dy_1 \\
& + \int_0^1 \int_0^1 u(y_1, m(y_1))\varphi(y_1, m(y_1))(H_\varepsilon(y_1) - \mu_h)dy_2dy_1 + \mu_h \int_0^1 u(y_1, m(y_1))\varphi(y_1, m(y_1))dy_1 \\
& \longrightarrow \mu_h \int_0^1 u(y_1, m(y_1))\varphi(y_1, m(y_1))\mu_h dy_1, \quad \text{as } \varepsilon \rightarrow 0,
\end{aligned}$$

since  $G_\varepsilon \rightarrow m$  when  $\varepsilon \rightarrow 0$  by Hypothesis **H**(i) from the domain (2.2). Finally, we obtain

$$\int_0^1 \mu_h u(y_1, m(y_1))\varphi(y_1, m(y_1))dy_1 = \int_\Gamma \hat{\mu}\gamma(u)\gamma(\varphi)dS$$

changing variables on the line integral, where  $\hat{\mu}$  is given by (2.5), proving the result using density and trace operator properties.  $\square$

**Remark 4.4.** *The function  $\hat{\mu}$  given by (2.5) is independent of the parametrization chosen in  $\Gamma$  and, therefore, is unique.*

We also have similar results to nonlinearities  $\Phi, f$ .

**Corollary 4.5.** *Let  $U \subset \mathbb{R}^2$  an open set such that  $\Omega_\varepsilon \subset U$  for all  $\varepsilon > 0$ . If  $u, \varphi \in H^1(U)$  and  $\Phi, f : \mathbb{R} \rightarrow \mathbb{R}$  bounded functions of class  $C^1$ , then*

$$\int_{\Omega_\varepsilon} \Phi(u(x_1, x_2))\varphi(x_1, x_2)dx_2dx_1 \rightarrow \int_\Omega \Phi(u(x_1, x_2))\varphi(x_1, x_2)dx_2dx_1$$

and

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u(x_1, x_2))\varphi(x_1, x_2)dx_2dx_1 \rightarrow \int_\Gamma \hat{\mu}\gamma(f(u))\gamma(\varphi)dS,$$

as  $\varepsilon \rightarrow 0$ , where  $\gamma$  is the trace operator given by [20, Theorem 1.5.1.3] and  $\hat{\mu} \in L^\infty(\Gamma)$  is the coefficient given by (2.5).

*Proof.* Arguing as in the proof of Propositions 4.2 and 4.3, we can assume  $u, \varphi \in C_c^\infty(\bar{U})$ . Then, using the same change of variables as before and noting that  $f$  is  $C^1$ , we have, for instance,

$$\begin{aligned}
\frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u)\varphi dx_2 dx_1 & = \frac{1}{\varepsilon} \int_0^1 \int_{G_\varepsilon(x_1) - \varepsilon H_\varepsilon(x_1)}^{G_\varepsilon(x_1)} f(u(x_1, x_2))\varphi(x_1, x_2)dx_2 dx_1 \\
& = \int_0^1 \int_0^1 f(u(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2)))\varphi(y_1, G_\varepsilon(y_1) - \varepsilon H_\varepsilon(y_1)(1 - y_2))H_\varepsilon(y_1)dy_2 dy_1 \\
& \longrightarrow \int_\Gamma \hat{\mu}\gamma(f(u))\gamma(\varphi)dS, \quad \text{as } \varepsilon \rightarrow 0.
\end{aligned}$$

The other convergence is analogous.  $\square$

The following corollaries possess similar proofs.

**Corollary 4.6.** *Let  $U \subset \mathbb{R}^2$  an open set such that  $\Omega_\varepsilon \subset U$  for all  $\varepsilon > 0$ . If  $u, \varphi, \psi \in H^1(U)$ , then*

$$\int_{\Omega_\varepsilon} u(x_1, x_2)\varphi(x_1, x_2)\psi(x_1, x_2)dx_2dx_1 \rightarrow \int_\Omega u(x_1, x_2)\varphi(x_1, x_2)\psi(x_1, x_2)dx_2dx_1$$

and

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} u(x_1, x_2)\varphi(x_1, x_2)\psi(x_1, x_2)dx_2dx_1 \rightarrow \int_\Gamma \hat{\mu}\gamma(u)\gamma(\varphi)\gamma(\psi)dS,$$

as  $\varepsilon \rightarrow 0$ , where  $\gamma$  is the trace operator given by [20, Theorem 1.5.1.3] and  $\hat{\mu} \in L^\infty(\Gamma)$  is the coefficient given by (2.5).

**Corollary 4.7.** *Let  $U \subset \mathbb{R}^2$  an open set such that  $\Omega_\varepsilon \subset U$  for all  $\varepsilon > 0$ . If  $u, \varphi, \psi \in H^1(U)$  and  $f, \Phi : \mathbb{R} \rightarrow \mathbb{R}$  bounded functions of class  $C^1$ , then*

$$\int_{\Omega_\varepsilon} \Phi(u(x_1, x_2))\varphi(x_1, x_2)\psi(x_1, x_2)dx_2dx_1 \rightarrow \int_{\Omega} \Phi(u(x_1, x_2))\varphi(x_1, x_2)\psi(x_1, x_2)dx_2dx_1$$

and

$$\frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u(x_1, x_2))\varphi(x_1, x_2)\psi(x_1, x_2)dx_2dx_1 \rightarrow \int_{\Gamma} \hat{\mu}\gamma(f(u))\gamma(\varphi)\gamma(\psi)dS,$$

as  $\varepsilon \rightarrow 0$ , where  $\gamma$  is the trace operator given by [20, Theorem 1.5.1.3] and  $\hat{\mu} \in L^\infty(\Gamma)$  is the coefficient given by (2.5).

## 5. NONLINEAR MAPS

In this section we discuss the main properties of the maps used to describe the reaction terms on the nonlinearities of the elliptic problems (2.1) and (2.4). For  $1/2 < s < 1$ , consider the Sobolev-Bochner spaces

$$X_\varepsilon = L^2(0, 1; H^s(0, G_\varepsilon(x_1))) \text{ and } X'_\varepsilon = L^2(0, 1; \{H^s(0, G_\varepsilon(x_1))\}').$$

Then define

$$\begin{aligned} F_\varepsilon : H^1(\Omega_\varepsilon) &\rightarrow X'_\varepsilon \\ u &\mapsto F_\varepsilon(u) : X_\varepsilon \rightarrow \mathbb{R} \end{aligned} \tag{5.1}$$

$$v \mapsto \langle F_\varepsilon(u), v \rangle = \int_{\Omega_\varepsilon} \Phi(u)v + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u)v,$$

where  $\Phi, f \in C^2(\mathbb{R})$  are bounded functions with bounded derivatives.

**Remark 5.1.** *Notice that the assumption  $\Phi, f \in C^2(\mathbb{R})$  bounded with bounded derivatives it is not a big restriction since we are interested in analyze  $f(u)$  when  $u$  is uniformly bounded in  $L^\infty(\Omega_\varepsilon)$ . More details can be found in [9, Remark 2.2] or [7, Remark 2.2].*

**Remark 5.2.** *Notice that  $L^2(\Omega_\varepsilon) \subset X_\varepsilon$  with constant independent of  $\varepsilon$ . Indeed, it follows from [25, Proposition 2.1] that, if  $u \in X_\varepsilon$ ,*

$$\begin{aligned} \|u\|_{L^2(\Omega_\varepsilon)}^2 &= \int_0^1 \int_0^{G_\varepsilon(x_1)} |u(x_1, x_2)|^2 dx_2 dx_1 = \int_0^1 \|u(x_1, \cdot)\|_{L^2(0, G_\varepsilon(x_1))}^2 dx_1 \\ &\leq \int_0^1 C \|u(x_1, \cdot)\|_{H^s(0, G_\varepsilon(x_1))}^2 dx_1 = C \|u\|_{X_\varepsilon}^2, \end{aligned}$$

where  $C > 0$  is independent of the domain and, furthermore, of  $\varepsilon$ .

Now we prove an analogous result to [6, Lemma 3.6] and [26, Lemma 3.1].

**Proposition 5.3.** *The function  $F_\varepsilon$  defined in (5.1) satisfies for constants independent of  $\varepsilon$ :*

(a) *there exists  $K > 0$  such that*

$$\sup_{u^\varepsilon \in H^1(\Omega_\varepsilon)} \|F_\varepsilon(u^\varepsilon)\|_{X'_\varepsilon} \leq K;$$

(b)  *$F_\varepsilon$  is globally Lipschitz continuous, that is, there exists  $L > 0$  such that*

$$\|F_\varepsilon(u_1^\varepsilon) - F_\varepsilon(u_2^\varepsilon)\|_{X'_\varepsilon} \leq L \|u_1^\varepsilon - u_2^\varepsilon\|_{H^1(\Omega_\varepsilon)}, \quad \forall u_1^\varepsilon, u_2^\varepsilon \in H^1(\Omega_\varepsilon).$$

(c)  *$F_\varepsilon$  is Frechet differentiable, with*

$$F'_\varepsilon : H^1(\Omega_\varepsilon) \rightarrow \mathcal{L}(H^1(\Omega_\varepsilon), X'_\varepsilon)$$

$$u^\varepsilon \mapsto F'_\varepsilon(u^\varepsilon) : H^1(\Omega_\varepsilon) \rightarrow X'_\varepsilon$$

$$w^\varepsilon \mapsto F'_\varepsilon(u^\varepsilon)(w^\varepsilon) : X_\varepsilon \rightarrow \mathbb{R}$$

$$v^\varepsilon \mapsto \langle F'_\varepsilon(u^\varepsilon)(w^\varepsilon), v^\varepsilon \rangle = \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon)w^\varepsilon v^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)w^\varepsilon v^\varepsilon;$$

(d) fixed  $u^\varepsilon \in H^1(\Omega_\varepsilon)$ , there is  $\bar{C} > 0$  such that

$$\|F'_\varepsilon(u^\varepsilon)(w_2^\varepsilon - w_1^\varepsilon)\|_{X'_\varepsilon} \leq \bar{C}\|w_2^\varepsilon - w_1^\varepsilon\|_{H^1(\Omega_\varepsilon)}, \quad \forall w_1^\varepsilon, w_2^\varepsilon \in H^1(\Omega_\varepsilon);$$

(e) there are  $\vartheta \in (0, 1)$  and  $M > 0$  such that

$$\|F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(v^\varepsilon)\|_{\mathcal{L}(H^1(\Omega_\varepsilon), X'_\varepsilon)} \leq M\|u^\varepsilon - v^\varepsilon\|_{X_\varepsilon}^\vartheta, \quad \forall u^\varepsilon, v^\varepsilon \in X_\varepsilon;$$

(f) there is  $k > 0$  such that

$$\|F_\varepsilon(u^\varepsilon + v^\varepsilon) - F_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)v^\varepsilon\|_{X'_\varepsilon} \leq k\|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta}, \quad \forall \delta \in (0, 1), \quad \forall u^\varepsilon, v^\varepsilon \in H^1(\Omega_\varepsilon).$$

*Proof.* (a) For  $u^\varepsilon \in H^1(\Omega_\varepsilon)$ ,

$$\|F_\varepsilon(u^\varepsilon)\|_{X'_\varepsilon} = \sup_{\|v^\varepsilon\|_{X_\varepsilon}=1} |\langle F_\varepsilon(u^\varepsilon), v^\varepsilon \rangle|.$$

Hence, if  $v^\varepsilon \in X_\varepsilon$ , it follows from Theorem 4.1 and Remark 5.2 that

$$\begin{aligned} |\langle F_\varepsilon(u^\varepsilon), v^\varepsilon \rangle| &\leq \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon)v^\varepsilon| + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon)v^\varepsilon| \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon)|^2 \right)^{\frac{1}{2}} \left( \int_{\Omega_\varepsilon} |v^\varepsilon|^2 \right)^{\frac{1}{2}} + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon)|^2 \right)^{\frac{1}{2}} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon|^2 \right)^{\frac{1}{2}} \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi(x)| \right) G_1^{1/2} \|v^\varepsilon\|_{L^2(\Omega_\varepsilon)} + \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} C \|v^\varepsilon\|_{X_\varepsilon} \leq K \|v^\varepsilon\|_{X_\varepsilon} \end{aligned}$$

Therefore

$$\sup_{u^\varepsilon \in H^1(\Omega_\varepsilon)} \|F_\varepsilon(u^\varepsilon)\|_{X'_\varepsilon} \leq K.$$

(b) Indeed, for any  $u_1^\varepsilon, u_2^\varepsilon \in H^1(\Omega_\varepsilon)$ , we have

$$\|F_\varepsilon(u_1^\varepsilon) - F_\varepsilon(u_2^\varepsilon)\|_{X'_\varepsilon} = \sup_{\|v^\varepsilon\|_{X_\varepsilon}=1} |\langle F_\varepsilon(u_1^\varepsilon), v^\varepsilon \rangle - \langle F_\varepsilon(u_2^\varepsilon), v^\varepsilon \rangle|.$$

Using Mean Value Theorem, with Theorem 4.1 and Remark 5.2 again, we get

$$\begin{aligned} |\langle F_\varepsilon(u_1^\varepsilon), v^\varepsilon \rangle - \langle F_\varepsilon(u_2^\varepsilon), v^\varepsilon \rangle| &= |\langle F_\varepsilon(u_1^\varepsilon) - F_\varepsilon(u_2^\varepsilon), v^\varepsilon \rangle| \\ &\leq \int_{\Omega_\varepsilon} |(\Phi(u_1^\varepsilon) - \Phi(u_2^\varepsilon))v^\varepsilon| + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |(f(u_1^\varepsilon) - f(u_2^\varepsilon))v^\varepsilon| \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u_1^\varepsilon) - \Phi(u_2^\varepsilon)|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |v^\varepsilon|^2 \right)^{1/2} + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u_1^\varepsilon) - f(u_2^\varepsilon)|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon|^2 \right)^{1/2} \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \|u_1^\varepsilon - u_2^\varepsilon\|_{L^2(\Omega_\varepsilon)} \|v^\varepsilon\|_{L^2(\Omega_\varepsilon)} + \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) C^2 \|u_1^\varepsilon - u_2^\varepsilon\|_{H^1(\Omega_\varepsilon)} \|v^\varepsilon\|_{X_\varepsilon} \\ &\leq L \|u_1^\varepsilon - u_2^\varepsilon\|_{H^1(\Omega_\varepsilon)} \|v^\varepsilon\|_{X_\varepsilon}. \end{aligned}$$

Thus

$$\|F_\varepsilon(u_1^\varepsilon) - F_\varepsilon(u_2^\varepsilon)\|_{X'_\varepsilon} \leq L \|u_1^\varepsilon - u_2^\varepsilon\|_{H^1(\Omega_\varepsilon)}$$

and, therefore,  $F_\varepsilon$  is globally Lipschitz with constant independent of  $\varepsilon$ .

(c) In fact, if  $u^\varepsilon, h^\varepsilon \in H^1(\Omega_\varepsilon)$  and  $v^\varepsilon \in X_\varepsilon$ , applying Mean Value Theorem,

$$\begin{aligned} |\langle F_\varepsilon(u^\varepsilon + h^\varepsilon) - F_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)h^\varepsilon, v^\varepsilon \rangle| &\leq \\ &\leq \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon + h^\varepsilon) - \Phi(u^\varepsilon) - \Phi'(u^\varepsilon)h^\varepsilon| |v^\varepsilon| + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon + h^\varepsilon) - f(u^\varepsilon) - f'(u^\varepsilon)h^\varepsilon| |v^\varepsilon| \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon + h^\varepsilon) - \Phi(u^\varepsilon) - \Phi'(u^\varepsilon)h^\varepsilon|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |v^\varepsilon|^2 \right)^{1/2} + \end{aligned}$$

$$\begin{aligned}
& + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon + h^\varepsilon) - f(u^\varepsilon) - f'(u^\varepsilon)h^\varepsilon|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon|^2 \right)^{1/2} \\
& \leq \left( \int_{\Omega_\varepsilon} |(\Phi'(\zeta^\varepsilon) - \Phi'(u^\varepsilon))h^\varepsilon|^2 \right)^{1/2} \|v^\varepsilon\|_{L^2(\Omega_\varepsilon)} + C \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |(f'(\xi^\varepsilon) - f'(u^\varepsilon))h^\varepsilon|^2 \right)^{1/2} \|v^\varepsilon\|_{X_\varepsilon} \quad (5.2)
\end{aligned}$$

where  $u^\varepsilon(x) \leq \xi^\varepsilon(x)$ ,  $\zeta^\varepsilon(x) \leq (u^\varepsilon + h^\varepsilon)(x)$ .

We will analyze the second part of (5.2). Notice that, applying Mean Value Theorem again, we get

$$|(f'(\xi^\varepsilon) - f'(u^\varepsilon))h^\varepsilon|^2 \leq |f''(\eta^\varepsilon)|^2 |\xi^\varepsilon - u^\varepsilon|^2 |h^\varepsilon|^2 \leq \left( \sup_{x \in \mathbb{R}} |f''(x)| \right) |h^\varepsilon|^4 \quad (5.3)$$

for  $\xi^\varepsilon(x) \leq \eta^\varepsilon(x) \leq u^\varepsilon(x)$ , for all  $x \in \Omega_\varepsilon$ .

On the other side,

$$|(f'(\xi^\varepsilon) - f'(u^\varepsilon))h^\varepsilon|^2 = |f'(\xi^\varepsilon) - f'(u^\varepsilon)|^2 |h^\varepsilon|^2 \leq 2 \left( \sup_{x \in \mathbb{R}} |f'(x)| \right)^2 |h^\varepsilon|^2. \quad (5.4)$$

Then putting (5.3) and (5.4) together, we have

$$|(f'(\xi^\varepsilon) - f'(u^\varepsilon))h^\varepsilon|^2 \leq K \min\{|h^\varepsilon|^2, 1\} |h^\varepsilon|^2. \quad (5.5)$$

However, for all  $\delta \in [0, 1]$ ,

$$\min\{|h^\varepsilon|^2, 1\} = \min\{|h^\varepsilon|^2, 1\}^\delta \min\{|h^\varepsilon|^2, 1\}^{1-\delta} \leq |h^\varepsilon|^{2\delta}$$

and, thus, (5.5) became

$$|(f'(\xi^\varepsilon) - f'(u^\varepsilon))h^\varepsilon|^2 \leq K_2 |h^\varepsilon|^{2(1+\delta)}, \quad \forall \delta \in [0, 1].$$

Analogously, using the properties of  $\Phi$  we may say that, for the first part of (5.2),

$$|(\Phi'(\zeta^\varepsilon) - \Phi'(u^\varepsilon))h^\varepsilon|^2 \leq K_1 |h^\varepsilon|^{2(1+\delta)}, \quad \forall \delta \in [0, 1].$$

Then it follows from (5.2) and using Remark 5.2 that

$$|\langle F'_\varepsilon(u^\varepsilon + h^\varepsilon) - F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)h^\varepsilon, v^\varepsilon \rangle| \leq K_1 \left( \int_{\Omega_\varepsilon} |h^\varepsilon|^{2(1+\delta)} \right)^{1/2} \|v^\varepsilon\|_{X_\varepsilon} + K_2 \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |h^\varepsilon|^{2(1+\delta)} \right)^{1/2} \|v^\varepsilon\|_{X_\varepsilon}.$$

Furthermore, if  $\delta \in (0, 1)$ , we can use Theorem 4.1 to get

$$\|F'_\varepsilon(u^\varepsilon + h^\varepsilon) - F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)h^\varepsilon\|_{X'_\varepsilon} \leq K_1 \left( \int_{\Omega_\varepsilon} |h^\varepsilon|^{2(1+\delta)} \right)^{1/2} + K_2 \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |h^\varepsilon|^{2(1+\delta)} \right)^{1/2} \leq C \|h^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta}.$$

Consequently,

$$\lim_{\|h^\varepsilon\|_{H^1(\Omega_\varepsilon)} \rightarrow 0} \frac{\|F'_\varepsilon(u^\varepsilon + h^\varepsilon) - F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)h^\varepsilon\|_{X'_\varepsilon}}{\|h^\varepsilon\|_{H^1(\Omega_\varepsilon)}} \leq \lim_{\|h^\varepsilon\|_{H^1(\Omega_\varepsilon)} \rightarrow 0} C^2 \|h^\varepsilon\|_{H^1(\Omega_\varepsilon)}^\delta = 0$$

and thus  $F_\varepsilon$  is Frechet differentiable.

(d) Indeed, since  $\Phi, \Phi', \Phi'', f, f', f''$  are bounded, if  $w_1^\varepsilon, w_2^\varepsilon \in H^1(\Omega_\varepsilon)$  and  $v^\varepsilon \in X_\varepsilon$ , we have by Theorem 4.1

$$\begin{aligned}
& |\langle F'_\varepsilon(u^\varepsilon)w_2^\varepsilon - F'_\varepsilon(u^\varepsilon)w_1^\varepsilon, v^\varepsilon \rangle| \leq \int_{\Omega_\varepsilon} |(\Phi'(u^\varepsilon)w_2^\varepsilon - \Phi'(u^\varepsilon)w_1^\varepsilon)v^\varepsilon| + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |(f'(u^\varepsilon)w_2^\varepsilon - f'(u^\varepsilon)w_1^\varepsilon)v^\varepsilon| \\
& \leq \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \left( \int_{\Omega_\varepsilon} |w_2^\varepsilon - w_1^\varepsilon|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |v^\varepsilon|^2 \right)^{1/2} + \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w_2^\varepsilon - w_1^\varepsilon|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon|^2 \right)^{1/2} \\
& \leq C \|w_2^\varepsilon - w_1^\varepsilon\|_{H^1(\Omega_\varepsilon)} \|v^\varepsilon\|_{X_\varepsilon}.
\end{aligned}$$

It follows that

$$\|F'_\varepsilon(u^\varepsilon)(w_2^\varepsilon - w_1^\varepsilon)\|_{X'_\varepsilon} \leq C \|w_2^\varepsilon - w_1^\varepsilon\|_{H^1(\Omega_\varepsilon)},$$

proving the result.

(e) If  $u^\varepsilon, v^\varepsilon \in H^1(\Omega_\varepsilon)$  and  $w^\varepsilon \in X_\varepsilon$ ,

$$\|F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(v^\varepsilon)\|_{\mathcal{L}(H^1(\Omega_\varepsilon), X'_\varepsilon)} = \sup_{\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}=1} \sup_{\|z^\varepsilon\|_{X_\varepsilon}=1} \langle (F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(v^\varepsilon))w^\varepsilon, z^\varepsilon \rangle.$$

Hence, if  $w^\varepsilon \in H^1(\Omega_\varepsilon)$  and  $z^\varepsilon \in X_\varepsilon$ , it follows from Theorem 4.1 and Hölder Inequality Generalized with  $3 < q < 4$  and  $4 < p < 6$  (since  $1/p + 1/q = 1/2$ ) that

$$\begin{aligned} | \langle (F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(v^\varepsilon))w^\varepsilon, z^\varepsilon \rangle | &\leq \int_{\Omega_\varepsilon} |(\Phi'(u^\varepsilon) - \Phi'(v^\varepsilon))w^\varepsilon z^\varepsilon| + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |(f'(u^\varepsilon) - f'(v^\varepsilon))w^\varepsilon z^\varepsilon| \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi'(u^\varepsilon) - \Phi'(v^\varepsilon)|^p \right)^{1/p} \left( \int_{\Omega_\varepsilon} |w^\varepsilon|^q \right)^{1/q} \left( \int_{\Omega_\varepsilon} |z^\varepsilon|^2 \right)^{1/2} + \\ &\quad + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f'(u^\varepsilon) - f'(v^\varepsilon)|^p \right)^{1/p} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w^\varepsilon|^q \right)^{1/q} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |z^\varepsilon|^2 \right)^{1/2} \\ &\leq C \left[ \left( \int_{\Omega_\varepsilon} |\Phi'(u^\varepsilon) - \Phi'(v^\varepsilon)|^p \right)^{1/p} + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f'(u^\varepsilon) - f'(v^\varepsilon)|^p \right)^{1/p} \right] \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \|z^\varepsilon\|_{X_\varepsilon}. \end{aligned}$$

Thus,

$$\|F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(v^\varepsilon)\|_{\mathcal{L}(H^1(\Omega_\varepsilon), X'_\varepsilon)} \leq \left[ \left( \int_{\Omega_\varepsilon} |\Phi'(u^\varepsilon) - \Phi'(v^\varepsilon)|^p \right)^{1/p} + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f'(u^\varepsilon) - f'(v^\varepsilon)|^p \right)^{1/p} \right].$$

Now, for all  $x \in \Omega_\varepsilon$ , we have

$$|f'(u^\varepsilon(x)) - f'(v^\varepsilon(x))| \leq 2 \left( \sup_{x \in \mathbb{R}} |f'(x)| \right).$$

On the other hand, by Mean Value Theorem,

$$|f'(u^\varepsilon(x)) - f'(v^\varepsilon(x))| \leq \left( \sup_{x \in \mathbb{R}} |f''(x)| \right) |u^\varepsilon(x) - v^\varepsilon(x)|.$$

Thus, if  $\vartheta \in (0, 1)$ ,

$$\begin{aligned} |f'(u^\varepsilon) - f'(v^\varepsilon)|^p &\leq K_1 \min\{1, |u^\varepsilon - v^\varepsilon|^p\} \\ &= K_1 \min\{1, |u^\varepsilon - v^\varepsilon|^p\}^\vartheta \min\{1, |u^\varepsilon - v^\varepsilon|^p\}^{1-\vartheta} \\ &\leq K_1 |u^\varepsilon - v^\varepsilon|^{\vartheta p}. \end{aligned}$$

Taking  $\vartheta$  such that  $\vartheta p = 2$  (ie, for some  $1/3 < \vartheta < 1/2$ ), it follows that

$$\left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f'(u^\varepsilon) - f'(v^\varepsilon)|^p \right)^{1/p} \leq \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} K_1 |u^\varepsilon - v^\varepsilon|^2 \right)^{1/p} \leq M_1 \|u^\varepsilon - v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^\vartheta.$$

In a similar way,

$$\left( \int_{\Omega_\varepsilon} |\Phi'(u^\varepsilon) - \Phi'(v^\varepsilon)|^p \right)^{1/p} \leq \left( \int_{\Omega_\varepsilon} \bar{K}_2 |u^\varepsilon - v^\varepsilon|^2 \right)^{1/p} \leq \bar{M}_2 \|u^\varepsilon - v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^\vartheta.$$

Furthermore, for some  $\vartheta \in (0, 1)$ ,

$$\|F'_\varepsilon(u^\varepsilon) - F'_\varepsilon(v^\varepsilon)\|_{\mathcal{L}(H^1(\Omega_\varepsilon), X'_\varepsilon)} \leq M \|u^\varepsilon - v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^\vartheta$$

(f) If  $u^\varepsilon, v^\varepsilon \in H^1(\Omega_\varepsilon)$ ,

$$\begin{aligned} \langle F_\varepsilon(u^\varepsilon + v^\varepsilon) - F_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)v^\varepsilon, w^\varepsilon \rangle &= \\ &= \int_{\Omega_\varepsilon} (\Phi(u^\varepsilon + v^\varepsilon) - \Phi(u^\varepsilon) - \Phi'(u^\varepsilon)v^\varepsilon)w^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f(u^\varepsilon + v^\varepsilon) - f(u^\varepsilon) - f'(u^\varepsilon)v^\varepsilon)w^\varepsilon. \end{aligned}$$

Hence, we can argue as in the proof of item (c) to obtain, for any  $\delta \in (0, 1)$ , that

$$\begin{aligned}
& \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon + v^\varepsilon) - f(u^\varepsilon) - f'(u^\varepsilon)v^\varepsilon| |w^\varepsilon| \leq \\
& \leq \left( \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon + v^\varepsilon) - \Phi(u^\varepsilon) - \Phi'(u^\varepsilon)v^\varepsilon|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |w^\varepsilon|^2 \right)^{1/2} + \\
& \quad + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon + v^\varepsilon) - f(u^\varepsilon) - f'(u^\varepsilon)v^\varepsilon|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w^\varepsilon|^2 \right)^{1/2} \\
& \leq C_1 \left( \int_{\Omega_\varepsilon} |v^\varepsilon|^{2(1+\delta)} \right)^{1/2} \|w^\varepsilon\|_{X_\varepsilon} + C_2 \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon|^{2(1+\delta)} \right)^{1/2} \|w^\varepsilon\|_{X_\varepsilon} \\
& \leq k \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta} \|w^\varepsilon\|_{X_\varepsilon}.
\end{aligned}$$

Therefore,

$$\|F_\varepsilon(u^\varepsilon + v^\varepsilon) - F_\varepsilon(u^\varepsilon) - F'_\varepsilon(u^\varepsilon)v^\varepsilon\|_{X'_\varepsilon} \leq k \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta}, \quad \forall \delta \in (0, 1),$$

which concludes the proof.  $\square$

**Remark 5.4.** *The results from Proposition 5.3 are also valid if*

$$\begin{aligned}
& F_\varepsilon : H^1(\Omega_\varepsilon) \rightarrow H^{-1}(\Omega_\varepsilon) \\
& u \mapsto F_\varepsilon(u) : H^1(\Omega_\varepsilon) \rightarrow \mathbb{R} \\
& v \mapsto \langle F_\varepsilon(u), v \rangle = \int_{\Omega_\varepsilon} \Phi(u)v + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u)v
\end{aligned}$$

or

$$\begin{aligned}
& F_\varepsilon : X_\varepsilon \rightarrow H^{-1}(\Omega_\varepsilon) \\
& u \mapsto F_\varepsilon(u) : H^1(\Omega_\varepsilon) \rightarrow \mathbb{R} \\
& v \mapsto \langle F_\varepsilon(u), v \rangle = \int_{\Omega_\varepsilon} \Phi(u)v + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u)v.
\end{aligned}$$

*This is a consequence of Proposition 3.6 and Theorem 4.1.*

## 6. UPPER AND LOWER SEMICONTINUITY OF SOLUTIONS

Our main goal in this section is to prove Theorem 2.7, passing to the limit in problem (2.1). First of all, we write equations (2.1) and (2.4) in an abstract way. Next, we combine the results from the previous sections with those ones from [7, 9] concerned with compact convergence to obtain upper and lower semicontinuity to  $\mathcal{E}_\varepsilon$  at  $\varepsilon = 0$ .

**6.1. Abstract setting and existence of solutions.** In order to write problem (2.1) in an abstract way, we consider the linear operator

$$\begin{aligned}
& A_\varepsilon : D(A_\varepsilon) \subset L^2(\Omega_\varepsilon) \rightarrow L^2(\Omega_\varepsilon) \\
& u^\varepsilon \mapsto A_\varepsilon u^\varepsilon = -\Delta u^\varepsilon + u^\varepsilon
\end{aligned}$$

with  $D(A_\varepsilon) = \{u^\varepsilon \in H^2(\Omega_\varepsilon); \frac{\partial u^\varepsilon}{\partial \nu^\varepsilon} = 0\}$ .

Let  $Z_\varepsilon^0 = L^2(\Omega_\varepsilon)$ ,  $Z_\varepsilon^1 = D(A_\varepsilon)$  and consider the scale of Hilbert spaces constructed by complex interpolation between  $Z_\varepsilon^0$  and  $Z_\varepsilon^1$ . In our context, such spaces isometrically coincide with the fractional power space of the operator  $A_\varepsilon$  (see [27, Theorem 16.1]). Such scale can be extended to negative exponents such



as  $Z_\varepsilon^{-\alpha} = (Z_\varepsilon^\alpha)'$  for  $\alpha > 0$ . Notice that  $Z_\varepsilon^{1/2} = H^1(\Omega_\varepsilon)$  and  $Z_\varepsilon^{-1/2} = (H^1(\Omega_\varepsilon))'$ . Hence, if we consider the realizations of  $A_\varepsilon$  in this scale, we have  $A_{\varepsilon, -1/2} \in \mathcal{L}(Z_\varepsilon^{1/2}, Z_\varepsilon^{-1/2})$  with

$$\langle A_{\varepsilon, -1/2} u^\varepsilon, \varphi^\varepsilon \rangle = \int_{\Omega_\varepsilon} \nabla u^\varepsilon \nabla \varphi^\varepsilon + u^\varepsilon \varphi^\varepsilon, \quad \forall \varphi^\varepsilon \in H^1(\Omega_\varepsilon).$$

With some abuse of notation we identify all different realizations of this operator writing as  $A_\varepsilon$ . Then the problem (2.1) can be rewrite as

$$A_\varepsilon u^\varepsilon = F_\varepsilon(u^\varepsilon), \quad (6.1)$$

where the map  $F_\varepsilon$  is given by

$$\begin{aligned} F_\varepsilon : H^1(\Omega_\varepsilon) &\rightarrow X'_\varepsilon \\ u^\varepsilon &\mapsto F_\varepsilon(u^\varepsilon) : L^2(0, 1; H^s(0, G_\varepsilon(x_1))) \rightarrow \mathbb{R} \\ v^\varepsilon &\mapsto \langle F_\varepsilon(u^\varepsilon), v^\varepsilon \rangle = \int_{\Omega_\varepsilon} \Phi(u^\varepsilon) v^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) v^\varepsilon, \end{aligned}$$

with  $1/2 < s < 1$ .

Thus,  $u^\varepsilon \in H^1(\Omega_\varepsilon)$  is a solution of (6.1) if, and only if,  $u^\varepsilon = A_\varepsilon^{-1} F_\varepsilon(u^\varepsilon)$ . Then,  $u^\varepsilon \in H^1(\Omega_\varepsilon)$  must be a fixed point of  $A_\varepsilon^{-1} F_\varepsilon : H^1(\Omega_\varepsilon) \rightarrow H^1(\Omega_\varepsilon)$ . The existence of such solutions follows from Schaefer Fixed Point Theorem [18, Section 9.2.2, Theorem 4].

Indeed, as we will see in Proposition 6.8, we have that the operator  $A_\varepsilon^{-1} F_\varepsilon$  is compact. Hence, to conclude the existence, we just need to prove that

$$O_\varepsilon = \{\varphi^\varepsilon \in H^1(\Omega_\varepsilon); \varphi^\varepsilon = A_\varepsilon^{-1} F_\varepsilon(\varphi^\varepsilon)\}$$

is a bounded set. Now, it is a direct consequence from Hölder Inequality and Theorem 4.1 since

$$\begin{aligned} \|\varphi^\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 &\leq \int_{\Omega_\varepsilon} |\Phi(\varphi^\varepsilon) \varphi^\varepsilon| + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(\varphi^\varepsilon) \varphi^\varepsilon| \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi(x)| \right) G_1^{1/2} \left( \int_{\Omega_\varepsilon} |\varphi^\varepsilon|^2 \right)^{1/2} + \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |\varphi^\varepsilon|^2 \right)^{1/2} \leq C \|\varphi^\varepsilon\|_{H^1(\Omega_\varepsilon)}, \end{aligned}$$

for any  $\varphi^\varepsilon \in O_\varepsilon$ .

In a similar way, we can analyze the limit problem given in (2.4). We first consider the linear operator  $A_0 \in \mathcal{L}(H^1(\Omega), H^1(\Omega)')$  with

$$\langle A_0 u, \varphi \rangle = \int_{\Omega} \nabla u \nabla \varphi + u \varphi, \quad \forall \varphi \in H^1(\Omega),$$

and then, we set the nonlinearity

$$\begin{aligned} F_0 : H^1(\Omega) &\rightarrow L^2(0, 1; \{H^s(0, m(x_1))\}') \\ u &\mapsto F_0(u) : L^2(0, 1; H^s(0, m(x_1))) \rightarrow \mathbb{R} \\ v &\mapsto \langle F_0(u), v \rangle = \int_{\Omega} \Phi(u) v + \int_{\Gamma} \hat{\mu} \gamma(f(u)) \gamma(v) dS. \end{aligned}$$

Then the limit problem (2.4) can be rewritten as

$$A_0 u = F_0(u) \quad (6.2)$$

and, with this notation,  $u \in H^1(\Omega)$  is a solution of (6.2) if, and only if,  $u = A_0^{-1} F_0(u)$ . In other words,  $u \in H^1(\Omega)$  is a fixed point of  $A_0^{-1} F_0 : H^1(\Omega) \rightarrow H^1(\Omega)$ . Again, the existence of a solution follows from Schauder's Fixed Point Theorem.

**6.2. Extension Operator.** A particular continuous linear extension operator is useful here. For the proof see [7, Proposition 4.1].

**Proposition 6.1.** *Let  $\Omega_\varepsilon$  be the family of domains defined in (2.2). Then, for each  $1 \leq p \leq \infty$ , there are  $\varepsilon_0 > 0$  and a continuous extension operator  $P_{\Omega_\varepsilon} : L^1(\Omega_\varepsilon) \rightarrow L^1(\mathbb{R}^2)$  such that, with the notation  $X(V) = L^p(V)$  or  $W^{1,p}(V)$  for an open set  $V \subset \mathbb{R}^2$ ,  $P_{\Omega_\varepsilon}$  transforms functions of  $X(\Omega_\varepsilon)$  in  $X(\mathbb{R}^2)$  with*

$$\|P_{\Omega_\varepsilon}\|_{\mathcal{L}(X(\Omega_\varepsilon), X(\mathbb{R}^2))} \leq K, \text{ for } 0 < \varepsilon < \varepsilon_0,$$

for some  $K > 0$  independent of  $\varepsilon$ .

Moreover,  $P_{\Omega_\varepsilon}$  is constructed in such way that  $P_{\Omega_\varepsilon} u \equiv 0$  outside an open set  $U$ , where  $U$  contain the closure of  $\Omega_\varepsilon$  for all  $\varepsilon > 0$ .

**Remark 6.2.** *The construction of operators  $P_{\Omega_\varepsilon}$  allows us to introduce a new family of operator  $P_{\Omega_\varepsilon, V} : X(\Omega_\varepsilon) \rightarrow X(V)$  given by  $P_{\Omega_\varepsilon, V} = R_V P_{\Omega_\varepsilon}$ , where  $R_V$  is the restriction to the open set  $V$ . Using this notation,  $P_{\Omega_\varepsilon} = P_{\Omega_\varepsilon, \mathbb{R}^2}$ . We also have  $\|P_{\Omega_\varepsilon, V}\|_{\mathcal{L}(X(\Omega_\varepsilon), X(V))} \leq C$  independent of  $\varepsilon$  (see [7, Remark 4.2]).*

The next lemma is convenient to get  $E$ -convergence results in  $\Omega_\varepsilon$  (see [7, Lemma 4.3]).

**Lemma 6.3.** *Let  $\{u^\varepsilon\}$  be a family in  $H^1(\Omega_\varepsilon)$  with  $\|u^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq M$ . Then*

- (i) *there is a subsequence of  $u^\varepsilon$ , denoted by  $u^{\varepsilon^k}$ , and  $u_0 \in H^1(\Omega)$  such that  $u^{\varepsilon^k} \xrightarrow{E} u_0$ ;*
- (ii) *there is a subsequence of  $u^\varepsilon$ , denoted by  $u^{\varepsilon^n}$ , and  $u \in H^1(U)$  such that  $P_{\Omega_{\varepsilon^n}, U} u^{\varepsilon^n} \rightharpoonup u$  in  $H^1(U)$  and  $u^{\varepsilon^n} \xrightarrow{E} u|_\Omega$ .*

**6.3. Continuity of the equilibria set.** We first show that the solutions are uniformly bounded in  $L^\infty(\Omega_\varepsilon)$ .

**Proposition 6.4.** *If  $u^\varepsilon \in H^1(\Omega_\varepsilon)$  is a solution of (6.1), then there is  $C > 0$  independent of  $\varepsilon > 0$  such that  $\|u^\varepsilon\|_{L^\infty(\Omega_\varepsilon)} \leq C$ .*

*Proof.* If  $u^\varepsilon \in H^1(\Omega_\varepsilon)$  is solution of (6.1), we have for all  $\varphi^\varepsilon \in H^1(\Omega_\varepsilon)$  that

$$\int_{\Omega_\varepsilon} \frac{\partial u^\varepsilon}{\partial x_1} \frac{\partial \varphi^\varepsilon}{\partial x_1} + \int_{\Omega_\varepsilon} \frac{\partial u^\varepsilon}{\partial x_2} \frac{\partial \varphi^\varepsilon}{\partial x_2} + \int_{\Omega_\varepsilon} u^\varepsilon \varphi^\varepsilon = \int_{\Omega_\varepsilon} \Phi(u^\varepsilon) \varphi^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) \varphi^\varepsilon.$$

Now, for  $k > 0$  take  $\varphi^\varepsilon = (u^\varepsilon - k)^+ \in H^1(\Omega_\varepsilon)$ , where

$$(u^\varepsilon - k)^+(x_1, x_2) = \begin{cases} u^\varepsilon(x_1, x_2) - k, & \text{if } (x_1, x_2) \in A_{\varepsilon, k} := \{(x_1, x_2) \in \Omega_\varepsilon; u^\varepsilon(x_1, x_2) > k\}, \\ 0, & \text{otherwise.} \end{cases}$$

Then we have that

$$\int_{\Omega_\varepsilon} \frac{\partial u^\varepsilon}{\partial x_1} \frac{\partial (u^\varepsilon - k)^+}{\partial x_1} + \int_{\Omega_\varepsilon} \frac{\partial u^\varepsilon}{\partial x_2} \frac{\partial (u^\varepsilon - k)^+}{\partial x_2} + \int_{\Omega_\varepsilon} u^\varepsilon (u^\varepsilon - k)^+ = \int_{\Omega_\varepsilon} \Phi(u^\varepsilon) (u^\varepsilon - k)^+ + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) (u^\varepsilon - k)^+.$$

Thus using Hölder Inequality, Theorem 4.1 and the definition of  $A_{\varepsilon, k}$ , we get

$$\begin{aligned} \|(u^\varepsilon - k)^+\|_{H^1(\Omega_\varepsilon)}^2 &= \int_{\Omega_\varepsilon \cap A_{\varepsilon, k}} \Phi(u^\varepsilon) (u^\varepsilon - k)^+ + \frac{1}{\varepsilon} \int_{\theta_\varepsilon \cap A_{\varepsilon, k}} f(u^\varepsilon) (u^\varepsilon - k)^+ - \int_{\Omega_\varepsilon \cap A_{\varepsilon, k}} k (u^\varepsilon - k)^+ \\ &\leq \left( \int_{\Omega_\varepsilon \cap A_{\varepsilon, k}} |\Phi(u^\varepsilon)|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon \cap A_{\varepsilon, k}} |u^\varepsilon - k|^2 \right)^{1/2} + \\ &\quad + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon \cap A_{\varepsilon, k}} |f(u^\varepsilon)|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon \cap A_{\varepsilon, k}} |u^\varepsilon - k|^2 \right)^{1/2} \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi(x)| \right) |A_{\varepsilon, k}|^{1/2} \|u^\varepsilon - k\|_{H^1(\Omega_\varepsilon)} + \left( \sup_{x \in \mathbb{R}} |f(x)| \right) \left( \frac{|\theta_\varepsilon| |A_{\varepsilon, k}|}{\varepsilon} \right)^{1/2} \|u^\varepsilon - k\|_{H^1(\Omega_\varepsilon)}. \end{aligned}$$

Since the set  $\theta_\varepsilon$  has order  $\varepsilon$ , we obtain that

$$\|u^\varepsilon - k\|_{H^1(\Omega_\varepsilon)} \leq C_1 |A_{\varepsilon, k}|^{1/2} \tag{6.3}$$

where  $C_1 > 0$  is independent of  $\varepsilon > 0$ .

Otherwise, notice that for  $p, q$  conjugates (in other words,  $1/p + 1/q = 1$ ) we have

$$\begin{aligned} \|(u^\varepsilon - k)^+\|_{L^1(A_{\varepsilon,k})} &= \int_{A_{\varepsilon,k}} (u^\varepsilon - k) \leq \left( \int_{A_{\varepsilon,k}} 1^p \right)^{1/p} \left( \int_{A_{\varepsilon,k}} (u^\varepsilon - k)^q \right)^{1/q} \\ &\leq |A_{\varepsilon,k}|^{1/p} \|(u^\varepsilon - k)\|_{L^q(\Omega_\varepsilon)}. \end{aligned} \quad (6.4)$$

From Proposition 3.8(c), we have that  $H^1(\Omega_\varepsilon) \subseteq L^q(\Omega_\varepsilon)$  for  $2 \leq q \leq 4$ . Thus, taking  $2 < q < 4$  and its conjugate  $1 < p < 2$ , we obtain from (6.3) in (6.4) that

$$\|(u^\varepsilon - k)^+\|_{L^1(A_{\varepsilon,k})} \leq C_2 |A_{\varepsilon,k}|^{1/p} \|(u^\varepsilon - k)\|_{H^1(\Omega_\varepsilon)} \leq K |A_{\varepsilon,k}|^{1/2+1/p} = K |A_{\varepsilon,k}|^{1+\delta}$$

for some  $\delta > 0$  since  $1/2 < 1/p < 1$ .

Therefore, applying [23, Lemma 5.1] we obtain  $\|u^\varepsilon\|_{L^\infty(\Omega_\varepsilon)}$  uniformly bounded, proving the result.  $\square$

We also need the following lemma.

**Lemma 6.5.** *Let  $u^\varepsilon, w^\varepsilon \in H^1(\Omega_\varepsilon)$  given by  $w^\varepsilon = A_\varepsilon^{-1} F_\varepsilon(u^\varepsilon)$ . Then  $\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq C$  for some  $C > 0$  independent of  $\varepsilon$ .*

*Proof.* Since  $w^\varepsilon = A_\varepsilon^{-1} F_\varepsilon u^\varepsilon$ , it follows that, for any  $\varphi^\varepsilon \in H^1(\Omega_\varepsilon)$ ,

$$\int_{\Omega_\varepsilon} \frac{\partial w^\varepsilon}{\partial x_1} \frac{\partial \varphi^\varepsilon}{\partial x_1} + \int_{\Omega_\varepsilon} \frac{\partial w^\varepsilon}{\partial x_2} \frac{\partial \varphi^\varepsilon}{\partial x_2} + \int_{\Omega_\varepsilon} w^\varepsilon \varphi^\varepsilon = \int_{\Omega_\varepsilon} \Phi(u^\varepsilon) \varphi^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) \varphi^\varepsilon.$$

Therefore, taking  $\varphi^\varepsilon = w^\varepsilon$ , we have from Hölder Inequality, the limitation of  $\Phi, f$  and Theorem 4.1 that

$$\begin{aligned} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon)|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |w^\varepsilon|^2 \right)^{1/2} + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon)|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w^\varepsilon|^2 \right)^{1/2} \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi(x)| \right) G_1^{1/2} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} + \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq C \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}, \end{aligned}$$

which shows the result.  $\square$

Next, we analyze the asymptotic behavior of the nonlinear terms of the problem.

**Proposition 6.6.** *Let  $w^\varepsilon, u^\varepsilon \in H^1(\Omega_\varepsilon)$  and  $w, u \in H^1(U)$  such that  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightharpoonup u$  and  $P_{\Omega_\varepsilon, U}(w^\varepsilon) \rightharpoonup w$  in  $H^1(U)$ , where  $P_{\Omega_\varepsilon, U}$  is the extension operator given by Proposition 6.1. Then*

$$\int_{\Omega_\varepsilon} \Phi(u^\varepsilon) w^\varepsilon \rightarrow \int_{\Omega} \Phi(u) w \quad \text{and} \quad \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) w^\varepsilon \rightarrow \int_{\Gamma} \hat{\mu} \gamma(f(u)) \gamma(w) dS,$$

where  $\hat{\mu}$  is given by (2.5).

*Proof.* To prove the first convergence, notice that using the Main Value Theorem we obtain

$$\begin{aligned} \left| \int_{\Omega_\varepsilon} \Phi(u^\varepsilon) w^\varepsilon - \int_{\Omega} \Phi(u) w \right| &\leq \left| \int_{\Omega_\varepsilon} \Phi(u^\varepsilon) (w^\varepsilon - w) \right| + \left| \int_{\Omega_\varepsilon} (\Phi(u^\varepsilon) - \Phi(u)) w \right| + \left| \int_{\Omega_\varepsilon} \Phi(u) w - \int_{\Omega} \Phi(u) w \right| \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon)|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |w^\varepsilon - w|^2 \right)^{1/2} + \left( \int_{\Omega_\varepsilon} |\Phi(u^\varepsilon) - \Phi(u)|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |w|^2 \right)^{1/2} + \\ &\quad + \left| \int_{\Omega_\varepsilon} \Phi(u) w - \int_{\Omega} \Phi(u) w \right| \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi(x)| \right) G_1^{1/2} \|w^\varepsilon - w\|_{L^2(\Omega_\varepsilon)} + \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \|u^\varepsilon - u\|_{X_\varepsilon} \|w\|_{L^2(\Omega_\varepsilon)} + \\ &\quad + \left| \int_{\Omega_\varepsilon} \Phi(u) w - \int_{\Omega} \Phi(u) w \right| = i + ii + iii \end{aligned}$$

Since  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightharpoonup u$  and  $P_{\Omega_\varepsilon, U}(w^\varepsilon) \rightharpoonup w$  in  $H^1(U)$ , we have that  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightarrow u$  and  $P_{\Omega_\varepsilon, U}(w^\varepsilon) \rightarrow w$  in  $L^2(U)$ . Using that  $\Phi$  and  $\Phi'$  are uniformly bounded and properties from the extension operator given by Proposition 6.1, we obtain

$$i = \left( \sup_{x \in \mathbb{R}} |\Phi(x)| \right) G_1^{1/2} \|w^\varepsilon - w\|_{L^2(\Omega_\varepsilon)} \leq \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} \|P_{\Omega_\varepsilon, U} w^\varepsilon - w\|_{L^2(U)} \rightarrow 0$$

and

$$ii = \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \|u^\varepsilon - u\|_{L^2(\Omega_\varepsilon)} \|w\|_{L^2(\Omega_\varepsilon)} \leq \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \|P_{\Omega_\varepsilon, U} u^\varepsilon - u\|_{L^2(U)} \|w\|_{L^2(U)} \rightarrow 0.$$

Since  $iii \rightarrow 0$  by Corollary 4.5, we obtain the first result.

On the other side, to prove the second convergence we have

$$\begin{aligned} & \left| \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) w^\varepsilon - \int_{\Gamma} \hat{\mu} \gamma(f(u)) \gamma(w) dS \right| \leq \left| \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon) (w^\varepsilon - w) \right| + \left| \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f(u^\varepsilon) - f(u)) w \right| + \\ & \quad + \left| \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u) w - \int_{\Gamma} \hat{\mu} \gamma(f(u)) \gamma(w) dS \right| \\ & \leq \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon)|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w^\varepsilon - w|^2 \right)^{1/2} + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u^\varepsilon) - f(u)|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w|^2 \right)^{1/2} + \\ & \quad + \left| \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u) w - \int_{\Gamma} \hat{\mu} \gamma(f(u)) \gamma(w) dS \right| \\ & \leq \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} \|w^\varepsilon - w\|_{X_\varepsilon} + \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \|u^\varepsilon - u\|_{X_\varepsilon} \|w\|_{H^1(\Omega_\varepsilon)} + \\ & \quad + \left| \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u) w - \int_{\Gamma} \hat{\mu} \gamma(f(u)) \gamma(w) dS \right| = I + II + III, \end{aligned}$$

with  $X_\varepsilon = L^2(0, 1; H^s(0, G_\varepsilon(x_1)))$  for  $1/2 < s < 1$ .

Notice that, since we are working on  $\mathbb{R}^2$ ,  $U \subset U_1 \times U_2$ , with  $U_1, U_2 \subset \mathbb{R}$  open sets,  $(0, 1) \subset U_1$  and  $(0, G_\varepsilon(x_1)) \subset U_2$  for all  $x_1 \in (0, 1)$  and  $0 < \varepsilon < \varepsilon_0$ . Therefore  $H^1(U) \subset H^1(U_1 \times U_2) \subset L^2(U_1; H^s(U_2)) =: X_U$ , where the last inclusion is compact by Proposition 3.6. Thus

$$I = \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} \|w^\varepsilon - w\|_{X_\varepsilon} \leq \left( \sup_{x \in \mathbb{R}} |f(x)| \right) H_1^{1/2} \|P_{\Omega_\varepsilon, U} w^\varepsilon - w\|_{X_U} \rightarrow 0$$

and

$$II = \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \|u^\varepsilon - u\|_{X_\varepsilon} \|w\|_{H^1(\Omega_\varepsilon)} \leq \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \|P_{\Omega_\varepsilon, U} u^\varepsilon - u\|_{X_U} \|w\|_{H^1(U)} \rightarrow 0.$$

Finally  $III \rightarrow 0$  again by Corollary 4.5 and we conclude the proof.  $\square$

**Proposition 6.7.** *Let  $u^\varepsilon, v^\varepsilon \in H^1(\Omega_\varepsilon)$  and  $u, v \in H^1(U)$  such that  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightharpoonup u$  and  $P_{\Omega_\varepsilon, U}(v^\varepsilon) \rightharpoonup v$  in  $H^1(U)$ , where  $P_{\Omega_\varepsilon, U}$  is the extension operator given by Proposition 6.1. Then, for all  $\varphi \in H^1(U)$ ,*

$$\int_{\Omega_\varepsilon} \Phi'(u^\varepsilon) v^\varepsilon \varphi \rightarrow \int_{\Omega} \Phi'(u) v \varphi \quad \text{and} \quad \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon) v^\varepsilon \varphi \rightarrow \int_{\Gamma} \hat{\mu} \gamma(f'(u)) \gamma(v) \gamma(\varphi) dS,$$

where  $\hat{\mu}$  is given by (2.5).

*Proof.* Indeed, to prove the first result we have

$$\begin{aligned} & \left| \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon) v^\varepsilon \varphi - \int_{\Omega} \Phi'(u) v \varphi \right| \leq \left| \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon) (v^\varepsilon - v) \varphi \right| + \left| \int_{\Omega_\varepsilon} (\Phi'(u^\varepsilon) - \Phi'(u)) v \varphi \right| + \\ & \quad + \left| \int_{\Omega_\varepsilon} \Phi'(u) v \varphi - \int_{\Omega} \Phi'(u) v \varphi \right| = i + ii + iii \end{aligned}$$

Remembering that  $\Phi, \Phi'$  are uniformly bounded and that  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightharpoonup u$  and  $P_{\Omega_\varepsilon, U}(v^\varepsilon) \rightharpoonup v$  in  $H^1(U)$  implies  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightarrow u$  and  $P_{\Omega_\varepsilon, U}(v^\varepsilon) \rightarrow v$  in  $L^2(U)$ , we can analyze each term on the right:

$$\begin{aligned} i &= \left| \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon)(v^\varepsilon - v)\varphi \right| \leq \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \left( \int_{\Omega_\varepsilon} |v^\varepsilon - v|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |\varphi|^2 \right)^{1/2} \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \|v^\varepsilon - v\|_{L^2(\Omega_\varepsilon)} \|\varphi\|_{L^2(\Omega_\varepsilon)} \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \|P_{\Omega_\varepsilon, U}v^\varepsilon - v\|_{L^2(U)} \|\varphi\|_{L^2(U)} \rightarrow 0 \end{aligned}$$

and using the Sobolev inclusion [27, Theorem 1.36] we have, for some  $C > 0$  independent of  $\varepsilon$  that

$$\begin{aligned} ii &= \left| \int_{\Omega_\varepsilon} (\Phi'(u^\varepsilon) - \Phi'(u))v\varphi \right| \leq \int_{\Omega_\varepsilon} |(\Phi'(u^\varepsilon) - \Phi'(u))v\varphi| \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi''(x)| \right) \left( \int_{\Omega_\varepsilon} |u^\varepsilon - u|^2 \right)^{1/2} \left( \int_{\Omega_\varepsilon} |v|^4 \right)^{1/4} \left( \int_{\Omega_\varepsilon} |\varphi|^4 \right)^{1/4} \\ &\leq \|u^\varepsilon - u\|_{L^2(\Omega_\varepsilon)} \|v\|_{L^4(\Omega_\varepsilon)} \|\varphi\|_{L^4(\Omega_\varepsilon)} \leq \|P_{\Omega_\varepsilon, U}u^\varepsilon - u\|_{L^2(U)} \|v\|_{L^4(U)} \|\varphi\|_{L^4(U)} \\ &\leq C \|P_{\Omega_\varepsilon, U}u^\varepsilon - u\|_{L^2(U)} \|v\|_{H^1(U)} \|\varphi\|_{H^1(U)} \rightarrow 0. \end{aligned}$$

For *iii*, using Corollary 4.7,

$$iii = \left| \int_{\Omega_\varepsilon} \Phi'(u)v\varphi - \int_{\Omega} \Phi'(u)v\varphi \right| \rightarrow 0,$$

proving the first result.

To prove the second convergence, we have

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)v^\varepsilon\varphi &= \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)(v^\varepsilon - v)\varphi + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f'(u^\varepsilon) - f'(u))v\varphi + \\ &+ \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u)v\varphi = I + II + III. \end{aligned}$$

Analyzing each term separately and using the definition of  $X_U$  given in the proof of Proposition 6.6:

$$\begin{aligned} I &= \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)(v^\varepsilon - v)\varphi \leq \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon - v|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |\varphi|^2 \right)^{1/2} \\ &\leq C \|v^\varepsilon - v\|_{X_\varepsilon} \|\varphi\|_{H^1(\Omega_\varepsilon)} \leq C \|P_{\Omega_\varepsilon, U}v^\varepsilon - v\|_{X_U} \|\varphi\|_{H^1(U)} \rightarrow 0. \end{aligned}$$

Since  $f'$  is  $C^1$ , applying Corollary 4.7, we get

$$III = \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u)v\varphi \rightarrow \int_{\Gamma} \hat{\mu}\gamma(f'(u))\gamma(\varphi)\gamma(\psi)dS.$$

Finally, notice that we can rewrite *II* as

$$\begin{aligned} \Psi_\varepsilon : H^1(U) &\rightarrow \mathbb{R} \\ \varphi &\mapsto \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f'(u^\varepsilon) - f'(u))v\varphi. \end{aligned}$$

It follows that  $\Psi$  is a bounded linear operator in  $H^1(U)$  since, using Theorem 4.1,

$$|\Psi_\varepsilon(\varphi)| \leq 2 \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |\varphi|^2 \right)^{1/2} \leq C \|v\|_{H^1(U)} \|\varphi\|_{H^1(U)}.$$

Besides, for all  $\varphi \in C_c^\infty(\bar{U})$ ,

$$\Psi_\varepsilon(\varphi) = \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f'(u^\varepsilon) - f'(u))v\varphi \leq \left( \sup_{x \in \mathbb{R}} |f''(x)| \right) \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |u^\varepsilon - u|^2 \right)^{1/2} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v|^2 \right)^{1/2} \|\varphi\|_\infty$$

$$\leq K \|P_{\Omega_\varepsilon, U} u^\varepsilon - u\|_{X_U} \|v\|_{H^1(U)} \|\varphi\|_{H^1(U)} \rightarrow 0$$

and then, by density, we have  $II = \Psi_\varepsilon(\varphi) \rightarrow 0$ , for all  $\varphi \in H^1(U)$ . This concludes the proof.  $\square$

For now on, consider the spaces  $H_\varepsilon = H^1(\Omega_\varepsilon)$  and  $H_0 = H^1(\Omega)$  in the context of Definition 2.1. We prove the result which guarantee the upper and lower semicontinuity of the set of solutions from (6.1) at  $\varepsilon = 0$ .

**Proposition 6.8.** *Using the notations from (6.1) and (6.2), we have that  $A_\varepsilon^{-1}F_\varepsilon \xrightarrow{CC} A_0^{-1}F_0$ .*

*Proof.* To prove the compact convergence, we verify separately each item.

(a)  $A_\varepsilon^{-1}F_\varepsilon$  is a compact operator, for each  $\varepsilon > 0$ .

Since by Proposition 3.6  $H^1(\Omega_\varepsilon) \hookrightarrow X_\varepsilon$  with compact immersion, we have  $X'_\varepsilon \hookrightarrow H^{-1}(\Omega_\varepsilon)$  compactly. Also,  $F_\varepsilon$  is a Lipschitz function by Proposition 5.3(b). Thus, we get the result from

$$H^1(\Omega_\varepsilon) \xrightarrow{F_\varepsilon} X'_\varepsilon \xrightarrow{i} H^{-1}(\Omega_\varepsilon) \xrightarrow{A_\varepsilon^{-1}} H^1(\Omega_\varepsilon).$$

(b) If  $\|u^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K$ , then  $\{A_\varepsilon^{-1}F_\varepsilon(u^\varepsilon)\}$  is  $E$ -precompact.

Let  $\{u^\varepsilon\}$  such that  $\|u^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K$ . By Lemma 6.3 we obtain a subsequence, that we still call  $u^\varepsilon$ , such that  $P_{\Omega_\varepsilon, U} u^\varepsilon \rightharpoonup u$  in  $H^1(U)$  and  $u^\varepsilon \xrightarrow{E} u|_\Omega$  for some  $u \in H^1(U)$ . Consider  $w^\varepsilon = A_\varepsilon^{-1}F_\varepsilon(u^\varepsilon)$ . By Lemma 6.5,  $\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq C$  and, thus, again by Lemma 6.3, there exists a subsequence, also called  $w^\varepsilon$ , and  $w \in H^1(U)$  such that  $P_{\Omega_\varepsilon, U} w^\varepsilon \rightharpoonup w$  in  $H^1(U)$  and  $w^\varepsilon \xrightarrow{E} w|_\Omega$ .

If we call  $u_0 = u|_\Omega$  and  $w_0 = w|_\Omega$ , we have that  $w_0 = A_0^{-1}F_0(u_0)$ . Indeed,  $w^\varepsilon \xrightarrow{E} w_0$  implies for any  $v \in H^1(U)$  that  $(w^\varepsilon, v)_{H^1(\Omega_\varepsilon)} \rightarrow (w_0, v)_{H^1(\Omega)}$ . On other hand, by Proposition 6.6 we have

$$(w^\varepsilon, v)_{H^1(\Omega_\varepsilon)} = \int_{\Omega_\varepsilon} \Phi(u^\varepsilon)v + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon)v \rightarrow \int_{\Omega} \Phi(u_0)v + \int_{\Gamma} \hat{\mu}\gamma(f(u_0))\gamma(v)dS.$$

Thus, since the limit is unique, we get

$$\langle A_0 w_0, v \rangle = (w_0, v)_{H^1(\Omega)} = \int_{\Omega} \Phi(u_0)v + \int_{\Gamma} \hat{\mu}\gamma(f(u_0))\gamma(v)dS = \langle F_0(u_0), v \rangle, \quad \forall v \in H^1(U),$$

and, therefore,  $w_0 = A_0^{-1}F_0(u_0)$ . Now, let us prove  $\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \rightarrow \|w_0\|_{H^1(\Omega)}$ , implying  $w^\varepsilon \xrightarrow{E} w_0$  by [7, Proposition 3.2]. As a matter of fact, using Proposition 6.6 again, we have

$$\begin{aligned} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 &= (w^\varepsilon, w^\varepsilon)_{H^1(\Omega)} = (A_\varepsilon^{-1}F_\varepsilon(u^\varepsilon), w^\varepsilon)_{H^1(\Omega)} = \int_{\Omega_\varepsilon} \Phi(u^\varepsilon)w^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f(u^\varepsilon)w^\varepsilon \\ &\rightarrow \int_{\Omega} \Phi(u_0)w_0 + \int_{\Gamma} \hat{\mu}\gamma(f(u_0))\gamma(w_0)dS = (A_0^{-1}F_0(u_0), w_0)_{H^1(\Omega)} = (w_0, w_0)_{H^1(\Omega)} = \|w_0\|_{H^1(\Omega)}^2. \end{aligned}$$

(c) If  $u^\varepsilon \xrightarrow{E} u$ , then  $A_\varepsilon^{-1}F_\varepsilon(u^\varepsilon) \xrightarrow{E} A_0^{-1}F_0(u)$ .

Indeed, if we assume that  $u^\varepsilon \xrightarrow{E} u$ , we get  $\|u^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq C$ , for some  $C > 0$  independent of  $\varepsilon$ . In particular, for any subsequence of  $u^\varepsilon$ , we can find another subsequence, denoting all by  $u^\varepsilon$ , such that, using the same argument of the previous item, we have  $P_{\Omega_\varepsilon, U}(u^\varepsilon) \rightharpoonup u$ , with  $u_0 = u|_\Omega$  and, for this subsequence,  $A_\varepsilon^{-1}F_\varepsilon(u^\varepsilon) \xrightarrow{E} A_0^{-1}F_0(u_0)$ . As we can prove this for any subsequence, we obtain the  $E$ -convergence of all family, that is,  $A_\varepsilon^{-1}F_\varepsilon(u^\varepsilon) \xrightarrow{E} A_0^{-1}F_0(u_0)$ .  $\square$

Finally, we can conclude the upper and lower semicontinuity of the equilibrium set at  $\varepsilon = 0$  proving Theorem 2.7. Indeed, from Proposition 6.8 and [9, Proposition 5.6], we have:

**Proposition 6.9.** *For any family  $\{u^\varepsilon\}$ ,  $u^\varepsilon \in H^1(\Omega_\varepsilon)$  solution of (6.1), there is  $u_* \in H^1(\Omega)$  solution of (6.2) and a subsequence still denoted by  $u^\varepsilon$ , such that  $u^\varepsilon \xrightarrow{E} u_*$ .*

Moreover, with the assumption that the limit solution is hyperbolic, we can get lower semicontinuity of the equilibrium set. More precisely, from Proposition 6.8 and [9, Proposition 5.7] we have

**Proposition 6.10.** *If  $u_* \in H^1(\Omega)$  solution of (6.2) is hyperbolic, then there is a sequence  $\{u_*^\varepsilon\}$ ,  $u_*^\varepsilon \in H^1(\Omega_\varepsilon)$  solution of (6.1), such that  $u_*^\varepsilon \xrightarrow{E} u_*$ .*

**Remark 6.11.** *In the case when all equilibria points of the limit equation (6.2) are hyperbolic, we have that all of them are isolated and there is only a finite number of them (see [9, Corollary 5.4 or Proposition 5.5]).*

Notice that the continuity above does not exclude the possibility that near an equilibrium point of the limiting equation may exist several different equilibrium points of the perturbed problem. We show that is possible to obtain some sort of uniqueness of the equilibrium points concluding the proof of Theorem 2.7.

First we will prove an important result about the compact convergence of  $A_\varepsilon^{-1}F'_\varepsilon(u_*^\varepsilon)$  if  $u_*^\varepsilon \in H^1(\Omega_\varepsilon)$  is a sequence of solutions from (6.1) that is  $E$ -convergent.

**Proposition 6.12.** *If  $\{u^\varepsilon\}$  is a sequence of solutions of (6.1),  $u^\varepsilon \in H^1(\Omega_\varepsilon)$ , and  $u_0 \in H^1(\Omega)$  is solution of (6.2) then  $A_\varepsilon^{-1}F'_\varepsilon(u^\varepsilon) \xrightarrow{CC} A_0^{-1}F'_0(u_0)$  whenever  $u^\varepsilon \xrightarrow{E} u_0$ .*

*Proof.* We prove by steps, as in Proposition 6.8.

(i)  $A_\varepsilon^{-1}F'_\varepsilon(u^\varepsilon)$  is compact, for each  $\varepsilon > 0$ .

Since  $H^1(\Omega_\varepsilon) \hookrightarrow X_\varepsilon$  with compact immersion by Proposition 3.6, we have

$$H^1(\Omega_\varepsilon) \xrightarrow{F'_\varepsilon(u^\varepsilon)} X'_\varepsilon \xrightarrow{i} H^{-1}(\Omega_\varepsilon) \xrightarrow{A_\varepsilon^{-1}} H^1(\Omega_\varepsilon),$$

where  $F'_\varepsilon(u^\varepsilon)$  is continuous by Proposition 5.3(d), proving the affirmation.

(ii)  $A_\varepsilon^{-1}F'_\varepsilon(u^\varepsilon)v^\varepsilon$  is  $E$ -precompact whenever  $\|v^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq C$ .

Let  $\{v^\varepsilon\}$  family in  $H^1(\Omega_\varepsilon)$  such that  $\|v^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq C$  and define  $w^\varepsilon = A_\varepsilon^{-1}F'_\varepsilon(u^\varepsilon)v^\varepsilon$ . Then for any  $\varphi^\varepsilon \in H^1(\Omega_\varepsilon)$ ,

$$\int_{\Omega_\varepsilon} \frac{\partial w^\varepsilon}{\partial x_1} \frac{\partial \varphi^\varepsilon}{\partial x_1} + \int_{\Omega_\varepsilon} \frac{\partial w^\varepsilon}{\partial x_2} \frac{\partial \varphi^\varepsilon}{\partial x_2} + \int_{\Omega_\varepsilon} w^\varepsilon \varphi^\varepsilon = \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon)v^\varepsilon \varphi^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)v^\varepsilon \varphi^\varepsilon.$$

If  $\varphi^\varepsilon = w^\varepsilon$  follows by Theorem 4.1

$$\begin{aligned} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 &= \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon)v^\varepsilon w^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)v^\varepsilon w^\varepsilon \\ &\leq \left( \sup_{x \in \mathbb{R}} |\Phi'(x)| \right) \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} + \left( \sup_{x \in \mathbb{R}} |f'(x)| \right) C^2 \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \end{aligned}$$

and, thus,  $\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K$ , for some  $K > 0$  independent of  $\varepsilon$ . Therefore, by Lemma 6.3 we obtain subsequences, also denoted by  $v^\varepsilon$ ,  $w^\varepsilon$ , and  $v, w \in H^1(U)$  such that  $P_{\Omega_\varepsilon, U}(v^\varepsilon) \rightharpoonup v$  and  $P_{\Omega_\varepsilon, U}(w^\varepsilon) \rightharpoonup w$  both in  $H^1(U)$ , with  $v^\varepsilon \xrightarrow{E} v|_\Omega$  and  $w^\varepsilon \xrightarrow{E} w|_\Omega$ .

Now if we call  $v_0 = v|_\Omega$  and  $w_0 = w|_\Omega$ , we may prove that  $w_0 = A_0^{-1}F'_0(u_0)v_0$ . Indeed, for  $\varphi \in H^1(U)$

$$(w^\varepsilon, \varphi)_{H^1(\Omega_\varepsilon)} = \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon)v^\varepsilon \varphi + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)v^\varepsilon \varphi. \quad (6.5)$$

On one hand, using Proposition 6.7, we have

$$\begin{aligned} \int_{\Omega_\varepsilon} \Phi'(u^\varepsilon)v^\varepsilon \varphi + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} f'(u^\varepsilon)v^\varepsilon \varphi &\rightarrow \int_{\Omega} \Phi'(u_0)v_0 \varphi + \int_{\Gamma} \hat{\mu} \gamma(f'(u_0)) \gamma(v_0) \gamma(\varphi) dS \\ &= (A_0^{-1}F'_0(u_0)v_0, \varphi)_{H^1(\Omega)}. \end{aligned}$$

However, since  $w^\varepsilon \xrightarrow{E} w|_\Omega$ ,

$$(w^\varepsilon, \varphi)_{H^1(\Omega_\varepsilon)} \rightarrow (w_0, \varphi)_{H^1(\Omega)}.$$

Thus  $w_0 = A_0^{-1}F'_0(u_0)v_0$ .

Finally, we show that  $w^\varepsilon \xrightarrow{E} w_0$ . By [7, Proposition 3.2], it is enough to prove  $\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \rightarrow \|w_0\|_{H^1(\Omega)}$ . But, if we take  $\varphi = w^\varepsilon$  in (6.5) we obtain arguing as in the proof of Proposition 6.8, the norm convergence.

(iii)  $A_\varepsilon^{-1}F'_\varepsilon(u^\varepsilon)v^\varepsilon \xrightarrow{E} A_0^{-1}F'_0(u_0)v_0$  se  $v^\varepsilon \xrightarrow{E} v_0$ .

To prove that  $w^\varepsilon \xrightarrow{E} w_0$  for the whole sequence it is enough to use an analogous proof of this step in Proposition 6.8.  $\square$

The following lemma is the last one that we need to conclude the uniqueness of equilibrium points near a hyperbolic limit solution.

**Lemma 6.13.** *If  $u_*^\varepsilon \in H^1(\Omega_\varepsilon)$  is a solution of (6.1) then there is  $K > 0$  such that, for all  $v^\varepsilon \in H^1(\Omega_\varepsilon)$  with  $\|v^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq 1$ , we have*

$$\|A_\varepsilon^{-1}(F_\varepsilon(u_*^\varepsilon + v^\varepsilon) - F_\varepsilon(u_*^\varepsilon) - F'_\varepsilon(u_*^\varepsilon)v^\varepsilon)\|_{H^1(\Omega_\varepsilon)} \leq K\|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta}, \quad \text{for some } \delta \in (0, 1).$$

*Proof.* Let  $w^\varepsilon = A_\varepsilon^{-1}(F_\varepsilon(u_*^\varepsilon + v^\varepsilon) - F_\varepsilon(u_*^\varepsilon) - F'_\varepsilon(u_*^\varepsilon)v^\varepsilon)$ . This implies that, for all  $\varphi^\varepsilon \in H^1(\Omega_\varepsilon)$ ,

$$\begin{aligned} \int_{\Omega_\varepsilon} \frac{\partial w^\varepsilon}{\partial x_1} \frac{\partial \varphi^\varepsilon}{\partial x_1} + \int_{\Omega_\varepsilon} \frac{\partial w^\varepsilon}{\partial x_2} \frac{\partial \varphi^\varepsilon}{\partial x_2} + \int_{\Omega_\varepsilon} w^\varepsilon \varphi^\varepsilon &= \\ &= \int_{\Omega_\varepsilon} (\Phi(u_*^\varepsilon + v^\varepsilon) - \Phi(u_*^\varepsilon) - \Phi'(u_*^\varepsilon)v^\varepsilon)\varphi^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f(u_*^\varepsilon + v^\varepsilon) - f(u_*^\varepsilon) - f'(u_*^\varepsilon)v^\varepsilon)\varphi^\varepsilon. \end{aligned}$$

Taking  $\varphi^\varepsilon = w^\varepsilon$ , the left side of the equation becomes  $\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}^2$ . For the right side, with a fixed  $1 < p < 2$  in a way that its conjugate  $q$  is  $2 < q < 4$ , follows by Theorem 4.1 that

$$\begin{aligned} &\int_{\Omega_\varepsilon} (\Phi(u_*^\varepsilon + v^\varepsilon) - \Phi(u_*^\varepsilon) - \Phi'(u_*^\varepsilon)v^\varepsilon)w^\varepsilon + \frac{1}{\varepsilon} \int_{\theta_\varepsilon} (f(u_*^\varepsilon + v^\varepsilon) - f(u_*^\varepsilon) - f'(u_*^\varepsilon)v^\varepsilon)w^\varepsilon \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u_*^\varepsilon + v^\varepsilon) - \Phi(u_*^\varepsilon) - \Phi'(u_*^\varepsilon)v^\varepsilon|^p \right)^{1/p} \left( \int_{\Omega_\varepsilon} |w^\varepsilon|^q \right)^{1/q} + \\ &\quad + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u_*^\varepsilon + v^\varepsilon) - f(u_*^\varepsilon) - f'(u_*^\varepsilon)v^\varepsilon|^p \right)^{1/p} \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |w^\varepsilon|^q \right)^{1/q} \\ &\leq \left( \int_{\Omega_\varepsilon} |\Phi(u_*^\varepsilon + v^\varepsilon) - \Phi(u_*^\varepsilon) - \Phi'(u_*^\varepsilon)v^\varepsilon|^p \right)^{1/p} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} + \\ &\quad + \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |f(u_*^\varepsilon + v^\varepsilon) - f(u_*^\varepsilon) - f'(u_*^\varepsilon)v^\varepsilon|^p \right)^{1/p} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}. \end{aligned}$$

By Proposition 5.3(f) we obtain, for  $\delta \in (0, 1)$  such that  $p(1 + \delta) = 2$  or, in other words,  $2/p = (1 + \delta)$ ,

$$\begin{aligned} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 &\leq \left( \int_{\Omega_\varepsilon} |v^\varepsilon|^2 \right)^{\frac{1}{2} \frac{2}{p}} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} + C \left( \frac{1}{\varepsilon} \int_{\theta_\varepsilon} |v^\varepsilon|^2 \right)^{\frac{1}{2} \frac{2}{p}} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \\ &\leq C^2 \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{2/p} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} = K \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta} \|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \end{aligned}$$

and, thus,

$$\|w^\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K \|v^\varepsilon\|_{H^1(\Omega_\varepsilon)}^{1+\delta}$$

proving the result.  $\square$

Now we can conclude the uniqueness of the equilibrium as  $\varepsilon$  is close to zero.

**Proposition 6.14.** *If  $u_0^*$  is a hyperbolic equilibrium of (6.2), then there exist  $\eta > 0$  and  $\varepsilon_0 > 0$  such that, for  $0 < \varepsilon < \varepsilon_0$ , there exists one, and only one,  $u_*^\varepsilon$  solution of (6.1) such that  $\|u_*^\varepsilon - E_\varepsilon u_0^*\|_{H^1(\Omega_\varepsilon)} \leq \eta$ . Furthermore  $u_*^\varepsilon \xrightarrow{E} u_0^*$ .*

*Proof.* This is a consequence of [7, Proposition 5.5] or [9, Theorem 5.8].  $\square$

Finally, we can prove the main result of this section.



*Proof of Theorem 2.7.* The item (a) follows from Theorem 6.9. On the other hand, (b) follows from Theorem 6.10 and Proposition 6.14.  $\square$

**Acknowledgements.** The first author (JMA)\* is partially supported by grants MTM2016-75465, IC-MAT Severo Ochoa project SEV-2015-0554, MINECO, Spain and Grupo de Investigación CADEDIF, UCM. The second author (AN) $^\diamond$  was supported by CNPq 141869/2013-5, Brazil. Third one (MCP) $^\dagger$  is partially supported by CNPq 303253/2017-7 and FAPESP 2017/02630-2 Brazil.

Finally, we would like to thank the anonymous referee whose comments have considerably improved the writing of the paper.

#### REFERENCES

- [1] Aragão, G. S.; Bruschi, S. M. Limit of nonlinear elliptic equations with concentrated terms and varying domains: the non uniformly Lipschitz case. *Electron. J. Differential Equations* 217 (2015).
- [2] Aragão, G. S.; Bruschi, S. M. Concentrated terms and varying domains in elliptic equations: Lipschitz case. *Math. Methods Appl. Sci.* 39 (12) (2016) 3450-3460.
- [3] Aragão, G. S.; Oliva, S. M. *Delay nonlinear boundary conditions as limit of reactions concentrating in the boundary*, J. of Diff. Equations 253 (2012), no. 9, 2573-2592.
- [4] Aragão, G. S.; Oliva, S. M. *Asymptotic behavior of a reaction-diffusion problem with delay and reaction term concentrated in the boundary*, São Paulo J. Math. Sci. 5 (2011), no. 2, 347-376.
- [5] Aragão, G. S., Pereira, A. L. and Pereira, M. C. *A nonlinear elliptic problem with terms concentrating in the boundary*, Mathematical Methods in the Applied Sciences 35 (2012), no. 9, 1110-1116.
- [6] Aragão, G. S.; Pereira, A. L.; Pereira, M. C. Attractors for a nonlinear parabolic problem with terms concentrating on the boundary. *J. Dynam. Differential Equations*, (2014), 26, 871-888.
- [7] Arrieta J M, Bruschi S M. Rapidly varying boundaries in equations with nonlinear boundary conditions. The case of a Lipschitz deformation. *Math. Models and Meth. in Appl. Sciences* 17 (10) (2007) 1555-1585.
- [8] Arrieta, J M.; Carvalho A. N. Spectral convergence and nonlinear dynamics of reaction-diffusion equations under perturbations of the domain. *Journal of Differential Equations* 199 (2004) 143-178.
- [9] Arrieta, J M.; Carvalho A. N.; Lozada-Cruz, G. Dynamics in dumbbell domains I. Continuity of the set of equilibria. *Journal of Differential Equations* 231.2 (2006): 551-597.
- [10] Arrieta, J M; Carvalho, A N; Pereira, M C; Silva, R P. Semilinear parabolic problems in thin domains with a highly oscillatory boundary. *Nonlinear Analysis: Theory, Methods & Applications* 74 (2011) 5111-5132.
- [11] Arrieta J M, Jiménez-Casas A, Rodríguez-Bernal A. Flux terms and Robin boundary conditions as limit of reactions and potentials concentrating at the boundary. *Revista Matemática Iberoamericana* (2008); 24 (1): 183-211.
- [12] Arrieta J M, Nogueira A, Pereira M C. Semilinear elliptic equations in thin regions with terms concentrating on oscillatory boundaries. <https://arxiv.org/abs/1803.09345> (2018).
- [13] Brizzi, R.; Chalot, J.P. Boundary homogenization and Neumann boundary problem. *Ric. Mat. XLVI* 2 (1997) 341-387.
- [14] Carvalho, A. N.; Piskarev, S. A general approximation scheme for attractors of abstract parabolic problems. *Numer. Funct. Anal. Optim.* 27 (2006) 785-829.
- [15] Cazenave, T; Haraux, A. An introduction to semilinear evolution equations. Vol. 13. Oxford Univ. Press on Demand, 1998.
- [16] Chandler-Wilde, S. N.; Hewett, D. P.; Moiola, A. Interpolation of Hilbert and Sobolev spaces: Quantitative estimates and counterexamples. *Mathematika* 61.2 (2015) 414-443.
- [17] Cioranescu D; Donato P. *An introduction to homogenization*. Oxford Lecture Series in Mathematics and its Applications vol.17. Oxford University Press, New York, 1999.
- [18] Evans, L. C. Partial Differential Equations. Graduate studies in mathematics. American Mathematical Society. (1998)
- [19] Gaudiello A.; Guibé, O. Homogenization of an elliptic second-order problem with  $L \log L$  data in a domain with oscillating boundary. *Communications in Contemporary Mathematics* 15 (6) (2013)
- [20] Grisvard, P. Elliptic problems in nonsmooth domains. Society for Industrial and Applied Mathematics, 2011.
- [21] Jiménez-Casas A, Rodríguez-Bernal A. Asymptotic behaviour of a parabolic problem with terms concentrated in the boundary. *Nonlinear Analysis: Theory, Methods & Applications* 71 (2009) 2377-2383.
- [22] Jiménez-Casas A, Rodríguez-Bernal A. Singular limit for a nonlinear parabolic equation with terms concentrating on the boundary. *Journal of Mathematical Analysis and Applications* 379 (2) (2011) 567-588.
- [23] Ladyzhenskaya, O A; Ural'tseva, N N. Linear and quasilinear elliptic equations. Translated from the Russian by Scripta Technica, Inc. Translation editor: Leon Ehrenpreis." (1968)
- [24] Meier, S A; Böhm, M. A note on the construction of function spaces for distributed-microstructure models with spatially varying cell geometry. *Int. J. Numer. Anal. Model* 5.5, (2008): 109-125.
- [25] Nezza, E. D.; Palatucci, G.; Valdinoci, E. Hitchhiker's guide to the fractional Sobolev spaces. *Bulletin des Sciences Mathématiques*, 136 (2012) 521-573.

- [26] Pereira, M. C. Asymptotic analysis of a semilinear elliptic equation in highly oscillating thin domains. *Zeitschrift für Angewandte Mathematik und Physik*, 67 (2016) 1-14.
- [27] Yagi, A. Abstract parabolic evolution equations and their applications. Springer Science Business Media, 2009.

JOSÉ M. ARRIETA

DPTO. DE ANÁLISIS MATEMÁTICO Y MATEMÁTICA APLICADA, FAC. CIENCIAS MATEMÁTICAS  
UNIVERSIDAD COMPLUTENSE DE MADRID, 28040 MADRID, SPAIN

AND

INSTITUTO DE CIENCIAS MATEMÁTICAS, CSIC-UAM-UC3M-UCM  
C/ NICOLÁS CABRERA 13-15, 28049 MADRID, SPAIN.

*E-mail address:* `arrieta@mat.ucm.es`

ARIADNE NOGUEIRA

DEPTO. DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA  
UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, SÃO PAULO - SP - BRAZIL

*E-mail address:* `ariadnen@ime.usp.br`

MARCONE C. PEREIRA

DEPTO. MATEMÁTICA APLICADA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA  
UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, SÃO PAULO - SP - BRAZIL

*E-mail address:* `marcone@ime.usp.br`