

The Regularity Lemma of Szemerédi for Sparse Graphs

Y. Kohayakawa

Department of Pure Mathematics and Mathematical Statistics,
University of Cambridge, 16 Mill Lane, Cambridge CB2 1SB, England

Instituto de Matemática e Estatística, Universidade de São Paulo,
Caixa Postal 20570, 01452-990 São Paulo, SP, Brazil

Abstract. In this note we present a new version of the well-known lemma of Szemerédi [17] concerning regular partitions of graphs. Our result deals with subgraphs of pseudo-random graphs, and hence may be used to partition sparse graphs that do not contain dense subgraphs.

1. Introduction

Our aim in this note is to give a simple extension of the beautiful regularity lemma of Szemerédi [17]. As is well known, a version of this lemma for bipartite graphs was one of the ingredients in Szemerédi's celebrated proof [16] of the Erdős–Turán conjecture on arithmetic progressions in dense subsets of integers. Furthermore, this bipartite version was also used by Ruzsa and Szemerédi [15] to solve an extremal problem concerning set systems.

The regularity lemma for generic graphs given in Szemerédi [17] has been used by many authors, and it has proved to play a crucial rôle in extremal graph theory. A few papers in which this lemma is important are Alon and Yuster [1], Bollobás, Erdős, Simonovits, and Szemerédi [2], Chvátal, Rödl, Szemerédi, and Trotter [4], Chvátal and Szemerédi [5],

Erdős, Frankl, and Rödl [6], Füredi [8], Rödl [12], and Rödl and Duke [14]. (We do not attempt to compile an exhaustive list here.)

More recently, generalisations of Szemerédi's lemma have been found and used by several authors. We mention Chung [3], Frankl and Rödl [7], and Prömel and Steger [11]. The novelty in these generalisations resides in that Szemerédi's result is extended to hypergraphs. Our aim here is to present a generalisation of this lemma to sparse graphs. Roughly speaking, we are concerned here in finding regular partitions of subgraphs of pseudo-random graphs. We remark that this new version of the regularity lemma is used in [9] and [10]. Moreover, we have been kindly informed that Professor Rödl [13] has also observed that this version of the regularity lemma holds.

The necessary definitions and the statement of our result, Theorem 1, is given in Section 2 below. We stress that our proof of Theorem 1, which we give in Section 3, is simply an adaptation of Szemerédi's original proof [17] to our context. Finally, we note that suitable generalisations of Theorem 1 to subhypergraphs of pseudo-random hypergraphs can be readily proved. Here, however, we restrict ourselves to the simplest case.

2. The regularity lemma for pseudo-random graphs

Let a graph $G = G^n$ of order $|G| = n$ be fixed. For $U, W \subset V = V(G)$, we write $E(U, W) = E_G(U, W)$ for the set of edges of G that have one endvertex in U and the other in W . We set $e(U, W) = e_G(U, W) = |E(U, W)|$. Now, let a partition $P_0 = (V_i)_1^\ell$ ($\ell \geq 1$) of V be fixed. For convenience, let us write $(U, W) \prec P_0$ if $U \cap W = \emptyset$ and either $\ell = 1$ or else $\ell \geq 2$ and for some $i \neq j$ ($1 \leq i, j \leq \ell$) we have $U \subset V_i, W \subset V_j$. We may now define the pseudo-random property that we shall be interested in.

Suppose $0 \leq \eta \leq 1$. We say that G is (P_0, η) -uniform if, for some $0 \leq p \leq 1$, we have that for all $U, W \subset V$ with $(U, W) \prec P_0$ and $|U|, |W| \geq \eta n$, we have

$$|e_G(U, W) - p|U||W|| \leq \eta p|U||W|. \quad (1)$$

We remark that the partition P_0 is introduced to handle the case of ℓ -partite graphs ($\ell \geq 2$). If $\ell = 1$, that is if the partition P_0 is trivial, then we are thinking of the case of ordinary graphs. In this case, we shorten the term (P_0, η) -uniform to η -uniform.

The prime example of an η -uniform graph is of course a random graph G_p . Note that for $\eta > 0$ a random graph $G_p \in \mathcal{G}(n, p)$ with $p = p(n) = C/n$ is almost surely η -uniform provided $C \geq C_0 = C_0(\eta)$, where $C_0(\eta)$ depends only on η .

Now let us go back to some definitions. Recall a graph $G = G^n$ is fixed. Let $H \subset G$ be a spanning subgraph of G . For $U, W \subset V$, let

$$d_{H,G}(U, W) = \begin{cases} e_H(U, W)/e_G(U, W) & \text{if } e_G(U, W) > 0 \\ 0 & \text{if } e_G(U, W) = 0. \end{cases}$$

Suppose $\varepsilon > 0$, $U, W \subset V$, and $U \cap W = \emptyset$. We say that the pair (U, W) is (ε, H, G) -regular, or simply ε -regular, if for all $U' \subset U$, $W' \subset W$ with $|U'| \geq \varepsilon|U|$ and $|W'| \geq \varepsilon|W|$, we have

$$|d_{H,G}(U', W') - d_{H,G}(U, W)| \leq \varepsilon.$$

We say that a partition $Q = (C_i)_0^k$ of $V = V(G)$ is (ε, k) -equitable if $|C_0| \leq \varepsilon n$, and $|C_1| = \dots = |C_k|$. Also, we say that C_0 is the *exceptional* class of Q . When the value of ε is not relevant, we refer to an (ε, k) -equitable partition as a k -equitable partition. Similarly, Q is an *equitable* partition of V if it is a k -equitable partition for some k . If P and Q are two equitable partitions of V , we say that Q *refines* P if every non-exceptional class of Q is contained in some non-exceptional class of P . If P' is an arbitrary partition of V , then Q *refines* P' if every non-exceptional class of Q is contained in some block of P' . Finally, we say that an (ε, k) -equitable partition $Q = (C_i)_0^k$ of V is (ε, H, G) -regular, or simply ε -regular, if at most $\varepsilon \binom{k}{2}$ pairs (C_i, C_j) with $1 \leq i < j \leq k$ are not ε -regular. We can now state the extension of Szemerédi's lemma to subgraphs of (P_0, η) -uniform graphs.

Theorem 1. *Let $\varepsilon > 0$ and $k_0, \ell \geq 1$ be fixed. Then there are constants $\eta = \eta(\varepsilon, k_0, \ell) > 0$ and $K_0 = K_0(\varepsilon, k_0, \ell) \geq k_0$ satisfying the following. For any (P_0, η) -uniform graph $G = G^n$, where $P_0 = (V_i)_1^\ell$ is a partition of $V = V(G)$, if $H \subset G$ is a spanning subgraph of G , then there exists an (ε, H, G) -regular (ε, k) -equitable partition of V refining P_0 with $k_0 \leq k \leq K_0$.*

3. The proof of Theorem 1

We now proceed to give the proof Theorem 1. As in [17], the following 'defect' form of the Cauchy–Schwarz inequality is used in the proof.

Lemma 2. *Let $y_1, \dots, y_v \geq 0$ be given. Suppose $0 \leq \rho = u/v < 1$, and $\sum_{1 \leq i \leq u} y_i = \alpha \rho \sum_{1 \leq i \leq v} y_i$. Then*

$$\sum_{1 \leq i \leq v} y_i^2 \geq \frac{1}{v} \left(1 + (\alpha - 1)^2 \frac{\rho}{1 - \rho} \right) \left\{ \sum_{1 \leq i \leq v} y_i \right\}^2. \quad \square$$

We now fix $G = G^n$ and put $V = V(G)$. Also, we assume that $P_0 = (V_i)_1^\ell$ is a fixed partition of V , and that G is (P_0, η) -uniform for some $0 \leq \eta \leq 1$. Moreover, we let $p = p(G)$ be as in (1).

Lemma 3. *Let $0 < \delta \leq 10^{-2}$ be fixed. Let $U, W \subset V(G)$ be such that $(U, W) \prec P_0$, and $\delta|U|, \delta|W| \geq \eta n$. If $U^* \subset U, W^* \subset W, |U^*| \geq (1 - \delta)|U|$, and $|W^*| \geq (1 - \delta)|W|$, then*

- (i) $|d_{H,G}(U^*, W^*) - d_{H,G}(U, W)| \leq 5\delta$,
- (ii) $|d_{H,G}(U^*, W^*)^2 - d_{H,G}(U, W)^2| \leq 9\delta$.

Proof. Note first that we have $\eta \leq \delta$, as $\eta n \leq \delta|U|, \delta|W| \leq \delta n$. Let U^*, W^* be as given in the lemma. We first check (i).

(i) We start by noticing that

$$\begin{aligned} d_{H,G}(U^*, W^*) &\geq \frac{e_H(U, W) - 2(1 + \eta)p\delta|U||W|}{e_G(U, W)} \\ &\geq d_{H,G}(U, W) - 2\delta \frac{1 + \eta}{1 - \eta} \geq d_{H,G}(U, W) - 3\delta. \end{aligned}$$

Moreover,

$$\begin{aligned} d_{H,G}(U^*, W^*) &\leq \frac{e_H(U, W)}{e_G(U^*, W^*)} \leq \frac{e_H(U, W)}{(1 - \eta)p|U^*||W^*|} \leq \frac{e_H(U, W)}{(1 - \eta)p(1 - \delta)^2|U||W|} \\ &\leq \frac{1 + \eta}{(1 - \eta)(1 - \delta)^2} d_{H,G}(U, W) \leq d_{H,G}(U, W) + 5\delta. \end{aligned}$$

Thus (i) follows.

(ii) The argument here is similar. First

$$\begin{aligned} d_{H,G}(U^*, W^*) &\geq \frac{(e_H(U, W) - 2(1 + \eta)p\delta|U||W|)^2}{e_G(U, W)^2} \\ &\geq d_{H,G}(U, W)^2 - \frac{4(1 + \eta)p\delta|U||W|e_H(U, W)}{e_G(U, W)(1 - \eta)p|U||W|} \\ &\geq d_{H,G}(U, W)^2 - 4\delta \frac{1 + \delta}{1 - \delta} \geq d_{H,G}(U, W)^2 - 5\delta. \end{aligned}$$

Secondly,

$$\begin{aligned} d_{H,G}(U^*, W^*)^2 &\leq \frac{e_H(U, W)^2}{e_G(U^*, W^*)^2} \\ &\leq \frac{e_H(U, W)^2}{(1 - \eta)^2 p^2 |U^*|^2 |W^*|^2} \leq \frac{e_H(U, W)^2}{(1 - \eta)^2 (1 - \delta)^4 p^2 |U||W|} \\ &\leq \left(\frac{1 + \eta}{(1 - \eta)(1 - \delta)^2} \right)^2 d_{H,G}(U, W)^2 \leq d_{H,G}(U, W)^2 + 9\delta. \end{aligned}$$

Thus (ii) follows. \square

In the sequel, a constant $0 < \varepsilon \leq 1/2$ and a spanning subgraph $H \subset G$ of G is fixed. Also, we let $P = (C_i)_0^k$ be an (ε, k) -equitable partition of $V = V(G)$ refining P_0 , where $4^k \geq \varepsilon^{-5}$. Moreover, we assume that $\eta \leq \eta_0 = \eta_0(k) = 1/k4^{k+1}$ and that $n = |G| \geq n_0 = n_0(k) = k4^{1+2k}$.

We now define an equitable partition $Q = Q(P)$ of $V = V(G)$ from P as follows. First, for each (ε, H, G) -irregular pair (C_s, C_t) of P with $1 \leq s < t \leq k$, we choose $X = X(s, t) \subset C_s$, $Y = Y(s, t) \subset C_t$ such that (i) $|X|, |Y| \geq \varepsilon|C_s| = \varepsilon|C_t|$, and (ii) $|d_{H,G}(X, Y) - d_{H,G}(C_s, C_t)| \geq \varepsilon$. For fixed $1 \leq s \leq k$, the sets $X(s, t)$ in

$$\{X = X(s, t) \subset C_s : 1 \leq t \leq k \text{ and } (C_s, C_t) \text{ is not } (\varepsilon, H, G)\text{-regular}\}$$

define a natural partition of C_s into at most 2^{k-1} blocks. Let us call such blocks the *atoms* of C_s . Now let $q = 4^k$ and set $m = \lfloor |C_s|/q \rfloor$ ($1 \leq s \leq k$). Note that $\lfloor |C_s|/m \rfloor = q$ as $|C_s| \geq n/2k \geq 2q^2$. Moreover, for later use, note that $m \geq \eta n$. We now let Q' be a partition of $V = V(G)$ refining P such that (i) C_0 is a block of Q' , (ii) all other blocks of Q' have cardinality m , except for possibly one, which has cardinality at most $m - 1$, (iii) for all $1 \leq s \leq k$, every atom $A \subset C_s$ contains exactly $\lfloor |A|/m \rfloor$ blocks of Q' , (iv) for all $1 \leq s \leq k$, the set C_s contains exactly $q = \lfloor |C_s|/m \rfloor$ blocks of Q' .

Let C'_0 be the union of the blocks of Q' that are not contained in any class C_s ($1 \leq s \leq k$), and let C'_i ($1 \leq i \leq k'$) be the remaining blocks of Q' . We are finally ready to define our equitable partition $Q = Q(P)$: we let $Q = (C'_i)_1^{k'}$.

Lemma 4. *The partition $Q = Q(P) = (C'_i)_0^{k'}$ defined from P as above is a k' -equitable partition of $V = V(G)$ refining P , where $k' = kq = k4^k$, and $|C'_0| \leq |C_0| + n4^{-k}$.*

Proof. Clearly Q refines P . Moreover, clearly $m = |C'_1| = \dots = |C'_{k'}|$ and, for all $1 \leq s \leq k$, we have $|C'_0| \leq |C_0| + k(m - 1) \leq |C_0| + k|C_s|/q \leq |C_0| + n4^{-k}$. \square

In what follows, for $1 \leq s \leq k$, we let $C_s(i)$ ($1 \leq i \leq q$) be the classes of Q' that are contained in the class C_s of P . Also, for all $1 \leq s \leq k$, we set $C_s^* = \bigcup_{1 \leq i \leq q} C_s(i)$. Now let $1 \leq s \leq k$ be fixed. Note that $|C_s^*| \geq |C_s| - (m - 1) \geq |C_s| - q^{-1}|C_s| \geq |C_s|(1 - q^{-1})$. As $q^{-1} \leq 10^{-2}$ and $q^{-1}|C_s| \geq m \geq \eta n$, by Lemma 3 we have, for all $1 \leq s < t \leq k$,

$$|d_{H,G}(C_s^*, C_t^*) - d_{H,G}(C_s, C_t)| \leq 5q^{-1} \quad (2)$$

and

$$|d_{H,G}(C_s^*, C_t^*)^2 - d_{H,G}(C_s, C_t)^2| \leq 9q^{-1} \quad (3)$$

Similarly to [17], we define the *index* $\text{ind}(R)$ of an equitable partition $R = (V_i)_0^r$ of $V = V(G)$ to be

$$\text{ind}(R) = \frac{2}{r^2} \sum_{1 \leq i < j \leq r} d_{H,G}(V_i, V_j)^2.$$

Note that trivially $0 \leq \text{ind}(R) < 1$. Our aim now is to show that, for $Q = Q(P)$ defined as above, we have $\text{ind}(Q) \geq \text{ind}(P) + \varepsilon^5/100$. We start with the following lemma.

Lemma 5. *Suppose $1 \leq s < t \leq k$. Then*

$$\frac{1}{q^2} \sum_{i,j=1}^q d_{H,G}(C_s(i), C_t(j))^2 \geq d_{H,G}(C_s, C_t)^2 - \frac{\varepsilon^5}{100}.$$

Proof. By the (P_0, η) -uniformity of G and the fact that $(C_s, C_t) \prec P_0$, we have

$$\begin{aligned} \frac{1}{q^2} \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} d_{H,G}(C_s(i), C_t(j)) &= \frac{1}{q^2} \sum_{i,j} \frac{e_H(C_s(i), C_t(j))}{e_G(C_s(i), C_t(j))} \\ &\geq \sum_{i,j} \frac{e_H(C_s(i), C_t(j))}{(1+\eta)q^2 p |C_s(i)||C_t(j)|} = \frac{e_H(C_s^*, C_t^*)}{(1+\eta)p |C_s^*||C_t^*|} \\ &\geq \frac{1-\eta}{1+\eta} d_{H,G}(C_s^*, C_t^*) \geq d_{H,G}(C_s^*, C_t^*) - 2\eta. \end{aligned}$$

Thus, by the Cauchy–Schwarz inequality, we have

$$\frac{1}{q^2} \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} d_{H,G}(C_s(i), C_t(j))^2 \geq d_{H,G}(C_s^*, C_t^*)^2 - 4\eta.$$

Furthermore, by (3), we have $d_{H,G}(C_s^*, C_t^*)^2 \geq d_{H,G}(C_s, C_t)^2 - 9q^{-1}$. Since $9q^{-1} + 4\eta \leq \varepsilon^5/100$, the lemma follows. \square

The inequality in Lemma 5 may be improved if (C_s, C_t) is an (ε, H, G) -irregular pair, as shows the following result.

Lemma 6. *Let $1 \leq s < t \leq k$ be such that (C_s, C_t) is not (ε, H, G) -regular. Then*

$$\frac{1}{q^2} \sum_{i,j=1}^q d_{H,G}(C_s(i), C_t(j))^2 \geq d_{H,G}(C_s, C_t)^2 + \frac{\varepsilon^4}{40} - \frac{\varepsilon^5}{100}.$$

Proof. Let $X = X(s, t) \subset C_s$, $Y = Y(s, t) \subset C_t$ be as in the definition of Q . Let $X^* \subset X$ be the maximal subset of X that is the union of blocks of Q , and similarly for $Y^* \subset Y$. Without loss of generality, we may assume that $X^* = \bigcup_{1 \leq i \leq q_s} C_s(i)$, and $Y^* = \bigcup_{1 \leq j \leq q_t} C_t(j)$. Note that $|X^*| \geq |X| - 2^{k-1}(m-1) \geq |X|(1 - 2^{k-1}m/|X|) \geq |X|(1 - 2^{k-1}/q\varepsilon) = |X|(1 - 1/\varepsilon 2^{k+1})$, and similarly $|Y^*| \geq |Y|(1 - 1/\varepsilon 2^{k+1})$. However, we have $1/\varepsilon 2^{k+1} \leq 10^{-2}$ and $|X|/\varepsilon 2^{k+1}$, $|Y|/\varepsilon 2^{k+1} \geq \eta n$. Thus, by Lemma 3, we have $|d_{H,G}(X^*, Y^*) - d_{H,G}(X, Y)| \leq 5/\varepsilon 2^{k+1}$. Moreover, by (2), we have $|d_{H,G}(C_s^*, C_t^*) - d_{H,G}(C_s, C_t)| \leq 5q^{-1}$. Since $|d_{H,G}(X, Y) - d_{H,G}(C_s, C_t)| \geq \varepsilon$ and $5q^{-1} + 5/\varepsilon 2^{k+1} \leq \varepsilon/2$, we have

$$|d_{H,G}(X^*, Y^*) - d_{H,G}(C_s^*, C_t^*)| \geq \varepsilon/2. \quad (4)$$

For later reference, let us note that $q_s m = |X^*| \geq |X| - 2^{k-1}m \geq \varepsilon|C_s| - 2^{k-1}m \geq \varepsilon q m - 2^{k-1}m$, and hence $q_s \geq \varepsilon q - 2^{k-1} \geq \varepsilon q/2$. Similarly, we have $q_t \geq \varepsilon q/2$. Let us now set $y_{ij} = d_{H,G}(C_s(i), C_t(j))$ for $i, j = 1, \dots, q$. In the proof of Lemma 5 we checked that

$$\sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij} \geq \frac{1-\eta}{1+\eta} q^2 d_{H,G}(C_s^*, C_t^*) \geq (1-2\eta) q^2 d_{H,G}(C_s^*, C_t^*).$$

Similarly, one has $\sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij} \leq (1+3\eta) q^2 d_{H,G}(C_s^*, C_t^*)$, $\sum_{1 \leq i \leq q_s} \sum_{1 \leq j \leq q_t} y_{ij} \geq (1-2\eta) q_s q_t d_{H,G}(X^*, Y^*)$, and $\sum_{1 \leq i \leq q_s} \sum_{1 \leq j \leq q_t} y_{ij} \leq (1+3\eta) q_s q_t d_{H,G}(X^*, Y^*)$. Let us set $\rho = q_s q_t / q^2 \geq \varepsilon^2/4$, and $d_{s,t}^* = d_{H,G}(C_s^*, C_t^*)$. We now note that by (4) we either have

$$\begin{aligned} \sum_{1 \leq i \leq q_s} \sum_{1 \leq j \leq q_t} y_{ij} &\geq \frac{1-2\eta}{1+3\eta} \cdot \frac{q_s q_t}{q^2} \left(1 + \frac{\varepsilon}{2(d_{s,t}^*)^2}\right) \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij} \\ &\geq \rho \left(1 + \frac{\varepsilon}{3(d_{s,t}^*)^2}\right) \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij}, \end{aligned}$$

or else

$$\begin{aligned} \sum_{1 \leq i \leq q_s} \sum_{1 \leq j \leq q_t} y_{ij} &\leq \frac{1+3\eta}{1-2\eta} \cdot \frac{q_s q_t}{q^2} \left(1 - \frac{\varepsilon}{2(d_{s,t}^*)^2}\right) \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij} \\ &\leq \rho \left(1 - \frac{\varepsilon}{3(d_{s,t}^*)^2}\right) \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij}. \end{aligned}$$

We may now apply Lemma 2 to conclude that

$$\begin{aligned} \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij}^2 &\geq \frac{1}{q^2} \left(1 + \frac{\varepsilon^2}{9(d_{s,t}^*)^2} \cdot \frac{\rho}{1-\rho}\right) \left\{ \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} y_{ij} \right\}^2 \\ &\geq \frac{1}{q^2} \left(1 + \frac{\varepsilon^2 \rho}{9(d_{s,t}^*)^2}\right) \{q^2(1-2\eta)d_{s,t}^*\}^2 \\ &\geq q^2(1-4\eta) \left((d_{s,t}^*)^2 + \frac{\varepsilon^2 \rho}{9} \right) \geq q^2 \left((d_{s,t}^*)^2 + \frac{\varepsilon^2 \rho}{10} - 4\eta \right). \end{aligned}$$

Therefore

$$\begin{aligned} \frac{1}{q^2} \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} d_{H,G}(C_s(i), C_t(j))^2 &\geq d_{H,G}(C_s^*, C_t^*)^2 + \frac{\varepsilon^2 \rho}{10} - 4\eta \\ &\geq d_{H,G}(C_s, C_t)^2 + \frac{\varepsilon^4}{40} - (9\eta^{-1} + 4\eta) \geq d_{H,G}(C_s, C_t)^2 + \frac{\varepsilon^4}{40} - \frac{\varepsilon^5}{100}, \end{aligned}$$

as required. \square

We are now ready to prove the main lemma needed in the proof of Theorem 1.

Lemma 7. *Suppose $k \geq 1$ and $0 < \varepsilon \leq 1/2$ are such that $4^k \geq 1800\varepsilon^{-5}$. Let $G = G^n$ be a (P_0, η) -uniform graph of order $n \geq n_0 = n_0(k) = k4^{2k+1}$, where $P_0 = (V_i)_1^\ell$ is a partition of $V = V(G)$, and assume that $\eta \leq \eta_0 = \eta_0(k) = 1/k4^{k+1}$. Let $H \subset G$ be a spanning subgraph of G . If $P = (C_i)_0^k$ is an (ε, H, G) -irregular (ε, k) -equitable partition of $V = V(G)$ refining P_0 , then there is a k' -equitable partition $Q = (C'_i)_0^{k'}$ of V such that (i) Q refines P , (ii) $k' = k4^k$, (iii) $|C'_0| \leq |C_0| + n4^{-k}$, and (iv) $\text{ind}(Q) \geq \text{ind}(P) + \varepsilon^5/100$.*

Proof. Let P be as in the lemma. We show that the k' -equitable partition $Q = (C'_i)_0^{k'}$ defined from P as above satisfies (i)–(iv). In view of Lemma 4, it only remains to check (iv). By Lemmas 5 and 6, we have

$$\begin{aligned} \text{ind}(Q) &= \frac{2}{(kq)^2} \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} d_{H,G}(C'_i, C'_j)^2 \\ &\geq \frac{2}{k^2} \sum_{1 \leq s < t \leq k} \frac{1}{q^2} \sum_{1 \leq i \leq q} \sum_{1 \leq j \leq q} d_{H,G}(C_s(i), C_t(j))^2 \\ &\geq \frac{2}{k^2} \left\{ \sum_{1 \leq s < t \leq k} \left(d_{H,G}(C_s, C_t)^2 - \frac{\varepsilon^5}{100} \right) + \varepsilon \binom{k}{2} \frac{\varepsilon^4}{40} \right\} \\ &\geq \text{ind}(P) - \frac{\varepsilon^5}{100} + \frac{\varepsilon^5}{50} \geq \text{ind}(P) + \frac{\varepsilon^5}{100}. \end{aligned}$$

This completes the proof of the lemma. \square

Proof of Theorem 1. Let $\varepsilon > 0$, $k_0 \geq 1$, and $\ell \geq 1$ be given. We may assume that $\varepsilon \leq 1/2$. Pick $s \geq 1$ such that $4^{s/4\ell} \geq 1800\varepsilon^{-5}$, $s \geq \max\{2k_0, 3\ell/\varepsilon\}$, and $\varepsilon 4^{s-1} \geq 1$. Let $f(0) = s$, and put inductively $f(t) = f(t-1)4^{f(t-1)}$ ($t \geq 1$). Let $t_0 = \lfloor 100\varepsilon^{-5} \rfloor$ and set $N = \max\{n_0(f(t)) : 0 \leq t \leq t_0\} = f(t_0)4^{2f(t_0)+1}$, $K_0 = \max\{6\ell/\varepsilon, N\}$, and $\eta = \eta(\varepsilon, k_0, \ell) =$

$\min\{\eta_0(f(t)) : 0 \leq t \leq t_0\} = 1/4f(t_0 + 1) > 0$. We claim that η and K_0 as defined above will do.

To prove our claim, let $G = G^n$ be a fixed (P_0, η) -uniform graph, where $P_0 = (V_i)_1^\ell$ is a partition of $V = V(G)$. Furthermore, let $H \subset G$ be a spanning subgraph of G . Note that we may clearly assume that $n \geq K_0$. Suppose $t \geq 0$. Let us say that an equitable partition $P^{(t)} = (C_i)_0^k$ of V is t -valid if (i) $P^{(t)}$ refines P_0 , (ii) $s/4\ell \leq k \leq f(t)$, (iii) $\text{ind}\{P^{(t)}\} \geq t\varepsilon^5/100$, and (iv) $|C_0| \leq \varepsilon n(1 - 2^{-(t+1)})$. We now verify that a 0-valid partition $P^{(0)}$ of V does exist. Let $m = \lceil n/s \rceil$, and let Q be a partition of V with all blocks of cardinality m , except for possibly one, which has cardinality at most $m - 1$, and moreover such that each V_i ($1 \leq i \leq \ell$) contains $\lfloor |V_i|/m \rfloor$ blocks of Q . Grouping at most ℓ blocks of Q into a single block C_0 , we arrive at an equitable partition $P^{(0)} = (C_i)_0^k$ of V that is 0-valid. Indeed, (i) is clear, and to check (ii) note that $k \leq n/m \leq s = f(0)$, and that there is $1 \leq i \leq \ell$ such that $|V_i| \geq n/\ell$, and so $k \geq \lfloor |V_i|/m \rfloor \geq \lfloor (n/\ell)/\lceil n/s \rceil \rfloor \geq (1/2)\{(n/\ell)/(2n/s)\} = s/4\ell$. Also, (iii) is trivial and (iv) does follow, since $|C_0| < \ell m \leq \ell \lceil n\varepsilon/3\ell \rceil \leq n\varepsilon/2$ as $n \geq K_0 \geq 6\ell/\varepsilon$.

Now note that if there is a t -valid partition $P^{(t)}$ of V , then $t \leq t_0 = \lfloor 100\varepsilon^{-5} \rfloor$, since $\text{ind}\{P^{(t)}\} \leq 1$. Suppose t is the maximal integer for which there is a t -valid partition $P^{(t)}$ of V . We claim that $P^{(t)}$ is (ε, H, G) -regular. Suppose to the contrary that $P^{(t)}$ is not (ε, H, G) -regular. Then simply note that Lemma 7 gives a $(t + 1)$ -valid equitable partition $P^{(t+1)} = Q = Q(P^{(t)})$, contradicting the maximality of t . This completes the proof of the theorem. \square

Acknowledgements. The author was supported by Churchill College, Cambridge, where he was a Junior Research Fellow while this work was being done.

References

- [1] Alon, N., Yuster, R., Almost H -factors in dense graphs, *Graphs and Combinatorics* **8** (1992), 95–102.
- [2] Bollobás, B., Erdős, P., Simonovits, M., Szemerédi, E., Extremal graphs without large forbidden subgraphs, in *Advances in Graph Theory* (Bollobás, B., ed.), Annals of Discrete Mathematics 3, North-Holland, Amsterdam 1978, pp. 29–41.
- [3] Chung, F.R.K., Regularity lemmas for hypergraphs and quasi-randomness, *Random Structures and Algorithms* **2** (1991), 241–252.
- [4] Chvátal, V., Rödl, V., Szemerédi, E., Trotter, W.T., The Ramsey number of a graph with

- bounded degree, *J. Combinatorial Theory (B)* **34** (1983), 239–243.
- [5] Chvátal, V., Szemerédi, E., On the Erdős–Stone theorem, *J. London Math. Soc.* **23** (1981), 207–214.
 - [6] Erdős, P., Frankl, P., Rödl, V., The asymptotic number of graphs not containing a fixed subgraph and a problem for hypergraphs having no exponent, *Graphs and Combinatorics* **2** (1986), 113–121.
 - [7] Frankl, P., Rödl, V., The uniformity lemma for hypergraphs, *Graphs and Combinatorics* **8** (1992), 309–312.
 - [8] Füredi, Z., The maximum number of edges in a minimal graph of diameter 2, *J. Graph Theory* **16** (1992), 81–98.
 - [9] Haxell, P.E., Kohayakawa, Y., On an anti-Ramsey property of Ramanujan graphs, in preparation.
 - [10] Haxell, P.E., Kohayakawa, Y., Ramsey and Turán type results for cycles in pseudo-random graphs, in preparation.
 - [11] Prömel, H.-J., Steger, A., Excluding induced subgraphs III: a general asymptotic, *Random Structures and Algorithms* **3** (1992), 19–31.
 - [12] Rödl, V., On universality of graphs with uniformly distributed edges, *Discrete Math.* **59** (1986), 125–143.
 - [13] Rödl, V., Personal communication, August, 1993.
 - [14] Rödl, V., Duke, R.A., On graphs with small subgraphs of large chromatic number, *Graphs and Combinatorics* **1** (1985), 91–96.
 - [15] Ruzsa, I.Z., Szemerédi, E., Triple systems with no six points carrying three triangles, *Coll. Math. Soc. János Bolyai* **18** (1978), 939–945.
 - [16] Szemerédi, E., On a set containing no k elements in arithmetic progression, *Acta Arithmetica* **XXVII** (1975), 199–245.
 - [17] Szemerédi, E., Regular partitions of graphs, in *Problèmes Combinatoires et Théorie des Graphes*, Proc. Colloque Inter. CNRS (Bermond, J.-C., Fournier, J.-C., Las Vergnas, M., Sotteau, D., eds), CNRS, Paris 1978, pp. 399–401.