

1

2

## DENSITY OF IDENTIFYING CODES OF HEXAGONAL GRIDS WITH FINITE NUMBER OF ROWS

3

4

RUDINI M. SAMPAIO<sup>1</sup>, GABRIEL A. G. SOBRAL<sup>2</sup> AND YOSHIKO  
WAKABAYASHI<sup>2</sup>

5

**Abstract.** In a graph  $G$ , a set  $C \subseteq V(G)$  is an identifying code if, for all vertices  $v$  in  $G$ , the sets  $N[v] \cap C$  are all nonempty and pairwise distinct, where  $N[v]$  denotes the closed neighbourhood of  $v$ . We focus on the minimum density of identifying codes of infinite hexagonal grids  $H_k$  with  $k$  rows, denoted by  $d^*(H_k)$ , and present optimal solutions for  $k \leq 5$ . Using the discharging method, we also prove a lower bound in terms of maximum degree for the minimum-density identifying codes of well-behaved infinite graphs. We prove that  $d^*(H_2) = 9/20$ ,  $d^*(H_3) = 6/13 \approx 0.4615$ ,  $d^*(H_4) = 7/16 = 0.4375$  and  $d^*(H_5) = 11/25 = 0.44$ . We also prove that  $H_2$  has a unique periodic identifying code with minimum density.

**Keywords:** identifying code, hexagonal grid, minimum density

**Mathematics Subject Classification.** 94B65, 68R10, 90C27, 05C69

6

### INTRODUCTION

7

8

9

10

The concept of *identifying code* (*idcode*, for short), was introduced in 1998 by Karpovsky *et al.* [27] to identify a faulty processor in a multiprocessor system. The vertices of an idcode correspond to special processors (the monitors) that are able to check themselves and their neighbours to identify a faulty processor.

---

Dates will be set by the publisher.

<sup>1</sup> Depto de Computação Universidade Federal do Ceará (UFC) – Fortaleza, CE – Brazil;  
e-mail: rudini@dc.ufc.br

<sup>2</sup> Instituto de Matemática e Estatística – Universidade de São Paulo – São Paulo, SP – Brazil;  
e-mail: gagsobral@ime.usp.br & yw@ime.usp.br

11 Problems on idcodes have been studied on finite and infinite graphs, being of  
 12 great interest both from theoretical as well as practical viewpoint. Particular in-  
 13 terest has been dedicated to grids as many processor networks have a grid topology  
 14 (see [34,35]). Among these, we mention the square grid  $\mathcal{G}_S$ , the triangular grid  $\mathcal{G}_T$   
 15 and the king grid  $\mathcal{G}_K$ , shown in Figure 1.

16 One fundamental problem on idcodes is that of finding idcodes of minimum  
 17 density. The density captures the proportion of vertices in the code with respect  
 18 to the whole graph. For finite graphs, Cohen *et al.* [7] proved that deciding the  
 19 existence of an idcode of size at most  $k$  in a graph is an NP-complete problem. On  
 20 infinite graphs, studies on minimum-density idcodes have considered grids with  
 21 infinite or with a finite number of rows (see [1–6, 9, 10, 12–14, 16–21, 24, 25, 27, 28]).  
 22 For an updated bibliography covering this topic and related ones, the reader is  
 23 referred to Jean [22].




FIGURE 1. Partial representation of infinite square, triangular and king grids, and the corresponding minimum-density idcodes

24 We denote by  $d^*(G)$  the minimum density of an idcode of a graph  $G$ . For the  
 25 infinite grids mentioned previously, it is known that  $d^*(\mathcal{G}_S) = 7/20$  [1],  $d^*(\mathcal{G}_T) =$   
 26  $1/4$  [27] and  $d^*(\mathcal{G}_K) = 2/9$  [5]. When these grids have a finite number  $k$  of rows,  
 27 idcodes of minimum density are known for  $k \leq 6$ , and for larger  $k$  only lower and  
 28 upper bounds have been found.

29 In this work we focus on infinite graphs, specially the hexagonal grids (see  
 30 Figure 2). We denote these grids by  $\mathcal{G}_H$  when the number of rows is infinite, and  
 31 by  $H_k$  when the number of rows is a positive integer  $k$ . For  $\mathcal{G}_H$ , new lower and  
 32 upper bounds have been proved in the last years. Just to mention the more recent  
 33 ones: in 2009, Cranston and Yu [9] proved a lower bound of  $12/29 \approx 0.4138$ , and  
 34 in 2013, Cuckierman and Yu [10] improved the lower bound to  $5/12 \approx 0.4166$ . In  
 35 2014, Stolee [33] presented a computer-assisted framework showing that  $d^*(\mathcal{G}_H) \geq$   
 36  $23/55 \approx 0.4181$ . As for upper bounds, in 2000, Cohen *et al.* [6] constructed two  
 37 idcodes of  $\mathcal{G}_H$  with density  $3/7 \approx 0.4285$ . Other idcodes with the same density  
 38 have also been reported in the literature. Recently, breaking the long-standing  
 39 bound of  $3/7$ , Salo and Törmä [29] showed that  $d^*(\mathcal{G}_H) \leq 53/126 \approx 0.4206$ . They  
 40 found a periodic idcode using a computer-assisted proof that uses automata theory  
 41 and Karp's minimum mean cycle algorithm. No results on lower or upper bounds  
 42 have appeared in the literature for  $d^*(H_k)$ .

43 We prove that idcodes of well-behaved infinite graphs with maximum degree  $\Delta$   
 44 have density at least  $2/(\Delta + 2)$ . This result and another one on infinite graphs  
 45 with maximum degree 3 imply that  $d^*(H_k) \geq 2/5$  for all  $k \geq 2$ , and that idcodes  
 46 of  $H_k$  that do not induce trivial components have density at least  $3/7$ . We prove  
 47 that  $d^*(H_2) = 9/20$ , and exhibit an idcode with this minimum density, which  
 48 we show to be unique. We also mention how we proved that  $d^*(H_3) = 6/13$ ,  
 49  $d^*(H_4) = 7/16$  and  $d^*(H_5) = 11/25$ , using computer-assisted tools.

50 In Section 1 we define the concepts used in this paper and establish the notation.  
 51 We also present a density result on the infinite 3-regular tree, to show that this  
 52 graph is not so well-behaved as the hexagonal grids, a fact (to be made precise) that  
 53 has caused an erroneous proof in the literature on a related concept called locating-  
 54 dominating set (and perhaps on other closed concepts as well). These preliminary  
 55 comments help understanding the property (named SG) that we require from the  
 56 infinite graphs to guarantee that some density proof techniques work. In Sections 2  
 57 and 3, we define SG-property and prove results on the discharging method and the  
 58 mentioned lower bound. In Section 4 we show a minimum-density idcode for  $H_2$ ,  
 59 and prove that it is unique. Section 5 contains results on minimum-density idcodes  
 60 for  $H_k$ ,  $k \in \{3, 4, 5\}$ .

61 A preliminary version of this work (an extended abstract) appeared in [30].  
 62 This work contains additional novel results and a simplified and complete proof of  
 63 Theorem 4.6.

## 64 1. DEFINITIONS, NOTATION, AND THE INFINITE 3-REGULAR TREE

65 The *hexagonal grid*, denoted by  $\mathcal{G}_H$ , is an infinite graph with vertex set  $V =$   
 66  $\mathbb{Z} \times \mathbb{Z}$  and edge set  $E = \{\{u, v\} : u = (i, j), u - v \in \{(\pm 1, 0), (0, (-1)^{i+j+1})\}\}$ .  
 67 See Figure 2. The *hexagonal grid with  $k$  rows*,  $k \geq 2$ , denoted by  $H_k$ , is a graph  
 isomorphic to the subgraph of  $\mathcal{G}_H$  induced by the vertex set  $\mathbb{Z} \times \{1, \dots, k\}$ .




FIGURE 2. Hexagonal grid  $\mathcal{G}_H$

68 Let  $G$  be a connected graph. If  $v$  is a vertex of  $G$ , and  $r$  is a natural number,  
 69 then  $N_r(v)$  denotes the set of vertices of  $v$  at distance at most  $r$  from  $v$ , and  
 70  $N_r[v] = N_r(v) \cup \{v\}$  denotes the *closed neighbourhood* of  $v$ . When  $r = 1$ , we  
 71 omit the subscript  $r$  and simply write  $N(v)$  and  $N[v]$ . Given  $C \subseteq V(G)$ , let  
 72  $C[v] = N[v] \cap C$ . An *idcode* of  $G$  is a set  $C \subseteq V(G)$  such that  $C[v] \neq \emptyset$  for every  
 73 vertex  $v$  of  $G$ , and  $C[v] \neq C[w]$  for any pair of distinct vertices  $v, w$  of  $G$ . Thus,  
 74

75 if a graph  $G$  has two distinct vertices  $v$  and  $w$  such that  $N[v] = N[w]$ , then  $G$  has  
 76 no idcode. Such vertices are called *twins*. Clearly, a graph has an idcode if and  
 77 only if it is twin-free. If  $C$  is an idcode, we say that  $C[v]$  is the *identifier* of  $v$ .

78 We are interested in minimum-density idcodes of countably infinite connected  
 79 graphs of bounded degree. For such a graph  $G$ , the *density* of a subset  $C \subseteq V(G)$ ,  
 80 denoted by  $d(C, G)$ , is defined as follows.

$$d(C, G) = \inf \{d_w(C, G) : w \in V(G)\},$$

81 where

$$d_w(C, G) = \limsup_{r \rightarrow \infty} \frac{|C \cap N_r[w]|}{|N_r[w]|}.$$

82 The *minimum density* of an idcode of a graph  $G$ , denoted by  $d^*(G)$ , is defined as

$$d^*(G) = \inf \{d(C, G) : C \text{ is an idcode of } G\}.$$

83 Notice that we use  $\inf$  (infimum) in the definition of  $d(C, G)$ , instead of  $\min$  (min-  
 84 imum), since the greatest lower bound does not always belong to the set. This  
 85 definition (with  $\inf$ ) is also given by Jiang [24] to study densities of idcodes of  
 86  $S_k$  (a topic to be mentioned in Section 5). Slater [31] defines density of locating-  
 87 dominating sets (a notion similar to idcode) with  $\min$ , but the definition of density  
 88  $d(C, G)$  makes sense for any set  $C$ . In the proof of Lemma 1.1 we show an example  
 89 of an infinite graph  $G$  for which  $d_w(C, G) > 0$  for all  $w \in V(G)$ , but  $d(C, G) = 0$ .

90 This definition of subset density given above has not always been used. In some  
 91 papers, such as [10–13, 23], the density  $d(C, G)$  was simply defined as  $d_w(C, G)$   
 92 where  $w$  is an “*arbitrary vertex*”. This contains an implicit assumption that  
 93  $d_w(C, G) = d_v(C, G)$  for any two vertices  $w, v$  of  $G$ , which is not always true  
 94 as we show in Lemma 1.1. In most of these papers, this problem in the density  
 95 definition did not lead to erroneous results, since the graphs considered were well-  
 96 behaved grids, all of them satisfy an important condition (named SG-property in  
 97 the next section) which guarantees that  $d_w(C, G) = d_v(C, G)$  for any two vertices  
 98  $w, v$  of  $G$  (see Lemma 2.1). However, some papers contain erroneous statements,  
 99 as we will see in Theorem 1.2.

100 **Lemma 1.1.** *There are infinite bounded degree graphs  $G$  with subsets  $C \subset V(G)$   
 101 for which there are distinct vertices  $w, v$  such that  $d_w(C, G) \neq d_v(C, G)$ .*

102 *Proof.* Let us consider the infinite 3-regular tree  $T$ , obtained from two infinite  
 103 binary trees  $T_1$  and  $T_2$  with roots  $r_1$  and  $r_2$ , respectively, by adding the edge  $r_1 r_2$ .  
 104 We exhibit two examples of sets  $C \subset V(T)$  and vertices  $w, v$  of  $V(T)$  for which  
 105  $d_w(C, T) \neq d_v(C, T)$ .

106 As a first example, consider  $C = V(T_2)$ . Let  $w$  be a vertex of  $T_1$  that is a  
 107 neighbour of  $r_1$ . Then  $d_w(C, T) = 1/6$ . (More generally, If  $w$  is at distance  $d$   
 108 from  $r_1$ , we have that  $d_w(C, T) = 2^{-d}/3$ .) Let  $v = r_2$ . Then,  $d_v(C, T) = 2/3$ .  
 109 (Note that here  $d(C, T) = 0$ .)

110 As a second example, let  $C$  be the set consisting of all vertices of  $T_2$  together  
 111 with all vertices of  $T_1$  whose distance to  $r_1$  is even ( $r_1$  included). In this case,  $C$

112 is an idcode of  $T$ . Let  $w$  (resp.  $v$ ) a vertex in  $T_1$  (resp.  $T_2$ ) that is at distance  $d$   
 113 from  $r_1$  (resp.  $r_2$ ). It is not difficult to check that  $d_w(C, T)$  converges to  $2/3$  and  
 114  $d_v(C, T)$  converges to 1 when  $d$  tends to  $\infty$ .  $\square$

115 Even considering the correct definition of subset density  $d(C, G)$ , some papers  
 116 calculate it in an informal way, covering the entire graph with periodic patterns  
 117 and assuming that the density of  $C$  will be the density of the pattern. As an  
 118 example, consider the infinite 3-regular tree  $T$ , used in the proof of Lemma 1.1,  
 119 which is obtained from two infinite binary trees with roots  $r_1$  and  $r_2$  and the edge  
 120  $r_1 r_2$ . Consider that  $T$  is rooted at  $r_1$ . Let  $C$  be the set of vertices in  $T$  whose  
 121 distance to  $r_1$  is even ( $r_1$  included). Then, the vertices of  $T$  can be covered by the  
 122 pattern (a matching) formed by a vertex and its leftmost child (being one in  $C$   
 123 and the other not in  $C$ ), whose density is  $1/2$ . Also, by ignoring  $r_2$ , the vertices  
 124 of  $T$  can be covered by the pattern (a cherry) formed by a vertex in  $C$  and its two  
 125 children not in  $C$ , whose density is  $1/3$ . Finally, by ignoring  $r_1$ , the vertices of  $T$   
 126 can also be covered by the pattern (a cherry) formed by a vertex not in  $C$  and its  
 127 two children in  $C$ , whose density is  $2/3$ .

128 Thus, considering three distinct periodic patterns, this method gives three dif-  
 129 ferent values as the density of  $d(C, T)$ , indicating that such a method should not  
 130 be used in any graph. We will elaborate more on this in what follows, calling  
 131 attention to a property that the infinite graph should satisfy for this method to  
 132 work (see Lemma 2.1). Unfortunately, this informal way to calculate the density of  
 133 sets on infinite graphs led to some erroneous results in the literature. We will not  
 134 present here the proof (based on the definition we have given) that  $d(C, T) = 2/3$ ,  
 135 as it is not so short, but the reader may verify this.

136 The next theorem shows that one of the first results on locating-dominating  
 137 sets is wrong. We say that a set  $C \subseteq V(G)$  is a *locating-dominating set* (lds)  
 138 of  $G$  if  $C[v] \neq \emptyset$ , for every  $v \notin C$ , and  $C[v] \neq C[w]$ , for any two distinct vertices  
 139  $v, w \notin C$ . Notice that every identifying code is also a locating-dominating set  
 140 (the difference is that a locating-dominating set  $C$  only cares about the vertices  
 141 outside  $C$ ). In 2002, Slater [31] stated that “the density of any locating-dominating  
 142 set of a countably infinite  $d$ -regular graph is at least  $2/(d + 3)$ ”. We present an  
 143 lds of the infinite 3-regular tree whose density is at most  $5/16 = 0.3125$  (a value  
 144 smaller than  $2/(3 + 3)$ ), which is a counterexample to the result stated by Slater.

145 **Theorem 1.2.** *The minimum density of a locating-dominating set of the infinite*  
 146 *3-regular tree is at most  $5/16 = 0.3125$ .*

147 *Proof.* Let  $T$  be the infinite 3-regular tree with root  $R$ , and let layer  $L_i$  be the set  
 148 of vertices of  $T$  at a distance  $i$  from the root  $R$ . Thus,  $V(T) = \bigcup_{i \geq 0} L_i$ ,  $L_0 = \{R\}$ ,  
 149 and  $|L_i| = 3 \cdot 2^{i-1}$ , for  $i \geq 1$ . Thus, for  $i \geq 5$ ,  $|L_i|$  is a multiple of 16, and is  
 150 composed of 3 groups with  $2^{i-1}$  vertices.

151 To construct a set  $C \subset V(T)$  which we shall prove to be an lds of  $T$ , we label  
 152 first the vertices of  $T$ , and then we define which vertices belong to  $C$ . The labelling  
 153 procedure is the following.

- 154 (a) We assign label 1 to all vertices in  $L_0 \cup L_1 \cup \dots \cup L_4$ .  
 155 (b) We label the vertices of  $L_5$  as follows. We consider that  $L_5$  is composed of  
 156 3 consecutive groups of 16 vertices (each of these groups are the leaves of  
 157 the subtree of height 4 rooted at one of the children of root  $R$ ). We label  
 158 identically these groups of 16 vertices, according to the following pattern:

159 -----  
 160 1 2 3 5 1 2 3 5 2 3 5 5 3 4 5 5  
 161 -----

- 162 (c) Once the vertices in  $L_i$ ,  $i \geq 5$ , have been labelled, we label the vertices  
 163 in  $L_{i+1}$ . For that, we define for each vertex with label  $j$  (in  $L_i$ ) which  
 164 are the labels  $k, l$  of its children (in  $L_{i+1}$ ), writing  $j \rightarrow \{k, l\}$ . We let  
 165  $1 \rightarrow \{3, 4\}$ ,  $2 \rightarrow \{3, 3\}$ ,  $3 \rightarrow \{1, 5\}$ ,  $4 \rightarrow \{5, 5\}$  and  $5 \rightarrow \{2, 5\}$ .  
 166 Representing this in a tree-like structure, we have:

167           1               2               3               4               5  
 168           / \            / \            / \            / \            / \  
 169           3 4            3 3            1 5            5 5            2 5

170

171 Now that  $V(T)$  is labelled, let

$$C := \{v \in V(T) : v \text{ has label 1 or 2}\}.$$

172 Consider a group, say  $H$ , of 16 vertices in  $L_5$ , and let  $x_j$  be the number of  
 173 vertices in  $H$  with label  $j$ . Then,  $x_1 = 2$ ,  $x_2 = 3$ ,  $x_3 = 4$ ,  $x_4 = 1$ ,  $x_5 = 6$ ; or in a  
 174 condensed form,  $x(H) = (2, 3, 4, 1, 6)$ .

175 Now, let  $\text{chld}(H)$  be the group (in  $L_6$ ) formed by the children of the vertices  
 176 in  $H$ . Let now  $x'_j$  be the number of vertices with label  $j$  in  $\text{chld}(H)$ . Then,  
 177  $x'_1 = x_3 = 4 = 2x_1$ ,  $x'_2 = x_5 = 6 = 2x_2$ ,  $x'_3 = x_1 + 2x_2 = 8 = 2x_3$ ,  $x'_4 =$   
 178  $x_1 = 2 = 2x_4$ , and  $x'_5 = x_3 + 2x_4 + x_5 = 12 = 2x_5$ . That is,  $x'_j = 2x_j$  for  
 179  $j \in \{1, 2, \dots, 5\}$ , and therefore,  $x(\text{chld}(H)) = 2x(H)$ . Since, at each layer  $L_i$ ,  
 180  $i \geq 5$ , there are 3 groups with  $2^{i-1}$  vertices, and each such group  $G$  (by the  
 181 labelling rule) gives rise to a (children) group with  $x(\text{chld}(G)) = 2x(G)$ , in each  
 182 new layer the proportion of vertices with labels 1 or 2 (those in  $C$ ) is exactly the  
 183 proportion that holds in layer  $L_5$ . We have  $|C \cap L_5| = 15$  and  $|L_5| = 48$ . Thus,  
 184  $|C \cap L_5|/|L_5| = 15/48 = 5/16$ . Since  $|L_{i+1}| = 2|L_i|$  and  $|C \cap L_{i+1}| = 2|C \cap L_i|$ ,  
 185 for each layer  $L_i$  the ratio  $|C \cap L_i|/|L_i| = 5/16$  holds for all  $i \geq 5$ . Only for the  
 186 initial layers  $L_i$ ,  $0 \leq i \leq 4$ , we have  $|C \cap L_i|/|L_i| = 1$ . Thus, the density  $d_R(C, T)$   
 187 is precisely

$$d_R(C, T) = \limsup_{r \rightarrow \infty} \frac{|C \cap N_r[R]|}{|N_r[R]|} = \limsup_{h \rightarrow \infty} \frac{|C \cap T_h(R)|}{|T_h(R)|} = 5/16,$$

188 where  $T_h(R)$  is the subtree of  $T$  with height  $h$  rooted at  $R$ . Since  $d(C, T) =$   
 189  $\inf\{d_w(C, T) : w \in V(T)\}$ , we conclude that  $d(C, T) \leq 5/16 = 0.3125$ .

190 It remains to prove that  $C$  is an lds of  $T$ . For that, it suffices to check that  
 191 the vertices with labels 3, 4, 5 have distinct neighbourhood in  $C$ . The reader may

192 check that a vertex with label 3 is identified by its parent and one child (with  
 193 label 1); a vertex with label 4 is identified solely by its parent (which has label 1);  
 194 and a vertex with label 5, if it belongs to  $L_5$ , then is identified by its parent and  
 195 one child (with label 2), and if it belongs to layer  $L_i$ ,  $i \geq 6$ , then it is identified  
 196 solely by one child (the one with label 2). This concludes our proof that  $C$  is an  
 197 lds of  $T$  with  $d(C, T) \leq 5/16 = 0.3125$ .  $\square$

198 We understand that the erroneous proof of Theorem 2 stated in [31] happened  
 199 because the infinite graph under consideration does not satisfy a property that  
 200 would allow the application of the method that was used. The author used a  
 201 measure called *share*  $\gamma(v, C)$ , that is an application of the *discharging method* (to  
 202 be discussed in the next section) to obtain a lower bound proof for the density of  
 203 a set, say  $C$ .

204 Roughly speaking, the share method works as follows: each vertex of  $C$  starts  
 205 with charge  $q > 0$  and each vertex outside  $C$  starts with charge 0. For any vertex  
 206  $c \in C$  and  $u \in N[c]$ , the vertex  $c$  sends charge  $1/|C[u]|$  to  $u$  (this includes the  
 207 case in which  $u = c$ ). At the end of this procedure, all vertices outside  $C$  will  
 208 have charge exactly 1 and every vertex  $c \in C$  will have charge  $q + 1 - sh(c)$ ,  
 209 where  $sh(c) = \sum_{u \in N[c]} 1/|C[u]|$  is the total charge sent by  $c$ . The idea is that, if  
 210  $sh(c) \leq q$  for every  $c \in C$ , all vertices in  $G$  will have charge at least 1. Then, if  $G$   
 211 is finite,

$$1 \cdot |V(G)| \leq \sum_{c \in C} sh(c) \leq q \cdot |C|, \quad \text{and hence } d(C, G) = \frac{|C|}{|V(G)|} \geq \frac{1}{q}.$$

212 Now, let  $G$  be an infinite connected graph and let  $v$  be a vertex of  $G$ . To guarantee  
 213 charge at least 1 at every vertex in  $N_{r-1}[v]$ , it suffices to consider the vertices in  
 214  $C \cap N_r[v]$ . Thus,

$$1 \cdot |N_{r-1}[v]| \leq \sum_{c \in C \cap N_r[v]} sh(c) \leq q \cdot |C \cap N_r[v]|,$$

215 which implies that

$$d_v(C, G) = \limsup_{r \rightarrow \infty} \frac{|C \cap N_r[v]|}{|N_r[v]|} \geq \frac{1}{q} \cdot \limsup_{r \rightarrow \infty} \frac{|N_{r-1}[v]|}{|N_r[v]|}.$$

216 As we can see, the *share method* of [31] will work if  $\limsup_{r \rightarrow \infty} |N_{r-1}[v]|/|N_r[v]| = 1$ ,  
 217 which is a consequence (Lemma 2.1(a) with  $t = -1$ ) of our SG-property, defined  
 218 in the next section.

## 219 2. THE USE OF DISCHARGING METHOD TO PROVE LOWER BOUNDS 220 FOR THE DENSITY OF IDCODS

221 The discharging method is a proof technique in combinatorics, first used in  
 222 graph theory, that has now been used in many different contexts, such as in graph

223 colouring, decomposition, embedding, geometric and structural problems. For  
 224 a guide on the use of the this method to prove results on colouring and other  
 225 structural properties of graphs see [8].

226 To prove results on a graph  $G$ , this method involves two phases: *charging*  
 227 and *discharging*. In the charging phase, we assign charges (a rational number)  
 228 to certain structures of  $G$  using a *charging rule*, which describes the value of the  
 229 charge and the structures of  $G$  which will receive the charge. These structures  
 230 may be vertices, edges, faces (if  $G$  is planar), etc. In the discharging phase, we  
 231 re-assign the charges using the *discharging rules*, which describe the structures  
 232 that will send and/or receive charge from other vertices. The discharging must  
 233 preserve the total charge that was assigned in the charging phase.

234 Both the charging and discharging rules are designed to guarantee that, after  
 235 these phases some information on the charges of certain vertices/edges will help  
 236 us prove some property of the graph. In some applications, the initial charges or  
 237 the discharging rules may take into consideration the degree of the vertices.

238 The discharging method has been one of the main tools to prove lower bounds  
 239 for density of idcodes. Theorem 2.2, proved in this section, tells how this method  
 240 can be used to obtain density results in infinite graphs, once these graphs satisfy  
 241 certain properties. Before that, we define *SG-property* and present a general result  
 242 (Lemma 2.1) that is related to this property and is used in Theorem 2.2 and  
 243 Lemma 2.3. (Here, the mnemonic SG stands for “slow growth”, the concept we  
 244 want to emphasize.)

245 **Definition 1.** We say that a graph  $G$  satisfies the SG-property if  $G$  is connected  
 246 and has a vertex  $s$  such that  $\lim_{r \rightarrow \infty} \frac{|N_{r+1}[s]|}{|N_r[s]|} = 1$ .

247 Notice that, since  $N_r[s] \subseteq N_{r+1}[s]$ , then  $\lim_{r \rightarrow \infty} \frac{|N_{r+1}[s]|}{|N_r[s]|} = 1$  if and only if  
 248  $\limsup_{r \rightarrow \infty} \frac{|N_{r+1}[s]|}{|N_r[s]|} = 1$ . Also notice that the integer  $t$  in the item (a) of the  
 249 following lemma may be negative.

250 **Lemma 2.1.** *Let  $G$  be an infinite connected graph satisfying the SG-property, and*  
 251 *let  $s \in V(G)$  be such that  $\lim_{r \rightarrow \infty} \frac{|N_{r+1}[s]|}{|N_r[s]|} = 1$ . Then the following hold.*

- 252 (a) *For every vertex  $v$  and integer  $t$ , we have  $\lim_{r \rightarrow \infty} \frac{|N_{r+t}[v]|}{|N_r[v]|} = 1$ .*  
 253 (b) *For every vertex  $v$  and  $C \subseteq V(G)$ , we have  $d_v(C, G) = d_s(C, G)$ . Thus the*  
 254 *density of  $C$  is  $d(C, G) = d_w(C, G)$ , where  $w$  is an arbitrary vertex of  $G$ .*

255 *Proof.* To simplify notation, let  $n_k[w] = |N_k[w]|$  for any positive integer  $k$  and  
 256 vertex  $w$ . For the vertex  $s$  stated in the lemma, and any integer  $t > 0$ , we have

$$\lim_{r \rightarrow \infty} \frac{n_{r+t}[s]}{n_r[s]} = \lim_{r \rightarrow \infty} \left( \frac{n_{r+t}[s]}{n_{r+t-1}[s]} \cdot \frac{n_{r+t-1}[s]}{n_{r+t-2}[s]} \cdots \frac{n_{r+2}[s]}{n_{r+1}[s]} \cdot \frac{n_{r+1}[s]}{n_r[s]} \right) = 1. \quad (1)$$

257 It is immediate that  $\lim_{r \rightarrow \infty} \frac{n_{r+t}[s]}{n_r[s]} = 1$  also holds when  $t$  is negative (as long  
 258 as  $r + t \geq 0$ ). Now, to prove (a), consider a vertex  $v$  and let  $d := \text{dist}(v, s)$ . First,  
 259 we prove that (for  $r \geq d$ )

$$N_{r-d}[s] \subseteq N_r[v] \subseteq N_{r+d}[s]. \quad (2)$$

260 To prove the first inclusion, take a vertex  $y$  in  $N_{r-d}[s]$ . Thus,  $\text{dist}(y, s) \leq r - d$ .  
 261 Since  $\text{dist}(y, v) \leq \text{dist}(y, s) + \text{dist}(s, v)$ , it follows that  $\text{dist}(y, v) \leq r$ , and therefore,  
 262  $y \in N_r[v]$ . The proof of the second inclusion is analogous: take  $y \in N_r[v]$ , which  
 263 means that  $\text{dist}(y, v) \leq r$ . Since  $\text{dist}(y, s) \leq \text{dist}(y, v) + \text{dist}(v, s)$ , we have that  
 264  $\text{dist}(y, s) \leq r + d$ , and therefore,  $y \in N_{r+d}[s]$ . From (2), we have that

$$N_{r+1-d}[s] \subseteq N_{r+1}[v] \subseteq N_{r+1+d}[s]. \quad (3)$$

265 Combining (3) and (2), we have

$$\frac{n_{r+1-d}[s]}{n_{r+d}[s]} \leq \frac{n_{r+1}[v]}{n_r[v]} \leq \frac{n_{r+1+d}[s]}{n_{r-d}[s]}. \quad (4)$$

266 Since (1) holds for every integer  $t$  (see the observation in the paragraph follow-  
 267 ing (1)), it follows that the limit of the fraction on the left (resp. right) side of (4)  
 268 when  $r$  tends to  $\infty$  is 1, and therefore,

$$\lim_{r \rightarrow \infty} \frac{n_{r+1}[v]}{n_r[v]} = 1. \quad (5)$$

269 From (5), we may conclude that (1) holds when  $s$  is replaced by  $v$ , and this  
 270 completes the proof of statement (a).

271 Now, let us prove (b). For that, we first note that, from (2) we have that

$$\frac{n_{r-d}[s]}{n_r[s]} \leq \frac{n_r[v]}{n_r[s]} \leq \frac{n_{r+d}[s]}{n_r[s]}. \quad (6)$$

272 Since the limit of the fraction on the left (resp. right) when  $r$  tends to  $\infty$  is 1,  
 273 it follows that

$$\lim_{r \rightarrow \infty} \frac{n_r[v]}{n_r[s]} = 1. \quad (7)$$

274 By definition, we have that

$$d_v(C, G) = \limsup_{r \rightarrow \infty} \frac{|C \cap N_r[v]|}{n_r[v]}. \quad (8)$$

275 From (2), we obtain

$$C \cap N_{r-d}[s] \subseteq C \cap N_r[v] \subseteq C \cap N_{r+d}[s].$$

276 Thus,

$$\limsup_{r \rightarrow \infty} \frac{|C \cap N_{r-d}[s]|}{n_r[v]} \leq d_v(C, G) \leq \limsup_{r \rightarrow \infty} \frac{|C \cap N_{r+d}[s]|}{n_r[v]}. \quad (9)$$

277 The lower (resp. upper) bound of  $d_v(C, G)$  given by (9) is precisely  $d_s(C, G)$ .  
 278 Indeed, for the lower bound, using (8),(1) and (7), we have

$$\limsup_{r \rightarrow \infty} \frac{|C \cap N_{r-d}[s]|}{n_r[v]} = \limsup_{r \rightarrow \infty} \left( \frac{|C \cap N_{r-d}[s]|}{n_{r-d}[s]} \cdot \frac{n_{r-d}[s]}{n_r[s]} \cdot \frac{n_r[s]}{n_r[v]} \right) = d_s(C, G).$$

279 For the upper bound, the proof follows similarly. Thus,  $d_v(C, G) = d_s(C, G)$ ,  
 280 and hence  $d(C, G) = d_w(C, G)$ , where  $w$  is an arbitrary vertex in  $G$ .  $\square$

281 The SG-property is very important for the forthcoming proofs on the minimum  
 282 density based on the discharging method. Lemma 2.1 guarantees that if a con-  
 283 nected graph  $G$  has this property, then the density of a vertex set  $C$  in  $G$  may be  
 284 calculated by considering  $d_v(C, G)$  for an arbitrary vertex  $v$ .

285 It is not difficult to see that the infinite hexagonal grids ( $\mathcal{G}_H$  and  $H_k$ ), as well  
 286 as the grids mentioned in the introduction (square, triangular, king), and many  
 287 others have the SG-property. In particular, for the grid  $\mathcal{G}_H$ , it is known that  
 288  $n_{r+1}[s] = (3(r+2)(r+1))/2 + 1$  for any vertex  $s$ , from which we conclude that it  
 289 has the SG-property. (For more information on  $n_r[s]$ , see any reference on the  $r$ th  
 290 centered triangular number.) For the grid  $H_k$ , as  $k$  is fixed, it is easier to conclude  
 291 that it has the SG-property. Recall that we have shown (see Lemma 1.1) that the  
 292 infinite 3-regular tree does not have this property.

293 **Theorem 2.2** (Discharging Method). *Let  $G$  be an infinite graph with bounded*  
 294 *maximum degree which satisfies the SG-property. Let  $C$  be a vertex set in  $G$ .*  
 295 *Suppose that the discharging method is applied to  $G$  in the following way. In the*  
 296 *charging phase, charge 1 is assigned to each vertex in  $C$  and charge 0 is assigned to*  
 297 *the remaining vertices. In the discharging phase, among other rules, the following*  
 298 *one is respected: no vertex sends charge from it to a vertex at a distance greater*  
 299 *than  $d$ , for a fixed integer  $d$ . If, at the end, every vertex  $v$  of  $G$  has final charge*  
 300  *$\text{chg}(v)$  such that  $q \leq \text{chg}(v) \leq q'$ , where  $q$  and  $q'$  are rational numbers, then*  
 301  *$q \leq d(C, G) \leq q'$ .*

302 *Proof.* Given a set  $W \subseteq V(G)$ , let  $\text{chg}(W) = \sum_{w \in W} \text{chg}(w)$ . Let  $q$ ,  $q'$  and  $d$  be  
 303 as in the hypothesis of the lemma, and let  $s$  be an arbitrary vertex in  $G$ . As in  
 304 the proof of Lemma 2.1, to simplify notation, we let  $n_r[s] = |N_r[s]|$ . Note that  
 305  $q \cdot n_r[s] \leq \text{chg}(N_r[s]) \leq q' \cdot n_r[s]$ .

306 Moreover, notice that  $\text{chg}(N_r[s])$  is at most  $|C \cap N_r[s]|$  plus the charge received  
 307 from vertices outside  $N_r[s]$ , which are contained in  $N_{r+d}[s]$ . Then,  $q \cdot n_r[s] \leq$   
 308  $\text{chg}(N_r[s]) \leq |C \cap N_r[s]| + n_{r+d}[s] - n_r[s]$ . Therefore,

$$d_s(C, G) = \limsup_{r \rightarrow \infty} \frac{|C \cap N_r[s]|}{n_r[s]} \geq q - \limsup_{r \rightarrow \infty} \frac{n_{r+d}[s] - n_r[s]}{n_r[s]} = q.$$

309 The last equality holds because  $\lim_{r \rightarrow \infty} n_{r+d}[s]/n_r[s] = 1$ , by Lemma 2.1(a).

310 Moreover, for  $r > d$ ,  $\text{chg}(N_r[s])$  is at least  $|C \cap N_r[s]|$  minus the charge sent  
 311 to vertices outside  $N_r[s]$ , which comes from vertices in  $N_r[s] \setminus N_{r-d}[s]$ . Then,  
 312  $q' \cdot n_r[s] \geq \text{chg}(N_r[s]) \geq |C \cap N_r[s]| - (n_r[s] - n_{r-d}[s])$ . Therefore,

313

$$d_s(C, G) = \limsup_{r \rightarrow \infty} \frac{|C \cap N_r[s]|}{n_r[s]} \leq q' + \limsup_{r \rightarrow \infty} \frac{n_r[s] - n_{r-d}[s]}{n_r[s]} = q'.$$

314

315

Thus, from Lemma 2.1(b) we conclude that  $q \leq d(C, G) \leq q'$ .  $\square$

316

317

318

The next lemma shows that the usual method of determining the density of a set from periodic patterns, which we showed that is not always valid, works on graphs satisfying the SG-property.

319

320

321

322

323

**Lemma 2.3.** *Let  $G$  be an infinite connected graph with bounded maximum degree that satisfies the SG-property. Let  $\ell, c, c', d$  be positive integers, and let  $C$  be a subset of  $V(G)$ . Suppose that  $V(G)$  can be partitioned into subsets  $V_1, V_2, \dots$  of size  $\ell$  such that,  $c \leq |V_i \cap C| \leq c'$  for each  $i \geq 1$ , and the distance between any two vertices of  $V_i$  is at most  $d$ . Then  $c/\ell \leq d(C, G) \leq c'/\ell$ .*

324

325

326

327

328

329

330

*Proof.* We use the discharging method as stated in Lemma 2.2 with  $q = c/\ell$  and  $q' = c'/\ell$ . Recall that every vertex of  $C$  starts with charge 1 and the vertices outside  $C$  starts with charge 0. In the discharging phase, for every part  $V_i$  of  $V(G)$ , the set of vertices in  $C \cap V_i$  can guarantee charge at least  $q = c/\ell$  and at most  $q' = c'/\ell$  for every vertex of  $V_i$ . Since the distance between any two vertices of  $V_i$  is at most  $d$ , no vertex sends charge to a vertex at a distance greater than  $d$ . From Lemma 2.2, we conclude that  $c/\ell \leq d(C, G) \leq c'/\ell$ .  $\square$

331

332

333

334

335

336

337

338

339

In particular, for  $H_k$ , the above result indicates that to prove a lower bound for the density of an idcode  $C$ , one can show that if  $H_k$  can be covered with a periodic pattern  $H$ , then  $H$  is a pattern (subgraph of  $H_k$  containing vertices of  $C$ ) for which the ratio  $|C \cap V(H)|/|V(H)|$  is minimum possible (a result that might not be so easy to prove). This would lead us to the conclusion that this ratio gives a lower bound for  $d(C, H_k)$ . In Section 3, we prove a lower bound for  $d^*(H_2)$  using the discharging method, as stated in Theorem 2.2, and we also give another proof based on this idea of a pattern  $H$  with best possible ratio. The latter idea also yields a uniqueness proof of the minimum-density periodic idcode of  $H_2$ .

340

### 3. LOWER BOUNDS FOR THE DENSITY OF SOME IDCODS OF $H_k$

341

342

343

344

345

346

347

Karpovsky *et al.* [27] proved that for  $d \geq 2$ , every finite twin-free  $d$ -regular graph  $G$  satisfies  $d^*(G) \geq 2/(d+2)$ . This was done using a double counting argument on the set of possible idcodes. The next theorem shows that the same bound holds for infinite connected graphs with maximum degree bounded by a constant  $d$ , if the graph has the SG-property. To prove this result, we use the discharging method, in a similar way that Cranston and Yu [9] proved the lower bound  $2/5$  for the minimum density  $d^*(\mathcal{G}_H)$  of the hexagonal grid.

348 **Theorem 3.1.** *Let  $\Delta \geq 2$  be a fixed integer and  $G$  be a connected infinite twin-free*  
 349 *graph with maximum degree  $\Delta$ . If  $G$  has the SG-property, then  $d^*(G) \geq 2/(\Delta+2)$ .*  
 350 *In particular,  $d^*(H_k) \geq 2/5$  for every  $k \geq 2$ .*

351 *Proof.* Let  $C$  be an idcode of  $G$ , and let  $q = 2/(\Delta + 2)$ . We apply the discharg-  
 352 ing method with charging rules as stated in Lemma 2.2, and with the following  
 353 discharging rule:

354 (R) If  $v \notin C$  and  $|C[v]| = p$ , then  $v$  receives a charge of  $q/p$  from each vertex  
 355 in  $C[v]$ .

356 We note that only neighbouring vertices exchange charges (thus we may apply  
 357 Lemma 2.2 with  $d = 1$ ). We prove now that  $\text{chg}(v) \geq q$  for every vertex  $v$  in  $G$ .  
 358 Clearly, if  $v \notin C$ , then  $\text{chg}(v) = q$ ; so assume that  $v \in C$ . If  $v$  has no neighbours  
 359 in  $C$ , then for all  $w \in N(v)$  we have  $|C[w]| \geq 2$ , otherwise  $C[v] = C[w]$ . Thus,  
 360 vertex  $v$  sends a charge of at most  $q/2$  to each vertex in  $N(v)$ . As a vertex in  $G$   
 361 has degree at most  $\Delta$ , it follows that  $\text{chg}(v) \geq 1 - \Delta(q/2) = q$ .

362 Suppose now that  $v$  has a neighbour in  $C$ . Then for at most one vertex, say  $w$ ,  
 363 that is a neighbour of  $v$  outside  $C$ , we have that  $C[w] = \{v\}$ ; and for all the  
 364 remaining neighbours  $x$  of  $v$  outside  $C$ , we have that  $|C[x]| \geq 2$ . Thus  $v$  sends a  
 365 charge of at most  $q$  to  $w$  and at most  $q/2$  to the remaining neighbours  $x$  in  $N(v) \setminus C$ .  
 366 Since the degree of  $v$  is at most  $\Delta$ , it follows that  $\text{chg}(v) \geq 1 - q - (\Delta - 2)(q/2) = q$ .

367 As  $\text{chg}(v) \geq q$  for every vertex  $v$  in  $G$ , by Lemma 2.2 we have that  $d(C, G) \geq q$ .  
 368 As this holds for an arbitrary idcode  $C$ , it follows that  $d^*(G) \geq q = 2/(\Delta + 2)$ .  
 369 When  $G$  is the hexagonal grid  $H_k$  with  $k$  rows, the result we have shown implies  
 370 that  $d^*(H_k) \geq 2/5$  for every  $k \geq 2$ .  $\square$

371 If  $C$  is an idcode of a graph  $G$ , then a component of  $G[C]$ , the subgraph induced  
 372 by  $C$ , is called a *cluster* of  $G$  (w.r.t.  $C$ ). If a cluster has precisely (resp. at least)  
 373  $t$  vertices, then it is called a *t-cluster* (resp. *t<sup>+</sup>-cluster*). The unique vertex of a  
 374 1-cluster is also called a 1-cluster. Note that  $G[C]$  has no 2-clusters, otherwise,  
 375 the 2 vertices in such a cluster would have the same identifier. The idcodes shown  
 376 in Figures 1(B) and 1(C) induce only 1-clusters.

377 In what follows, we show that if  $C$  is an idcode of a graph  $G$  such that  $G[C]$   
 378 has no 1-clusters, and  $G$  satisfies certain conditions, then  $d(C, G) \geq 3/7$ .

379 **Theorem 3.2.** *Let  $G$  be a connected infinite twin-free graph with maximum de-*  
 380 *gree 3, and with the SG-property. If  $C$  is an idcode of  $G$  such that  $G[C]$  has no*  
 381 *1-clusters, then  $d(C, G) \geq 3/7$ . In particular,  $d(C, \mathcal{G}_H) \geq 3/7$  and  $d(C, H_k) \geq 3/7$*   
 382 *for every  $k \geq 2$ .*

383 *Proof.* We use the discharging method with charging rules as stated in Lemma 2.2.  
 384 We take  $q = 3/7$ , and consider the following discharging rules:

385 (R1) If  $v \notin C$  and  $|C[v]| = p$ , then  $v$  receives a charge of  $3/(7p)$  from each vertex  
 386 in  $C[v]$ .

387 (R2) If  $c \in C$  and  $|N[c] \cap C| \geq 2$ , then  $c$  sends a charge of  $1/14$  to each neighbour  
 388 in  $N(c) \cap C$ .

389 Let us prove now that  $\text{chg}(v) \geq 3/7$  for every vertex  $v$ . Clearly,  $\text{chg}(v) = 3/7$   
 390 if  $v \notin C$ . Consider now a vertex  $c \in C$ . By hypothesis, we have that  $c$  has

391 at least one neighbour in  $C$ . If  $c$  has exactly one neighbour  $c'$  in  $C$ , then  $c'$   
 392 must have another neighbour in  $C$ . Since  $c$  has at most 2 neighbours outside  $C$ ,  
 393 then  $c$  sends a charge of at most  $3/7$  to one of them, at most  $3/14$  to the other,  
 394 and receives  $1/14$  from  $c'$ . (Note that, if these two neighbours exist, then one  
 395 of them must have another neighbour in  $C$ , distinct from  $c$ ). Hence,  $\text{chg}(c) \geq$   
 396  $1 - 3/7 - 3/14 + 1/14 = 3/7$ . If  $c$  has exactly two neighbours in  $C$ , then  $c$  sends a  
 397 charge of at most  $3/7$  to some neighbour  $w \notin C$  and exactly  $1/14$  to each one of  
 398 the two neighbours in  $C$ . Thus,  $\text{chg}(c) \geq 1 - 3/7 - 2(1/14) = 3/7$ . If  $c$  has exactly  
 399 three neighbours in  $C$ , then  $c$  sends exactly  $1/14$  of charge to each of them. Hence,  
 400  $\text{chg}(c) \geq 1 - 3(1/14) = 11/14 > 3/7$ . The results follow from Lemma 2.2.  $\square$

401 **4. AN IDENTIFYING CODE OF  $H_2$  WITH MINIMUM DENSITY**

402 In this section we prove that  $d^*(H_2) = 9/20$ . For that, we prove first the  
 403 following result.

404 **Lemma 4.1.** *The minimum density of an idcode of  $H_2$  is at most  $9/20$ .*

405 *Proof.* Consider the subgraph, say  $T$ , indicated in Figure 3, which is a subgraph  
 406 of  $H_2$  induced by the vertices from columns 1 to 20. Let  $C$  the set of 18 black  
 407 vertices indicated in  $T$ .

408 Note that, the pattern defined by  $C$  in the first 10 columns of  $T$  is a reflected  
 409 form of the pattern defined by  $C$  in the next 10 columns. We claim that if we  
 410 concatenate infinite copies of  $T$  (side by side), the set of black vertices obtained is  
 411 an idcode of  $H_2$  (with period 20). We leave to the reader to check this fact (it is  
 412 enough to check the first 11 columns, and the columns 20 and 21). By Lemma 2.3  
 413 we conclude that  $d^*(H_2) \leq 9/20$ .  $\square$





FIGURE 3. An idcode of  $T \subset H_2$ , which gives an idcode of  $H_2$

414 To show that  $d^*(H_2) \geq 9/20$ , we present two different proofs, which are closely  
 415 related. Both are based on the patterns defined by an idcode  $C$  in the graph  $H_2$ .  
 416 To study these patterns, we consider that the graph  $H_2$  is an infinite strip that  
 417 can be “split” into “sequential” 4-vertex sets, defined formally in what follows.

418 For an integer  $x$ , we say that a vertex of column  $x$  of  $H_2$  is *cubic* if it has  
 419 degree 3 in  $H_2$ . We adopt the convention that when  $x$  is odd then the vertices  
 420 in column  $x$  are cubic. For an odd integer  $x$ , we denote by  $Q_x$  the set of vertices  
 421  $\{(x, 1), (x + 1, 1), (x, 2), (x + 1, 2)\}$ , and call it a *quartet*.

422 Note that  $H_2[Q_x]$  is a  $\square$ -shaped path in  $H_2$  with 4 vertices, and  $V(H_2)$  is the  
 423 disjoint union of quartets  $Q_x$  such that  $x$  is an odd integer. Given a quartet  $Q_x$ ,

FIGURE 4. Quartets  $Q_x^L$ ,  $Q_x$  and  $Q_x^R$ 


424 we also refer to  $Q_{x-2}$  (resp.  $Q_{x+2}$ ), its left (resp. right) quartet, as  $Q_x^L$  (resp.  $Q_x^R$ ),  
 425 see Figure 4.

426 For a given idcode  $C$ , we say that  $Q_x$  is *type  $i$*  (resp. *type  $i^+$* ) if  $|Q_x \cap C| = i$   
 427 (resp.  $|Q_x \cap C| \geq i$ ). Type 1 quartets  $Q_x$  play an important role in the proofs.  
 428 If the single vertex in the idcode that belongs to  $Q_x$  is cubic (resp. not cubic)  
 429 in  $H_2$ , we say that  $Q_x$  is *type 1-cubic* (resp. *type 1-noncubic*). See Figure 6. All  
 430 references to types assume that an idcode is clear from the context.

431 The next lemmas tell us, for each quartet  $Q_x$  of type  $i$  ( $1 \leq i \leq 3$ ), which  
 432 are the possible (or forbidden) types of its neighbouring quartets  $Q_x^L$  and/or  $Q_x^R$ .  
 433 Once we have these results, we can either use the discharging method or an idea  
 434 based on the average density of patterns defined by consecutive quartets.

435 We denote by  $(H_2, C, x)$  a triple consisting of the grid  $H_2$ , an idcode  $C$  of  $H_2$ ,  
 436 and an odd integer  $x$ . In the figures, vertices coloured black belong to  $C$ , vertices  
 437 coloured gray may belong to  $C$ .

438 **Lemma 4.2** ( $Q_x$  is type 0). *Consider a triple  $(H_2, C, x)$ . If  $Q_x$  is type 0, then*  
 439  *$Q_x^L$  is type 4; moreover,  $Q_x^R$  is type  $3^+$  and  $C \cap Q_x^R$  contains two cubic vertices.*

FIGURE 5. Quartet  $Q_x$  is type 0 implies quartet  $Q_x^L$  is type 4


440 *Proof.* If  $Q_x$  is type 0, it is immediate that all vertices in columns  $x-1$  and  $x+2$   
 441 must be in  $C$ , since all vertices of  $Q_x$  must have a nonempty identifier. As  $C$  is  
 442 an idcode, the vertices of column  $x-2$  must belong to  $C$ ; thus,  $Q_x^L$  is type 4. See  
 443 Figure 5. Since  $H_2[C]$  has no 2-clusters,  $Q_x^R$  is type  $3^+$ .  $\square$

444 **Lemma 4.3** ( $Q_x$  is type 1). *Consider a triple  $(H_2, C, x)$ . If  $Q_x$  is type 1, then*  
 445 *the following holds.*

- 446 (a) *If  $Q_x$  is type 1-cubic, then  $Q_x^L$  is type  $2^+$  and  $Q_x^R$  is type  $3^+$ .*  
 447 (b) *If  $Q_x$  is type 1-noncubic, then  $Q_x^L$  is type  $3^+$  and  $Q_x^R$  is type  $2^+$ .*

448 *Proof.* For simplicity, rename the vertices of  $Q_x^L \cup Q_x \cup Q_x^R$  as shown in Figure 6.

449 To prove (a), let  $Q_x$  be type 1-cubic, and assume without loss of generality that  
 450  $Q_x \cap C = \{x_1\}$ . See Figure 6(A).

FIGURE 6. Quartet  $Q_x$  is type 1

451 • If  $v_1 \in C$ , then  $u_1 \in C$ , otherwise  $\{v_1, x_1\}$  would induce a 2-cluster in  $H_2$ ,  
 452 a contradiction. Thus,  $Q_x^L \cap C \supseteq \{v_1, u_1\}$  and therefore  $Q_x^L$  is type  $2^+$ . If  $v_1 \notin C$ ,  
 453  $C$ , then  $v_2 \in C$ , otherwise  $C[x_1] = C[x_2]$ , a contradiction. Moreover,  $u_1 \in C$ ,  
 454 otherwise  $C[v_1] = C[x_1]$ . Thus,  $Q_x^L \cap C \supseteq \{v_2, u_1\}$  and therefore  $Q_x^L$  is type  $2^+$ .  
 455 • Clearly,  $z_2 \in C$ , otherwise  $C[y_2] = \emptyset$ . If  $z_1 \in C$ , then  $|Q_x^R \cap C| \geq 3$ , otherwise  
 456  $\{z_1, z_2\}$  would induce a 2-cluster in  $H_2$ . Hence,  $Q_x^R$  is type  $3^+$ . If  $z_1 \notin C$ , then  
 457  $w_1 \in C$ , otherwise  $C[z_1] = C[y_2]$ . Moreover,  $w_2 \in C$ , otherwise  $C[y_2] = C[z_2]$ .  
 458 Thus,  $Q_x^R \cap C = \{z_2, w_1, w_2\}$ , and  $Q_x^R$  is type 3.

459 To prove (b), let  $Q_x$  be type 1-noncubic, and assume without loss of generality  
 460 that  $Q_x \cap C = \{y_2\}$ . See Figure 6(B).


461 • Clearly,  $v_1 \in C$ , otherwise  $C[x_1] = \emptyset$ . Moreover,  $u_1 \in C$ , otherwise  $C[x_1] = C[v_1]$ .  
 462 If  $u_2 \notin C$ , then  $v_2 \in C$  (because  $C[v_2] \neq \emptyset$ ). Thus,  $Q_x^L$  is type  $3^+$ .

463 • Clearly,  $z_1 \in C$  (because  $C[y_1] \neq \emptyset$ ). If  $z_2 \in C$ , then  $Q_x^R$  is type  $2^+$ . If  $z_2 \notin C$ ,  
 464 then  $w_1 \in C$ , otherwise  $C[z_1] = C[y_1]$ , a contradiction. Hence,  $Q_x^R$  is type  $2^+$ .  $\square$

465 **Lemma 4.4** ( $Q_x$  is type 2). Consider a triple  $(H_2, C, x)$ . If  $Q_x$  is type 2, then  
 466  $Q_x^L$  and  $Q_x^R$  may not be both type 1.

467 *Proof.* Suppose, by contradiction, that both  $Q_x^L$  and  $Q_x^R$  are type 1. By Lemma 4.3,  
 468 if  $Q_x^L$  (resp.  $Q_x^R$ ) is type 1-cubic (resp. 1-noncubic), then  $Q_x$  is type  $3^+$ . Thus,  
 469 let us suppose now that  $Q_x^L$  is type 1-noncubic,  $Q_x^L \cap C = \{u\}$ ; and  $Q_x^R$  is type  
 470 1-cubic,  $Q_x^R \cap C = \{w\}$ .

471 First, assume that  $u$  and  $w$  are in the same row, say 2. See Figure 7(A). Then  
 472  $(x, 1) \in C$ , because  $C[(x-1, 1)] \neq \emptyset$ . Note that one of the vertices  $(x+1, 1)$  or  
 473  $(x, 2)$  belongs to  $C$ , because  $C[(x-1, 1)] \neq C[(x, 1)]$ . If  $(x+1, 1) \in C$ , then  $(x, 1)$   
 474 and  $(x+1, 1)$  would induce a 2-cluster in  $H_2$ , a contradiction. If  $(x, 2) \in C$ , then  
 475  $C[(x+2, 2)] = \{w\} = C[(x+2, 1)]$ , a contradiction.

FIGURE 7. Quartet  $Q_x$  is type 2

476 If  $u$  and  $w$  are in different rows, assume without loss of generality that  $u$  is in  
 477 row 2 and  $w$  is in row 1. See Figure 7(B). Then  $(x, 1) \in C$ , because  $C[(x-1, 1)] \neq \emptyset$ .  
 478 If both  $(x+1, 1)$  and  $(x+1, 2)$  do not belong to  $C$ , then  $C[(x+2, 2)] = C[(x+2, 1)]$ ,  
 479 a contradiction. Thus, exactly one of them belongs to  $C$ . If  $(x+1, 1) \in C$ ,  
 480 then  $C[(x+1, 2)] = \emptyset$ , a contradiction. Hence,  $(x+1, 2) \in C$ . But in this  
 481 case,  $C[(x-1, 1)] = C[(x, 1)]$ , a contradiction. This concludes the proof of the  
 482 lemma.  $\square$

483 We state now a lemma that will be helpful to simplify the proof of the next  
 484 theorem.

485 **Lemma 4.5.** *The grid  $H_2$  has idcodes of minimum density without type 0 quartets.*

486 *Proof.* Let  $C$  be an idcode of  $H_2$ , and  $Q_x$  be a quartet of type 0. By Lemma 4.2,  
 487  $Q_x^L$  is type 4. It is simple to verify that  $C' = C \setminus \{(x-1, 1)\} \cup \{(x, 1)\}$  is an idcode  
 488 of  $H_2$  such that  $Q_x$  is type 1 and  $Q_x^L$  is type 3. Thus,  $d(C', H_2) = d(C, H_2)$ . This  
 489 means that If  $C$  is an idcode of minimum density containing type 0 quartets, then  
 490  $H_2$  has also an idcode of the same density without type 0 quartets.  $\square$

491 **Remark.** The previous lemma does not guarantee anything about the elimination  
 492 of type 4 quartets. We note that by doing a local change (more involved than the  
 493 above one) we may also eliminate type 4 quartets and obtain an idcode of equal  
 494 or possibly smaller density. We do not prove this statement as we do not use it  
 495 here. Moreover, later we present arguments showing that type 4 quartets do not  
 496 occur in minimum density idcodes of  $H_2$ .

497 Before going to the next proof, the reader may highlight in Figure 3 the 1-cubic  
 498 and 1-noncubic quartets, and check the statements of Lemma 4.3 and Lemma 4.4  
 499 with respect to the quartets of this figure. This will help the understanding of the  
 500 discharging rule (resp. the idea based on the average density) used in the next  
 501 two proofs.

502 **Theorem 4.6.** *The minimum density of an idcode of  $H_2$  is precisely  $9/20$ .*

503 *Proof.* We use the discharging method to prove that  $d^*(H_2) \geq 9/20$ . For that,  
 504 let  $C$  be a minimum identifying code of  $H_2$  that has no quartets of type 0  
 505 (cf. Lemma 4.5). In the charging phase, we proceed as stated in Lemma 2.2:  
 506 we set  $\text{chg}(v) = 1$  if  $v \in C$ , and  $\text{chg}(v) = 0$ , otherwise. We shall prove that after  
 507 the discharging phase (to be defined), we have  $\text{chg}(Q_x) \geq 9/5$  for each quartet  
 508  $Q_x$ . If this happens, then the total charge of each  $Q_x$  can be distributed among  
 509 its 4 vertices, and we get  $\text{chg}(v) \geq 9/20$  for each vertex  $v$  in  $Q_x$ . Thus, we say  
 510 that a quartet  $Q_x$  is *satisfied* if  $\text{chg}(Q_x) \geq 9/5$ , otherwise, it is *unsatisfied*.

511 After the charging phase, only type 1 quartets are unsatisfied. Apply the fol-  
 512 lowing discharging rule.

- 513 (R) As long as there are type 1 quartets  $Q_x$  that are unsatisfied,  
 514 (a) if  $Q_x$  is 1-cubic, then it receives  $1/5$  from  $Q_x^L$ , and  $3/5$  from  $Q_x^R$ ;  
 515 (b) if  $Q_x$  is 1-noncubic, then it receives  $3/5$  from  $Q_x^L$ , and  $1/5$  from  $Q_x^R$ .

516 We prove now that each quartet  $Q_x$  is satisfied after the discharging phase.

517 **Case 1.**  $Q_x$  is type 1.

518 If  $Q_x$  is type 1, then by Lemma 4.3, both  $Q_x^L$  and  $Q_x^R$  have charge at least 2.  
 519 Thus, they have sufficient charge to send to  $Q_x$ . If  $Q_x$  is type 1-cubic, it received  
 520  $1/5$  from  $Q_x^L$  and  $3/5$  from  $Q_x^R$ . If  $Q_x$  is type 1-noncubic, then it received  $3/5$  from  
 521  $Q_x^L$ , and  $1/5$  from  $Q_x^R$ . Hence, in both cases,  $\text{chg}(Q_x) = 1 + 1/5 + 3/5 = 9/5$ , and  
 522 therefore  $Q_x$  is satisfied.

523 **Case 2.**  $Q_x$  is type 2.

524 If  $Q_x$  is type 2, then by Lemma 4.4,  $Q_x^L$  and  $Q_x^R$  are not both type 1. If  $Q_x^L$  is  
 525 type 1, then by Lemma 4.3, it is type 1-noncubic (because  $Q_x$  is type 2). Thus,  
 526 according to rule (R)(b),  $Q_x^L$  received  $1/5$  from  $Q_x$ . Since  $Q_x$  did not send charge to  
 527  $Q_x^R$  (because  $Q_x^R$  is not type 1) we have that  $\text{chg}(Q_x) = 2 - 1/5 = 9/5$ . Analogously,  
 528 if  $Q_x^R$  is type 1, then by Lemma 4.3, it is type 1-cubic (because  $Q_x$  is type 2). Thus,  
 529 according to rule (R)(a),  $Q_x^R$  received  $1/5$  from  $Q_x$ . Since  $Q_x$  did not send charge  
 530 to  $Q_x^L$  (because  $Q_x^L$  is not type 1), we have that  $\text{chg}(Q_x) = 2 - 1/5 = 9/5$ .

531 **Case 3.**  $Q_x$  is type  $3^+$ .

532 The only possibility for  $Q_x$  to decrease its initial charge is when it has type 1  
 533 neighbours. In the worst case, when both  $Q_x^L$  and  $Q_x^R$  are type 1,  $Q_x$  sends at  
 534 most  $3/5$  to each of them. Thus,  $\text{chg}(Q_x) \geq 3 - 3/5 - 3/5 = 9/5$ .

535 Since every quartet  $Q_x$  is satisfied, by Lemma 2.2, we have that  $d(C, H_2) \geq 9/20$ .  
 536 Using Lemma 4.1, we conclude that  $d^*(H_2) = 9/20$ .  $\square$

537 From the previous result and the fact that the idcode shown in Figure 4.1 has  
 538 density at most  $9/20$ , we conclude the following result.

539 **Corollary 2.** *The idcode shown in Figure 4.1 is a periodic idcode of  $H_2$  with*  
 540 *minimum density.*

541 In what follows we present a second proof of Theorem 4.6 which is based on the  
 542 idea of finding a periodic pattern that covers  $H_2$  and has the minimum possible  
 543 density. This proof also uses Lemmas 4.2 to 4.5, and it is based on the fact  
 544 (mentioned in Section 5) that  $H_2$  has a periodic idcode of minimum density. As  
 545 we will see, the information provided by this proof, combined with further tests,  
 546 will lead us to conclude that the periodic idcode that we have found is unique.

547 **Proof 2.** (of Theorem 4.6). Let  $C$  be an idcode of minimum density in  $H_2$  that  
 548 has no quartets of type 0. If  $C$  has no quartets of type 1, then all quartets in  $C$   
 549 are of type  $2^+$ , and in this case,  $d(C, H_2) \geq 1/2$ , contradicting Lemma 4.1. Thus,  
 550  $C$  has a quartet of type 1, and by Lemma 4.3 we conclude that  $C$  has a quartet  
 551 of type  $3^+$ .

552 Now let us consider that  $H_2$  (seen as a concatenation of quartets) can be split  
 553 into subgraphs corresponding to special sequences of consecutive quartets. We  
 554 are interested in sequences, which we call *S(3)-sequences*, defined as those *starting*  
 555 *with a quartet of type  $3^+$  and containing exactly one quartet of type  $3^+$* . The  
 556 *S(3)-sequences* whose second quartet is of type 1 (resp. type 2) are called *S(3, 1)-*  
 557 *sequences* (resp. *S(3, 2)-sequences*). (We remark that not allowing the presence of  
 558 another quartet of type  $3^+$  is not a restriction to the size of the periods of the  
 559 patterns we want to study. We may have different *S(3)-sequences*, and later we

560 allow them to be concatenated, so that periods with many occurrences of quartets  
561 of type  $3^+$  are made possible.)

562 For an  $S(3)$ -sequence  $S$ , let  $I(S) = (i_1, i_2, \dots)$  be the sequence where each  $i_j \in$   
563  $\{1, 2, 3, 4\}$  indicates the type of each of the  $j$ th quartet in  $S$ . In this proof,  $i_j = i^+$   
564 means that  $i_j \in \{i, i + 1\}$ . A simplified notation such as  $I(S) = (3^+, 1, 2, 2, 1^+)$   
565 stands for  $I(S) \in \{(3, 1, 2, 2, 1), (3, 1, 2, 2, 2), (4, 1, 2, 2, 1), (4, 1, 2, 2, 2)\}$ . We denote  
566 by  $H[S]$  the subgraph of  $H_2$  induced by the quartets in  $S$ , and denote by  $C(S)$   
567 the restriction of  $C$  to  $H[S]$ . We are interested in  $d(C(S), H[S])$ , the density of  
568  $C(S)$  with respect to  $H[S]$ .

569 Note that  $I(S)$  may not contain subsequences of the form  $(1, 2, 1)$ ,  $(2, 1, 2)$  or  
570  $(1, 1)$  because of Lemmas 4.3 and 4.4. If  $S$  is an infinite  $S(3)$ -sequence, then  $I(S) =$   
571  $(3^+, 1, 2, 2, \dots)$  or  $I(S) = (3^+, 2, 2, \dots)$ , and therefore  $d(C(S), H[S]) \geq 1/2$ . If  $S$   
572 is a finite  $S(3, 1)$ -sequence, then  $I(S)$  contains at most two (non-consecutive) 1's.

573 Let  $S_t$  be a finite  $S(3, 1)$ -sequence of length  $t$ , let  $I_t = I(S_t)$ , and let  $C_t$  be  
574 the restriction of  $C$  to  $S_t$ . The possibilities for  $I_t$  are:  $I_1 = (3^+)$ ,  $I_2 = (3^+, 1)$ ,  
575  $I_3 = (3^+, 1, 2)$ ,  $I_4 = (3^+, 1, 2, 2)$ ,  $I_5 = (3^+, 1, 2, 2, 1^+)$ , and  $I_t = (3^+, 1, 2, \dots, 2, 1^+)$   
576 if  $t > 5$ . Thus  $d(C_t, H[S_t]) \geq 1/2$ , for  $1 \leq t \leq 4$ ,  $d(C_5, H[S_5]) \geq 9/20$  and  
577  $d(C_t, H[S_t]) \geq (3 + 1 + 2(t - 3) + 1)/4t = (2t - 1)/4t > 9/20$  if  $t > 5$ . Thus  
578 the minimum density  $9/20$  may possibly occur for  $S(3)$ -sequences of length 5 with  
579 sequence of types  $(3, 1, 2, 2, 1)$ .

580 It is easy to see that if  $S$  is a finite  $S(3, 2)$ -sequence, then  $d(C(S), H[S]) \geq 1/2$   
581 (because  $I(S)$  contains at most one 1). This ends the proof that all  $S(3)$ -sequences  
582 of  $H_2$  have density at least  $9/20$ . Thus,  $d(C, H_2) \geq 9/20$  (as  $H_2$  has a minimum-  
583 density periodic idcode). Combining this result with Lemma 4.1, we conclude that  
584  $d^*(H_2) = 9/20$ .  $\square$

585 **Remark on the uniqueness of a periodic minimum-density idcode for  $H_2$ .**

586 By Corollary 2, the idcode shown in Figure 3 is a periodic idcode of  $H_2$  with  
587 minimum density. An interesting question is whether this idcode is unique, among  
588 the periodic ones. The meaning of uniqueness will be clear in what follows.

589 The second proof of Theorem 4.6 suggests that to construct a periodic minimum-  
590 density idcode for  $H_2$  we should look for idcodes that define  $S(3, 1)$ -sequences of  
591 length 5 of type  $(3, 1, 2, 2, 1)$ , and try to concatenate them to see whether they  
592 yield a periodic idcode.

593 As the reader may check, the  $S(3, 1)$ -sequence, say  $S$ , corresponding to the  
594 5 initial quartets (first 10 columns) shown in Figure 3 is of type  $(3, 1, 2, 2, 1)$ .  
595 However, the concatenation  $SS$  does not define an idcode of  $H_2$  restricted to these  
596 sequences. But, as one can see in Figure 3, after  $S$ , the next sequence of 5 quartets,  
597 say  $S'$ , which is a reflected form of  $S$  is also an  $S(3)$ -sequence of type  $(3, 1, 2, 2, 1)$ .  
598 As we mentioned before, this is an idcode of  $H_2$  with period 20. This is not  
599 the way we obtained this idcode. In fact, this idcode was obtained by an ad hoc  
600 method, and we used it as an inspiration to derive the properties (Lemmas 4.2-4.5)  
601 that we proved. These lemmas, in turn, helped us in the lower bound proof. If  
602 a sequence such as  $S$  could not be found, one should look for  $S(3)$ -sequences of

603 lengths  $t = 6, 7, \dots$ , as they would be the next candidates (if we did not know an  
604 idcode with density  $9/20$ ).

605 Let us now investigate whether the idcode shown in Figure 3 is the unique  
606 periodic idcode of  $H_2$  with density  $9/20$ . We note that  $S$  and  $S'$  are the unique  
607  $S(3)$ -sequences of type  $(3, 1, 2, 2, 1)$  (we have verified this by running a program).  
608 We also note that the concatenation  $S'S'$  does not define an idcode. So, for the  
609 moment we may say that the answer to this question is “yes”, if we consider  
610 minimum idcodes without type 0 quartets (as we proved).

611 The question now is whether there are minimum-density idcodes containing  
612 type 0 quartets. We will not go into details, but we can prove that carrying  
613 out analogous arguments as those we used for  $S(3, 1)$ - and  $S(3, 2)$ -sequences, the  
614 answer is “no”. By Lemma 4.2, a type 0 quartet is preceded by a type 4 quartet,  
615 and is succeeded by a type  $3^+$  quartet. Using this fact, we can show that any  
616  $S(3)$ -sequence that is of subtype  $S(4, 0)$  has density greater than  $9/20$ . Thus, we  
617 conclude that the idcode shown in Figure 3 is the unique periodic idcode of  $H_2$   
618 with minimum density. This idcode was also obtained by running a computer  
619 program, about which we report in the next section.

620 We note that, the idea we mentioned after Lemma 2.3 to prove lower bound for  
621 the density of idcodes of  $H_k$  —based on periodic patterns with minimum density—  
622 is basically the idea behind the study we have carried out on the types of sequences  
623 of  $H_2$ . This study led us to conclude that the periodic pattern  $H$  defined by the  
624 concatenation  $SS'$  is the shortest periodic pattern that has the minimum density  
625  $9/20$ . Of course, we may say that  $S'S$  is also such a shortest periodic pattern, but  
626 here we consider that they are equivalent.

## 627 5. MINIMUM-DENSITY IDENTIFYING CODES OF $H_3$ , $H_4$ AND $H_5$

628 In this section we present minimum-density idcodes for  $H_3$ ,  $H_4$  and  $H_5$  that we  
629 found with an algorithm implemented in C++. We describe briefly the algorithm,  
630 then exhibit some of these idcodes and the values  $d^*(H_3)$ ,  $d^*(H_4)$  and  $d^*(H_5)$ .

631 The algorithm that we implemented searches for a periodic idcode for these  
632 grids, and uses an idea that was already proposed in 2018 by Jiang [24], to find  
633 minimum-density idcodes for square grids  $S_k$  with finite number  $k$  of rows. We were  
634 not aware of his algorithm, although we knew about his results on  $S_k$ . Jiang [24]  
635 proved that such grids have idcodes with minimum density that are periodic, and  
636 described an algorithm to find them. His work presents in detail an algorithm that  
637 constructs a weighted directed graph (associated with  $S_k$ ) in which a minimum  
638 mean cycle corresponds to a periodic minimum-density idcode of  $S_k$ . Unfortu-  
639 nately, the size of this graph is exponential in  $k$ . With his implementation in C, in  
640 2018 Jiang was able to obtain optimum idcodes for  $S_4$  and  $S_5$ . We used basically  
641 the same idea for  $H_k$ . For completeness, we describe briefly the construction of  
642 this graph, using the terminology introduced by Jiang.

643 We do not prove here that  $H_k$  has finite periodic idcodes that have minimum  
644 density, but this result holds. A proof similar to the one presented by Jiang [24]

645 for  $S_k$  can be done for  $H_k$ , using the idea based on the concept of bars, which is  
 646 central here, and is defined in what follows.

647 For  $\ell \geq 1$  and  $k \geq 2$ , any subgraph of  $H_k$  induced by  $\{j_1, \dots, j_\ell\} \times [k]$ ,  
 648 where  $j_1 \leq j_2 \leq \dots \leq j_\ell$  are  $\ell$  consecutive columns of  $H_k$ , is called an  $\ell$ -bar (see  
 649 Figure 9). Let  $R$  be any  $\ell$ -bar with  $\ell \geq 3$  in  $H_k$ , and let  $R'$  be the  $(\ell - 2)$ -bar  
 650 consisting of the middle columns of  $R$  (obtained by excluding the first and the  
 651 last columns of  $R$ ). We say that a subset  $C$  of vertices of  $R$  is a *barcode* of  $R$  if  
 652  $C[v] \neq \emptyset$  and  $C[u] \neq C[v]$  for every distinct  $u, v \in R'$ . We adopt the convention  
 653 that the first column of each 4-bar of  $H_k$  is indexed by an odd number.

### 654 5.1. CONSTRUCTION OF THE ARC-WEIGHTED DIRECTED GRAPH $G_{k,4,j}$

655 For  $k \geq 2$  and  $5 \leq j \leq 8$ , let  $G_{k,4,j} = (V, A)$  denote the  $j$ -configuration graph  
 656 of the *idcodes* of  $H_k$  defined as follows. The vertex set  $V$  of this graph consists of  
 657 barcodes  $C$  of any 4-bar of  $H_k$ . There is an arc from  $C$  to  $C'$  if there is a barcode  $Q$   
 658 of a  $j$ -bar  $B$  of  $H_k$  such that  $C$  (resp.  $C'$ ) is the restriction of  $Q$  to the first (resp.  
 659 last) 4 columns of  $B$ . In this case, the arc from  $C$  to  $C'$  gets weight  $|Q| - |C|$ . Note  
 660 that,  $|V| \leq 2^{4k}$  and  $|A| \leq 2^{jk}$ . In our implementation, we used  $j = 6$  and  $j = 8$   
 661 (as in this case we have to deal only with 4-bars whose first column is indexed by  
 662 an odd number).

663 Jiang [24] considered, for the grid  $S_k$ , the graph  $G_{k,4,5}$ , described above for  $H_k$   
 664 (for  $S_k$ , the 4-bars correspond to subgraphs of  $S_k$ ). He showed that in this graph,  
 665 each 4-bar pattern of a periodic idcode for  $S_k$  corresponds to a directed cycle and  
 666 vice-versa. We defined  $G_{k,4,j}$  for  $5 \leq j \leq 8$ . It is not difficult to see that an  
 667 equivalent statement also holds for  $j = 6, 7, 8$ , and for the grid  $H_k$ . Thus, in this  
 668 case, the density of a minimum periodic idcode in  $G_{k,4,j}$  is  $w(Z)/pk$ , where  $w(Z)$   
 669 is the weight of a minimum mean cycle  $Z$  in the configuration graph  $G_{k,4,j}$  and  $p$   
 670 is the period. (If  $Z$  is a cycle, then the *mean weight* of  $Z$  is the ratio between the  
 671 total weight  $w(Z)$  of the arcs in  $Z$  and the number of arcs in  $Z$ .)

672 In Figure 9 we show a minimum density periodic idcode (with period 8) for  
 673  $H_4$  that was found in the 8-configuration graph  $G_{4,4,8}$ . The two curly braces  
 674 indicate two consecutive 4-bars (corresponding to two barcodes, say  $C$  and  $C'$ ,  
 675 which are adjacent vertices in this graph). In this case,  $Q$  is the barcode of the  
 676 8-bar (formed by the indicated 4-bars), and the weight of the arc from  $C$  to  $C'$  is  
 677  $|Q| - |C| = 14 - 7 = 7$ . This solution corresponds to the weighted directed cycle  
 678  $Z = (C, C')$  that has length  $|Z| = 2$  and weight  $w(Z) = 14$  (with mean weight  
 679  $w(Z)/2 = 14/2 = 7$ ). In this case, the period is  $p = 8$ . Thus, the density of this  
 680 solution is  $w(Z)/(8 \cdot 4) = 14/32 = 7/16$ . We observe that when  $j = 8$  the period  
 681 is  $|Z| \cdot 4$ . (but the period is  $|Z| \cdot 2$  if  $j = 6$ , as in this there is an overlap of 2  
 682 columns for each two adjacent barcodes).

683 It is well known that the *minimum mean cycle problem* on a graph with  $n$   
 684 vertices and  $m$  arcs can be solved in  $O(nm)$  time by Karp's algorithm [26]. This  
 685 is the algorithm that Jiang [24] used in his implementation for  $S_k$ . For  $H_k$ , we  
 686 use Hartmann-Orlin's algorithm [15], which is an improved version of Karp's algo-  
 687 rithm, to find a minimum mean cycle. We implemented a program in C++, using

TABLE 1. Sizes of the configuration graphs generated by our implementation and total running times.

(A) Data for  $j = 6$ 

Configuration graph	# vertices	# edges	Total running time
$G_{2,4,6}$ ( $H_2$ )	144	1359	8 ms
$G_{3,4,6}$ ( $H_3$ )	1896	57723	253 ms
$G_{4,4,6}$ ( $H_4$ )	5870	63095	8 s
$G_{5,4,6}$ ( $H_5$ )	63751	1650188	87 m

(B) Data for  $j = 8$ 

Configuration graph	# vertices	# edges	Total running time
$G_{2,4,8}$ ( $H_2$ )	144	12894	46 ms
$G_{3,4,8}$ ( $H_3$ )	1896	1784401	9 s
$G_{4,4,8}$ ( $H_4$ )	5870	3291346	820 s
$G_{5,4,8}$ ( $H_5$ )	63751	248161004	928 m

688 *lemon*<sup>1</sup> library for graphs: it builds the graph  $G_{k,4,j}$ , finds a minimum mean cycle  
 689 and outputs an idcode with minimum density for  $H_k$ . This implementation can  
 690 be found in [32].

691 We run this program to find minimum-density idcodes for  $H_3$ ,  $H_4$  and  $H_5$ .  
 692 This program constructed  $G_{3,4,6}$ ,  $G_{4,4,8}$ ,  $G_{5,4,6}$ , and obtained  $d^*(H_3) = 6/13$ ,  
 693  $d^*(H_4) = 7/16$  and  $d^*(H_5) = 11/25$ . The corresponding idcodes for these grids  
 694 are depicted in Figures 8, 9 and 10. In Table 1, we indicate the size of these  
 695 configuration graphs and the total running time the program needed to find an  
 696 optimal solution. The running times for  $j = 8$  are included to show the difference  
 697 when compared to  $j = 6$ . The code was compiled with g++ 11.4.0 and option  
 698 -O3, and executed in a computer with Intel(R) Xeon(R) CPU E7- 2870 @ 2.40GHz  
 699 processor with 512 GB of RAM.




FIGURE 8. A minimum-density idcode of  $H_3$  found in the graph  $G_{3,4,6}$  (density  $6/13 \approx 0.46153$ , period 26)

700 **Theorem 5.1.** For  $k = 3, 4, 5$ , the idcodes for  $H_k$  shown in Figures 8, 9 and 10  
 701 have minimum density. The corresponding densities of these idcodes are  $d^*(H_3) =$   
 702  $6/13$ ,  $d^*(H_4) = 7/16$  and  $d^*(H_5) = 11/25$ .

<sup>1</sup><https://lemon.cs.elte.hu/trac/lemon>




FIGURE 9. A minimum-density idcode of  $H_4$  found in the graph  $G_{4,4,8}$  (density  $7/16 = 0.4375$ , period 8)




FIGURE 10. A minimum-density idcode of  $H_5$  found in the graph  $G_{5,4,6}$  (density  $11/25 = 0.44$ , period 10)




FIGURE 11. A minimum-density idcode of  $H_5$  found in the graph  $G_{5,4,8}$  (density  $11/25 = 0.44$ )

703 As a side remark, we observe that if instead of considering 4-bars, we consider  
 704 3-bars (to define the vertices of the graph), and define adjacency of vertices in  
 705 an analogous way, the corresponding graphs  $G_{k,3,5}$  or  $G_{k,3,6}$  for  $S_k$  or  $H_k$  do not  
 706 have the desired property (as some arcs would indicate a wrong adjacency). We  
 707 leave to the reader finding examples to verify this statement. But such incorrect  
 708 adjacencies occur rarely. Since it is much faster to work with 3-bars, one possibility  
 709 is to work with 3-bars, and check whether the solution found does not have wrong  
 710 adjacencies, as in this case, an optimum solution may be found more quickly.

711 We conclude this section mentioning that with our implementation we were  
 712 not able to find a minimum-density idcode for  $H_6$  using the computer resources  
 713 available to us.

714

## 6. CONCLUDING REMARKS

715 We note that for  $H_3$  we have found only the minimum-density idcode shown  
716 in Figure 8. But we are not claiming that it is unique. For  $H_4$  and  $H_5$ , we  
717 have found other minimum-density idcodes with different periods. For  $H_5$  we note  
718 that the minimum-density idcode shown in Figure 11 is different from the idcode  
719 shown in Figure 10, but both have period 10. By considering the graph  $G_{5,4,8}$ ,  
720 the corresponding program output the solution of Figure 11 indicating that the  
721 period is 20. We noted that the columns from 1–10 of this idcode is equal to the  
722 columns from 11–20. Thus, we may say that the period of this idcode is 10. This  
723 does not indicate that the program is incorrect. Clearly, when  $j = 8$ , the program  
724 outputs a solution whose period is always a multiple of 4, while when  $j = 6$  the  
725 program outputs a solution whose period is a multiple of 2.

726 With this respect, we note that if  $H_k$  has a minimum-density idcode with pe-  
727 riod  $p$ , even when  $p$  is odd, an idcode with the same density and possibly different  
728 period can be found in the graph  $G_{k,4,6}$  and  $G_{k,4,8}$ . This is true because there is  
729 a (smallest) multiple of  $p$  which is always a multiple of 2 or of 4, and therefore  
730 such a solution will be present in the corresponding graphs. We observe that our  
731 program finds one optimal solution (a minimum mean cycle) but not all optimal  
732 solutions.

733 Our implementation may possibly be improved if we can eliminate from the  
734 graph  $G_{k,4,j}$  some vertices and arcs which we are sure will not occur in an optimal  
735 solution. For example, barcodes corresponding to the set of all vertices in a 4-bar,  
736 or possibly barcodes whose densities are much larger than some known upper  
737 bound for the minimum-density idcode. But to implement such steps safely, some  
738 proofs are needed. We also believe that a more substantial improvement is needed  
739 to be able to solve for larger  $k$ . We are working on this topic and hope that in a  
740 forthcoming paper we will be able to present good upper bounds for  $d^*(H_k)$ , for  
741 all  $k \geq 6$ .

742 **Acknowledgments.** The authors would like to thank the referees for the careful reading  
743 of the paper, for pointing out some references, and for the valuable suggestions that  
744 improved the content and presentation of this paper. The authors are also thankful to  
745 Antoine Lobstein for the updated bibliography on identifying codes and related topics  
746 that he maintained for so many years, and to Devin Jean [22] for continuing his work.  
747 The authors extend their gratitude to the Brazilian research agencies CAPES, CNPq,  
748 FAPESP and FUNCAP for the financial support.

749

## REFERENCES

- 750 1. Yael Ben-Haim and Simon Litsyn, *Exact minimum density of codes identifying vertices in*  
751 *the square grid*, SIAM J. Discrete Math. **19** (2005), no. 1, 69–82. MR2178184
- 752 2. Marwane Bouznif, Frédéric Havet, and Myriam Preissmann, *Minimum-density identifying*  
753 *codes in square grids*, Lecture Notes in Computer Science (2016), 77–88.

- 754 3. Irène Charon, Iiro Honkala, Olivier Hudry, and Antoine Lobstein, *General bounds for identifying codes in some infinite regular graphs*, Electron. J. Combin. **8** (2001), no. 1, Research Paper 39, 21. MR1877658
- 755
- 756
- 757 4. ———, *The minimum density of an identifying code in the king lattice*, Discrete Math. **276**
- 758 (2004), no. 1-3, 95–109. MR2046627
- 759 5. Irène Charon, Olivier Hudry, and Antoine Lobstein, *Identifying codes with small radius in some infinite regular graphs*, Electron. J. Combin. **9** (2002), no. 1, Research Paper 11, 25. MR1912793
- 760
- 761
- 762 6. Gérard D. Cohen, Iiro Honkala, Antoine Lobstein, and Gilles Zémor, *Bounds for codes identifying vertices in the hexagonal grid*, SIAM J. Discrete Math. **13** (2000), no. 4, 492–504. MR1787935
- 763
- 764
- 765 7. ———, *On identifying codes*, Codes and association schemes (Piscataway, NJ, 1999), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 56, Amer. Math. Soc., Providence, RI, 2001, pp. 97–109. MR1816390
- 766
- 767
- 768 8. Daniel W. Cranston and Douglas B. West, *An introduction to the discharging method via graph coloring*, Discrete Math. **340** (2017), no. 4, 766–793. MR3603558
- 769
- 770 9. Daniel W. Cranston and Gexin Yu, *A new lower bound on the density of vertex identifying codes for the infinite hexagonal grid*, Electron. J. Combin. **16** (2009), no. 1, Research Paper 113, 16. MR2546316
- 771
- 772
- 773 10. Ari Cukierman and Gexin Yu, *New bounds on the minimum density of an identifying code for the infinite hexagonal grid*, Discrete Appl. Math. **161** (2013), no. 18, 2910–2924. MR3126657
- 774
- 775 11. Marc Daniel, Sylvain Gravier, and Julien Moncel, *Identifying codes in some subgraphs of the square lattice*, Theoretical Computer Science **319** (2004), no. 1, 411–421.
- 776
- 777 12. Rennan Dantas, Frédéric Havet, and Rudini M. Sampaio, *Identifying codes for infinite triangular grids with a finite number of rows*, Discrete Math. **340** (2017), no. 7, 1584–1597. MR3634129
- 778
- 779
- 780 13. ———, *Minimum density of identifying codes of king grids*, Discrete Math. **341** (2018), no. 10, 2708–2719. MR3843256
- 781
- 782 14. Florent Foucaud, *Combinatorial and algorithmic aspects of identifying codes in graphs*, Ph.D. thesis, Université Sciences et Technologies, 2012.
- 783
- 784 15. Mark Hartmann and James B. Orlin, *Finding minimum cost to time ratio cycles with small integral transit times*, Networks **23** (1993), no. 6, 567–574.
- 785
- 786 16. Iiro Honkala and Tero Laihonen, *Codes for identification in the king lattice*, Graphs Combin. **19** (2003), no. 4, 505–516. MR2031005
- 787
- 788 17. ———, *On the identification of sets of points in the square lattice*, Discrete Comput. Geom. **29** (2003), no. 1, 139–152. MR1946799
- 789
- 790 18. ———, *On identification in the triangular grid*, J. Combin. Theory Ser. B **91** (2004), no. 1, 67–86. MR2047532
- 791
- 792 19. ———, *On identifying codes in the hexagonal mesh*, Inform. Process. Lett. **89** (2004), no. 1, 9–14. MR2025883
- 793
- 794 20. ———, *On identifying codes in the triangular and square grids*, SIAM J. Comput. **33** (2004), no. 2, 304–312. MR2048443
- 795
- 796 21. Iiro Honkala and Antoine Lobstein, *On the density of identifying codes in the square lattice*, J. Combin. Theory Ser. B **85** (2002), no. 2, 297–306. MR1912970
- 797
- 798 22. Devin Jean, *Watching systems, identifying, locating-dominating and discriminating codes in graphs*, <https://dragazo.github.io/bibdom/main.pdf>, July 2023, Accessed on July 2023.
- 799
- 800 23. Devin Jean and Suk Seo, *Fault-tolerant locating-dominating sets with error-correction*, arXiv cs.DM (2022), 2212.08193 version 1.
- 801
- 802 24. Minghui Jiang, *Periodicity of identifying codes in strips*, Inform. Process. Lett. **135** (2018), 77–84. MR3779981
- 803
- 804 25. Ville Junnila and Tero Laihonen, *Optimal lower bound for 2-identifying codes in the hexagonal grid*, Electron. J. Combin. **19** (2012), no. 2, Paper 38, 16. MR2946096
- 805
- 806 26. Richard M. Karp, *A characterization of the minimum cycle mean in a digraph*, Discrete Mathematics **23** (1978), no. 3, 309–311.
- 807

- 808 27. Mark G. Karpovsky, Krishnendu Chakrabarty, and Lev B. Levitin, *On a new class of codes*  
809 *for identifying vertices in graphs*, IEEE Trans. Inform. Theory **44** (1998), no. 2, 599–611.  
810 MR1607726
- 811 28. Ryan Martin and Brendon Stanton, *Lower bounds for identifying codes in some infinite*  
812 *grids*, Electron. J. Combin. **17** (2010), no. 1, Research Paper 122, 16. MR2729371
- 813 29. Ville Salo and Ilkka Törmä, *Finding codes on infinite grids automatically*, 2023.
- 814 30. Rudini Sampaio, Gabriel Sobral, and Yoshiko Wakabayashi, *Minimum density of identifying*  
815 *codes of hexagonal grids with a finite number of rows*, Anais do VII Encontro de Teoria da  
816 Computação (Porto Alegre, RS, Brasil), SBC, 2022, pp. 145–148.
- 817 31. Peter J. Slater, *Fault-tolerant locating-dominating sets*, vol. 249, 2002, Combinatorics, graph  
818 theory and computing (Louisville, KY, 1999), pp. 179–189. MR1898670
- 819 32. Gabriel Sobral, Rudini Sampaio, and Yoshiko Wakabayashi, *Identifying codes*, [https://](https://gitlab.uspdigital.usp.br/gagsobral/identifying-codes)  
820 [gitlab.uspdigital.usp.br/gagsobral/identifying-codes](https://gitlab.uspdigital.usp.br/gagsobral/identifying-codes), 2023.
- 821 33. Derrick Stolee, *Automated discharging arguments for density problems in grids*, 2014.
- 822 34. Zhibin Xiao and Bevan Baas, *A hexagonal processor and interconnect topology for many-*  
823 *core architecture with dense on-chip networks*, VLSI-SoC: From Algorithms to Circuits and  
824 System-on-Chip Design, Springer Berlin Heidelberg, 2013, pp. 125–143.
- 825 35. Janez Zerovnik and Ignasi Sau, *An optimal permutation routing algorithm on full-duplex*  
826 *hexagonal networks*, Discrete Mathematics & Theoretical Computer Science **10** (2008), 1–  
827 15.