

ON BIORTHOGONAL SYSTEMS WHOSE FUNCTIONALS ARE FINITELY SUPPORTED

CHRISTINA BRECH AND PIOTR KOSZMIDER

ABSTRACT. We show that for each natural $n > 1$ it is consistent that there is a compact Hausdorff space K_{2n} such that in $C(K_{2n})$ there is no uncountable (semi)biorthogonal sequence $(f_\xi, \mu_\xi)_{\xi \in \omega_1}$ where μ_ξ 's are atomic measures with supports consisting of at most $2n - 1$ points of K_{2n} , but there are biorthogonal systems $(f_\xi, \mu_\xi)_{\xi \in \omega_1}$ where μ_ξ 's are atomic measures with supports consisting of $2n$ points. This complements a result of Todorćević that it is consistent that each nonseparable Banach space $C(K)$ has an uncountable biorthogonal system where the functionals are measures of the form $\delta_{x_\xi} - \delta_{y_\xi}$ for $\xi < \omega_1$ and $x_\xi, y_\xi \in K$. It also follows that it is consistent that the irredundance of the Boolean algebra $Clop(K)$ or the Banach algebra $C(K)$ for K totally disconnected can be strictly smaller than the sizes of biorthogonal systems in $C(K)$. The compact spaces exhibit an interesting behaviour with respect to known cardinal functions: the hereditary density of the powers K_{2n}^k is countable up to $k = n$ and it is uncountable (even the spread is uncountable) for $k > n$.

1. INTRODUCTION

If X is a Banach space and X^* its dual, then $(x_i, x_i^*)_{i \in I} \subseteq X \times X^*$ is called a biorthogonal system if $x_i^*(x_i) = 1$ and $x_i^*(x_j) = 0$ if $i \neq j$ for each $i, j \in I$. If α is an ordinal, a transfinite sequence $(x_i, x_i^*)_{i < \alpha} \subseteq X \times X^*$ is called a semibiorthogonal sequence if $x_i^*(x_i) = 1$, $x_i^*(x_j) = 0$ if $j < i < \alpha$ and $x_i^*(x_j) \geq 0$ if $i < j < \alpha$.

Biorthogonal systems have always played an important role in the theory of Banach spaces ([9]) because all kinds of bases in Banach spaces are in particular X -parts of biorthogonal systems ([20] and [21]). Semibiorthogonal sequences have been introduced quite recently ([2]) in the relation with the sets in Banach spaces supported by all of their points ([17], [13], [8]).

We will mainly deal with biorthogonal systems in Banach spaces $C(K)$ of all continuous functions on a compact Hausdorff space K with the supremum norm. Its dual space is isometric to the Banach space $M(K)$ of all Radon measures on K with the variation norm, and so, we will identify this dual with $M(K)$. If K is a compact Hausdorff space and $x \in K$, δ_x denotes the functional on $C(K)$ defined by $\delta_x(f) = f(x)$ for all $f \in C(K)$.

The first author was supported by FAPESP fellowship (2007/08213-2), which is part of Thematic Project FAPESP (2006/02378-7). Part of the research was done at the Technical University of Łódź where the first author was partially supported by Polish Ministry of Science and Higher Education research grant N N201 386234.

The second author was partially supported by Polish Ministry of Science and Higher Education research grant N N201 386234. Part of the research was done at the State University of Campinas UNICAMP where the second author was partially supported by the Department of Mathematics.

This paper is motivated by the following question: *If there is an uncountable biorthogonal system $(x_\xi, x_\xi^*)_{\xi \in \omega_1}$ in $C(K) \times M(K)$, is there also one such that*

$$x_\xi^* = \delta_{x_\xi} - \delta_{y_\xi}$$

for some points $x_\xi, y_\xi \in K$? We will follow [4] and call such a biorthogonal system a nice biorthogonal system.

The origin of this question is that in all concrete situations so far analyzed in the literature the above question has positive answer. Moreover, it happens for a good reason, namely, it follows from a recent result of Todorćević that Martin's axiom together with the negation of the continuum hypothesis implies the positive answer to this question. Indeed, analyzing the proof of Theorem 11 of [24], one gets two cases: the first when K is hereditarily separable, which is the main part of that proof and the constructed biorthogonal system is nice; and the second case, when K is c.c.c. but contains a nonseparable subspace, then the proof of Theorem 10 of [24] provides the required nice system; if K is not c.c.c., one can easily obtain an uncountable nice biorthogonal system.

There is one more reason why nice biorthogonal systems appear frequently in the context of Banach spaces $C(K)$ and which makes them more meaningful. Namely, $(f_\alpha)_{\alpha \in \kappa}$ is the X -part of a nice biorthogonal system if and only if $(f_\alpha)_{\alpha \in \kappa}$ is irredundant in the Banach algebra $C(K)$, in the sense that no f_α belongs to the Banach subalgebra generated by the remaining elements. This is a consequence of the Stone-Weierstrass theorem. If K is totally disconnected and f_α 's are characteristic functions of clopen $A_\alpha \subseteq K$, we obtain a well-known notion of an irredundant set in a Boolean algebra (see e.g., [14]), i.e., a set where no element belongs to the Boolean algebra generated by the remaining elements. The irredundance of a Boolean algebra is the supremum of cardinalities of irredundant sets.

To formulate properly our main results we need the following:

Definition 1.1. *Let K be a compact Hausdorff space and $n \in \mathbb{N}$. We say that the functionals of a sequence $(f_\xi, \mu_\xi)_{\xi \in \omega_1} \subseteq C(K) \times M(K)$ are n -supported if each μ_ξ is an atomic measure whose support consists of no more than n points of K .*

Theorem 1.2. *For each natural $n > 1$, it is consistent that there is a compact Hausdorff space K_{2n} such that in $C(K_{2n})$ there is no uncountable semibiorthogonal sequence whose functionals are $2n-1$ -supported, but there are biorthogonal systems whose functionals are $2n$ -supported.*

Moreover, K_{2n}^n is hereditarily separable but K_{2n}^{n+1} has an uncountable discrete subspace. Neither the Banach algebra $C(K_{2n})$ nor the Boolean algebra $\text{Clop}(K_{2n})$ have an uncountable irredundant family. In particular, $C(K_4)$ has an uncountable biorthogonal system but it has no uncountable nice biorthogonal system.

Such a situation suggests many questions about the size of biorthogonal systems of various types in $C(K)$ spaces as well as in general Banach spaces. These more general discussions will appear elsewhere. In particular, we are unable to obtain K 's such that $C(K)$ contains biorthogonal systems whose functionals are $2n+1$ -supported but does not contain one whose functionals are $2n$ -supported. The reason why some fundamental change with the approach must be taken to obtain such a space is shown in Lemma 3.9.

On the other hand, if $n = 1$ one has absolute results. If K is the split interval, then K is hereditarily separable and so cannot have an uncountable semibiorthogonal system whose functionals are 1-supported, but $C(K)$ has an uncountable biorthogonal system (see [7]).

It seems that our compact space is the first example showing that the hereditary density or spread of finite powers of a compact space may change its value from countable to uncountable arbitrarily high in \mathbb{N} . Such an example can be only consistent since, for example, under $\text{MA} + \neg\text{CH}$ if K^3 is hereditarily separable for a compact K , then it is metrizable, and so, all finite powers are hereditarily separable. This follows from the fact that then there are no compact S -spaces ([22]), from the Katetov theorem ([10]) and from the fact that Lindelöf regular spaces are normal.

The paper is organized as follows: in the following second section we discuss a general form of the compact spaces we construct and call them unordered N -split Cantor sets. They are versions of the split interval whose relation with biorthogonal systems in Banach spaces was already demonstrated in [7]. Section 3 is devoted to a generic construction of Boolean algebras whose Stone spaces are the K_{2^n} 's. This is the only section that requires the knowledge of forcing. The partial order used is a new modification of that of [11], which produced nonseparable $C(K)$'s with no uncountable semibiorthogonal sequences. Thus our spaces are quite controllable members of the group of compact spaces constructed in [1], [18], [19], [11]. In this section we also prove the existence of an uncountable discrete subspace of $K_{2^n}^{n+1}$ and an uncountable biorthogonal system in $C(K)$ whose functionals are $2n$ -supported. The section ends with Theorem 3.8, which expresses the random character of the constructed compact space. Later on we use this theorem to prove further properties of the space. Hence a reader not familiar with forcing may use this theorem for other purposes and read only the following section.

The last, fourth section is devoted to applications of Theorem 3.8, that is the proof that $K_{2^n}^n$ is hereditarily separable and that $C(K_{2^n})$ has no uncountable semibiorthogonal sequences whose functionals are $2n - 1$ -supported.

We use standard notation. In particular $[n] = \{1, \dots, n\}$ and $n = \{0, \dots, n - 1\}$ for a positive natural number n . A^B denotes the set of all functions from B into A and so if $2 = \{0, 1\}$ we have that 2^ω denotes all infinite sequences with terms in $\{0, 1\}$, while 2^n stands for functions from n into $\{0, 1\}$; also $2^{<\omega} = \bigcup \{2^n : n \in \mathbb{N}\}$. $\langle s \rangle = \{x \in 2^\omega : s \subseteq x\}$ for $s \in 2^n$ for some $n \in \mathbb{N}$. If A, B are sets of ordinals, then $A < B$ means that $\alpha < \beta$ for any $\alpha \in A$ and any $\beta \in B$.

2. UNORDERED N -SPLIT CANTOR SETS

Fix a sequence of distinct elements $\mathcal{X} = \{x_\xi : \xi < \omega_1\} \subseteq 2^\omega$ and $N \in \mathbb{N}$. Let

$$K_N = (2^\omega \setminus \mathcal{X}) \cup (\mathcal{X} \times [N])$$

and define

$$V_s = (\langle s \rangle \cap (2^\omega \setminus \mathcal{X})) \cup ((\langle s \rangle \cap \mathcal{X}) \times [N]).$$

Definition 2.1. A family $(A_{\xi,i} : \xi < \omega_1, i \in [N])$ of subsets of K_N is called an N -splitting family if it satisfies the following conditions:

- (1) $(x_\xi, i) \in A_{\xi,i} \subseteq K_N$ for each $\xi < \omega_1$ and $i \in [N]$;
- (2) for each $\xi < \omega_1$ the sets $A_{\xi,i}$'s are pairwise disjoint;
- (3) for each $\xi < \omega_1$ we have $K_N = A_{\xi,1} \cup \dots \cup A_{\xi,N}$;
- (4) if $\eta < \xi$, then there is $k \in \mathbb{N}$ and $j \in [N]$ such that $A_{\eta,i} \cap V_{x_\eta|k} \subseteq A_{\xi,j} \cap V_{x_\eta|k}$;

(5) if $\eta > \xi$ and $x = x_\eta$ or $x \in 2^\omega \setminus \mathcal{X}$, then there is $k \in \mathbb{N}$ and $j \in [N]$ such that $V_{x|k} \subseteq A_{\xi,j}$.

Definition 2.2. Given an N -splitting family $(A_{\xi,i} : \xi < \omega_1, i \in [N])$, we call the space (K_N, \mathcal{T}) an unordered N -split Cantor set if the topology \mathcal{T} on K_N is defined by indicating neighbourhood bases \mathcal{B}_x at x for every $x \in K_N$ in the following way: if $x \in 2^\omega \setminus \mathcal{X}$, then

$$\mathcal{B}_x = \{V_s : s \subseteq x\}$$

and if $x = (x_\xi, j) \in K_N$, then

$$\mathcal{B}_x = \{V_s \cap A_{\xi,j} : s \subseteq x_\xi\}.$$

The intuitive meaning of the above definitions is as follows: each point x_ξ of 2^ω is split into N points $(x_\xi, 1), \dots, (x_\xi, N)$. If we view K_N as constructed inductively, when at step $\xi < \omega_1$ we construct the splitting clopen neighbourhoods $A_{\xi,1}, \dots, A_{\xi,N}$ of the points $(x_\xi, 1), \dots, (x_\xi, N)$, then these neighbourhoods split only x_ξ and no other previously constructed (x_η, i) for $\eta < \xi$ (condition 2.1.(4)) nor x_η for $\eta > \xi$ nor $x \in 2^\omega \setminus \mathcal{X}$ (condition 2.1.(5)). Note that, on the other hand, $A_{\xi,i}$'s may split x_η for $\eta < \xi$, and in this case, by condition 2.1.(4), they do it "the same way" as the $A_{\eta,j}$'s.

Proposition 2.3. Let $N \in \mathbb{N}$. If $(A_{\xi,i} : \xi < \omega_1, i \in [N])$ is an N -splitting family, then the corresponding unordered N -split Cantor set is a compact, Hausdorff, totally disconnected topological space.

Proof. Since $V_\emptyset = K_N$, conditions (1) - (3) of Definition 2.1 imply that $A_{\xi,i}$'s are clopen sets. Now using Proposition 1.2.3. of [5], we will prove that the above families satisfy the axioms for neighbourhood bases BP1-BP3 from [5]. The only nontrivial part is to prove that given $x \in V \in \mathcal{B}_y$, there is $U \in \mathcal{B}_x$ such that $x \in U \subseteq V$.

Suppose $x \in 2^\omega \setminus \mathcal{X}$ and $x \in V_s \in \mathcal{B}_y$. Then $s \subseteq x$ and so V_s itself is in \mathcal{B}_x . If $x \in V_s \cap A_{\xi,i}$, we also have $s \subseteq x$ and by (5) of Definition 2.1 there is $k \in \mathbb{N}$ such that $V_{x|k} \subseteq A_{\xi,j}$ for some $j \in [N]$. Put $t = s \cup x|k$ and note that we have that $V_t \subseteq A_{\xi,j}$, so by the disjointness (condition 2.1.(2)) we have $j = i$ with $x \in V_t \in \mathcal{B}_x$ and $V_t \subseteq V_s \cap A_{\xi,i}$.

Now suppose that $x = (x_\eta, i)$ and $x \in V_s \in \mathcal{B}_y$, hence $s \subseteq x$ and so $V_s \cap A_{\eta,i} \in \mathcal{B}_x$ and $x \in V_s \cap A_{\eta,i} \subseteq V_s$.

Finally, let $x = (x_\eta, i)$ and $x \in V_s \cap A_{\xi,j} \in \mathcal{B}_{(x_\xi, j)}$, then $s \subseteq x_\eta$.

First consider $\eta < \xi$, then by (5) of Definition 2.1 there are $k \in \mathbb{N}$ and j' such that $A_{\eta,i} \cap V_{x_\eta|k} \subseteq A_{\xi,j'} \cap V_{x_\eta|k}$ and by disjointness (2) we get that $j' = j$. So, if we put $t = s \cup x_\eta|k$, then $A_{\eta,i} \cap V_t \subseteq A_{\xi,j} \cap V_t \subseteq A_{\xi,j} \cap V_s$ and of course $A_{\eta,i} \cap V_t \in \mathcal{B}_{(x_\eta, i)}$.

Secondly if $\eta \geq \xi$ and $(x_\eta, i) \in V_s \cap A_{\xi,j}$, we also have $s \subseteq x_\eta$ and by 2.1.(4) there are $k \in \mathbb{N}$ and j' such that $V_{x_\eta|k} \subseteq A_{\xi,j'}$ for some j' . By the disjointness we have $j = j'$. If $t = s \cup x_\eta|k$ we have that $V_t \subseteq A_{\xi,j}$, so $x \in V_t \in \mathcal{B}_x$ and $V_t \subseteq V_s \cap A_{\xi,i}$. This completes the proof that \mathcal{B}_x 's form a local neighbourhood system.

The Hausdorff property is easy since basic sets are clopen.

To prove the compactness, suppose \mathcal{U} is an open cover of K_N . We may assume that it consists of basic open sets. For each $x \in 2^\omega \setminus \mathcal{X}$ define $s_x \in 2^{<\omega}$ such that $x \in V_s \subseteq U \in \mathcal{U}$ for some U , and for each $\xi < \omega_1$ define $s_\xi \in 2^{<\omega}$ such that $(x_\xi, i) \in V_{s_\xi} \cap A_{\xi,i} \subseteq U \in \mathcal{U}$ for some U , and for each $1 \leq i \leq N$. This actually gives by (3) of Definition 2.1 that V_{s_ξ} is covered by finitely many $U \in \mathcal{U}$.

Now $\{\langle s_x \rangle, \langle s_\xi \rangle : x \in 2^\omega \setminus \mathcal{X}, \xi < \omega_1\}$ forms an open cover of 2^ω which is compact and so it has a finite subcover, which easily yields a finite subcover of \mathcal{U} . \square

Definition 2.4. Suppose $N \in \mathbb{N}$ and K_N is an unordered N -split Cantor set. Under the notation as above, we define the following:

- $R_\xi = \{(x_\xi, 1), \dots, (x_\xi, N)\}$,
- \mathcal{A}_α is the subalgebra of $\text{Clop}(K_N)$ generated by $(V_s : s \in 2^{<\omega})$ and $\{A_{\xi,i} : \xi < \alpha, i \in [N]\}$ for $\alpha \leq \omega_1$.
- C_α is the closure (in the norm) of finite linear combinations of characteristic functions of elements of \mathcal{A}_α inside $C(K)$.

Note that C_0 can be naturally identified with $C(2^\omega)$ inside $C(K)$.

Lemma 2.5. Let $N \in \mathbb{N}$ and let K_N be an unordered N -split Cantor set. For every $n \in \mathbb{N}$ and for every $\alpha \in \omega_1$ and every $i \in [N]$ we have

$$A_{\alpha,i} \setminus V_{x_\alpha|n} \in \mathcal{A}_\alpha.$$

Proof. By the properties 2.1.(4) and (5) of $A_{\xi,i}$'s any point of $K_N \setminus R_\alpha$ has a neighbourhood V such that for every $i \in [N]$ it is included in $A_{\alpha,i}$ or disjoint from $A_{\alpha,i}$ and moreover $V \in \mathcal{A}_\alpha$.

Since $A_{\alpha,i} \setminus V_{x_\alpha|n}$ is a compact subspace of $K_N \setminus R_\alpha$, we have a finite subcover consisting of subsets i.e., $A_{\alpha,i} \setminus V_{x_\alpha|n}$ is the supremum of a finite family of elements of \mathcal{A}_α as required. \square

Let us see the general form of continuous rational simple functions on an unordered N -split Cantor set. By a rational simple function we mean a function assuming only finitely many rational values.

Lemma 2.6. Suppose that $N \in \mathbb{N}$ and that K_N is an unordered N -split Cantor set, $\varepsilon > 0$, μ is a (regular) Radon measure on K_N and that f is a continuous rational simple function on K_N . Then there is a simple rational function $g \in C(2^\omega)$, distinct $\xi_1, \dots, \xi_k < \omega_1$ and there are rationals $q_{i,l}$, non-negative integers m_i and $s_i \in 2^{m_i}$ such that $s_i = x_{\xi_i}|m_i$ for $1 \leq i \leq k$ and for $1 \leq l < N$ such that

$$f = g + \sum_{1 \leq i \leq k} \sum_{1 \leq l < N} q_{i,l} \chi_{A_{\xi_i,l} \cap V_{s_i}}$$

and such that

$$\sum_{1 \leq i \leq k} \max_{1 \leq l < N} (|q_{i,l}|) |\mu|(V_{s_i} \setminus R_{\xi_i}) \leq \varepsilon.$$

Proof. By induction on ξ we prove that any continuous simple rational function in C_ξ can be written in the form as in the lemma. The Stone-Weierstrass theorem and the uncountable cofinality of ω_1 imply that the union of C_ξ 's is the entire $C(K_N)$.

The limit stage is trivial. So, suppose we are done for C_ξ and we are given a continuous simple rational function f in $C_{\xi+1}$. Note that

$$\bigcap_{m \in \mathbb{N}} V_{x_\xi|m} = R_\xi.$$

Hence, the regularity of the Radon measures implies that $|\mu|(V_{x_\xi|m} \setminus R_\xi)$'s converge to 0. Let m_1 be such that

$$|\mu|(V_{x_\xi|m} \setminus R_\xi) \leq \frac{\varepsilon}{4\|f\|}$$

for $m \geq m_1$.

Note also that a simple function is a linear combination of characteristic functions of clopen sets, hence there are $\xi_1, \dots, \xi_{k-1} < \xi < \omega_1$ and m_2 such that preimages under f of each of its finite rational values belong to the subalgebra of $\mathcal{A}_{\xi+1}$ generated by V_t 's for $|t| < m_2$ and $A_{\xi_1, j}, \dots, A_{\xi_{k-1}, j}, A_{\xi, j}$ for $1 \leq j \leq N$. Now let $n \geq m_1, m_2$ be such that for every $1 \leq i < k$ there is $1 \leq j \leq N$ such that $V_{x_\xi | m} \subseteq A_{\xi_i, j}$ which can be obtained by the property (5) (of Definition 2.1) of A_ξ 's and $\eta = \xi_i$.

It follows that f is constant on $A_{\xi, j} \cap V_{x_\xi | m}$ for every $1 \leq j \leq N$. Let $q'_1, \dots, q'_N \in \mathbb{Q}$ be the corresponding values and note that $|q'_l - q'_N| \leq 2\|f\|$ for any $1 \leq l \leq N$. So, by conditions (2) and (3) (of Definition 2.1) of $A_{\xi, j}$'s we have

$$f = [f|(K \setminus V_{x_\xi | m}) + q'_N \chi_{V_{x_\xi | m}}] + \sum_{1 \leq l < N} (q'_l - q'_N) \chi_{A_{\xi, l} \cap V_{x_\xi | m}}.$$

Note that $f|(K \setminus V_{x_\xi | m})$ belongs to C_ξ by Lemma 2.5, and so

$$f = h + \sum_{1 \leq l < N} q_l \chi_{A_{\xi, l} \cap V_{x_\xi | m}}, \quad \max_{1 \leq l < N} |q_l| \mu(V_{x_\xi | m} \setminus R_\xi) \leq \frac{\varepsilon}{2}$$

where $q_l = q'_l - q'_N$ and $h \in C_\xi$. Hence the inductive assumption for $\varepsilon/2$ can be used, which completes the proof of the lemma. \square

Definition 2.7. We say that an N -splitting family $(A_{\xi, i} : \xi < \omega_1, i \in [N])$ is balanced if it satisfies the following additional condition:

(6) for all distinct $\xi, \eta \in \omega_1$ and all $j \in [2n]$,

$$|\{i \in \{1, 3, \dots, 2n-1\} : (x_\eta, i) \in A_{\xi, j}\}| = |\{i \in \{2, 4, \dots, 2n\} : (x_\eta, i) \in A_{\xi, j}\}|.$$

Lemma 2.8. Suppose that $n \in \mathbb{N}$ and that K_{2n} is an unordered $2n$ -split Cantor set, where the N -splitting family $(A_{\xi, i} : \xi < \omega_1, i \in [2n])$ is balanced. Then we have that:

- (a) K_{2n}^{n+1} contains an uncountable discrete subspace;
- (b) there is an uncountable biorthogonal system in $C(K_{2n})$ with $2n$ -supported functionals.

Proof. To prove (a), let us show that the subset $\{((x_\xi, 1), (x_\xi, 2), (x_\xi, 4), \dots, (x_\xi, 2n)) : \xi < \omega_1\}$ of K_{2n}^{n+1} is relatively discrete.

Let $U_\xi = A_{\xi, 1} \times A_{\xi, 2} \times A_{\xi, 4} \times \dots \times A_{\xi, 2n}$, which clearly is an open neighbourhood of $((x_\xi, 1), (x_\xi, 2), (x_\xi, 4), \dots, (x_\xi, 2n))$. Now, fix distinct $\xi, \eta < \omega_1$ and let us prove that $((x_\eta, 1), (x_\eta, 2), (x_\eta, 4), \dots, (x_\eta, 2n)) \notin U_\xi$.

By contradiction, suppose that $((x_\eta, 1), (x_\eta, 2), (x_\eta, 4), \dots, (x_\eta, 2n)) \in U_\xi$, that is, $(x_\eta, j) \in A_{\xi, j}$ for each $j = 1, 2, 4, \dots, 2n$. By condition 2.7.(6), we have that for each $j \in [2n]$,

$$|\{i \in \{1, 3, \dots, 2n-1\} : (x_\eta, i) \in A_{\xi, j}\}| = |\{i \in \{2, 4, \dots, 2n\} : (x_\eta, i) \in A_{\xi, j}\}|.$$

Hence, each set $A_{\xi, 2}, A_{\xi, 4}, \dots, A_{\xi, 2n}$ must contain at least one of the $(x_\eta, 1), (x_\eta, 3), \dots, (x_\eta, 2n-1)$. By the disjointness of the $A_{\xi, j}$'s (Property (2) of Definition 2.1), $(x_\eta, 1)$ has to be in one of the sets $A_{\xi, 2}, A_{\xi, 4}, \dots, A_{\xi, 2n}$. But by our assumption, $(x_\eta, 1) \in A_{\xi, 1}$ and again by the disjointness of the $A_{\xi, j}$'s, this is a contradiction.

To show (b), for each $\xi < \omega_1$, let $f_\xi = \chi_{A_{\xi, 2n}}$ and

$$\mu_\xi = \sum_{k=1}^n (\delta_{(x_\xi, 2i)} - \delta_{(x_\xi, 2i-1)})$$

and note that $(f_\xi, \mu_\xi)_{\xi < \omega_1} \subseteq C(K_{2n}) \times M(K_{2n})$. Let us prove that this is a biorthogonal system.

For each $\xi < \omega_1$, since $(x_\xi, i) \in A_{\xi, i}$ and these sets are disjoint (Property (2) of Definition 2.1), we get that

$$\begin{aligned} \mu_\xi(f_\xi) &= \sum_{k=1}^n (\delta_{(x_\xi, 2k)} - \delta_{(x_\xi, 2k-1)})(\chi_{A_{\xi, 2n}}) \\ &= \sum_{k=1}^n (\chi_{A_{\xi, 2n}}((x_\xi, 2k)) - \chi_{A_{\xi, 2n}}((x_\xi, 2k-1))) = \chi_{A_{\xi, 2n}}((x_\xi, 2n)) = 1. \end{aligned}$$

On the other hand, for distinct $\xi, \eta < \omega_1$, by Property (6), we have that for all $j \in [2n]$,

$$|\{i \in \{1, 3, \dots, 2n-1\} : (x_\eta, i) \in A_{\xi, j}\}| = |\{i \in \{2, 4, \dots, 2n\} : (x_\eta, i) \in A_{\xi, j}\}|.$$

Hence,

$$\begin{aligned} \mu_\xi(f_\eta) &= \sum_{k=1}^n (\delta_{(x_\xi, 2k)} - \delta_{(x_\xi, 2k-1)})(\chi_{A_{\eta, 2n}}) \\ &= \sum_{k=1}^n (\chi_{A_{\eta, 2n}}((x_\xi, 2k)) - \chi_{A_{\eta, 2n}}((x_\xi, 2k-1))) \\ &= \sum_{k=1}^n \chi_{A_{\eta, 2n}}((x_\xi, 2k)) - \sum_{k=1}^n \chi_{A_{\eta, 2n}}((x_\xi, 2k-1)) \\ &= |\{i \in \{2, 4, \dots, 2n\} : (x_\xi, i) \in A_{\eta, 2n}\}| - |\{i \in \{1, 3, \dots, 2n-1\} : (x_\xi, i) \in A_{\eta, 2n}\}| \\ &= 0, \end{aligned}$$

concluding that $(f_\xi, \mu_\xi)_{\xi < \omega_1} \subseteq C(K_{2n}) \times M(K_{2n})$ is a biorthogonal system. \square

3. THE GENERIC CONSTRUCTION

This section is devoted to a generic construction of an unordered $2n$ -split Cantor set which exhibits quite random features. This type of uncountable structures was first investigated systematically in [19]. One can describe this random behavior as: in any uncountable sequence of finite substructures we have two which are related as we wish (up to constrains). We fix an uncountable sequence $(x_\xi : \xi < \omega_1) \subseteq 2^\omega$ consisting of distinct elements.

Definition 3.1. *Let \mathbb{P} be the forcing formed by conditions*

$$p = (F_p, n_p, (f_\xi^p : \xi \in F_p)),$$

where:

1. $F_p \in [\omega_1]^{<\omega}$;
2. $n_p \in \omega$ is such that for all $\xi \neq \eta$ in F_p , $x_\xi|n_p \neq x_\eta|n_p$;
3. for all $\xi \in F_p$,

$$f_\xi^p : 2^{n_p} \setminus \{x_\xi|n_p\} \rightarrow [2n]^{[2n]} \times [F_p \cap (\xi + 1)]$$

is such that

- a) if $f_\xi^p(s) = (\varphi, \xi)$, then φ is a constant function;
- b) if $f_\xi^p(s) = (\varphi, \eta)$ for some $\eta < \xi$, then

$$\forall j \in [2n] \quad |\varphi^{-1}(j) \cap \{1, 3, 5, \dots, 2n-1\}| = |\varphi^{-1}(j) \cap \{2, 4, \dots, 2n\}|.$$

We put $q \leq p$ if $F_q \supseteq F_p$, $n_q \geq n_p$ and for all $\xi \in F_p$, all $s \in 2^{n_q} \setminus \{x_\xi|n_q\}$ and all $t \in 2^{n_p} \setminus \{x_\xi|n_p\}$,

$$t \subseteq s \Rightarrow f_\xi^p(t) = f_\xi^q(s).$$

Intuitively, we are, of course, trying to build a $2n$ -split Cantor set which is determined by the choice of the balanced $2n$ -splitting family formed by $A_{\xi,i}$'s. Thus the coordinate $f_\xi^p(s)$ describes the behavior of $A_{\xi,i}$'s on V_s . The formal description is the subject of Definition 3.3. The value $f_\xi^p(s) = (\varphi, \xi)$, where φ has to be a constant function, say equal to i , means that the entire V_s is included in $A_{\xi,i}$. The value $f_\xi^p = (\varphi, \eta)$ for some $\eta < \xi$ means that $A_{\xi,i}$'s divide V_s as coded by φ i.e., $A_{\eta,j} \cap V_s \subseteq A_{\xi,\varphi(j)}$ for each $j \in [N]$. Note that a condition $p \in \mathbb{P}$ does not carry any information about the behavior of $A_{\xi,i}$'s on $V_{x_\xi|n_p}$, other than $(x_\xi, i) \in A_{\xi,i}$. This is the degree of freedom we have and which can be controlled by passing to an appropriate extension $q \leq p$. Condition (b) is to guarantee that the family of $A_{\xi,i}$'s is balanced, that is, that it satisfies property (6) of Definition 2.7.

Lemma 3.2. *The following subsets of \mathbb{P} are dense in \mathbb{P} :*

- (i) $\{p \in \mathbb{P} : n_p \geq k\}$, for some fixed $k \in \mathbb{N}$;
- (ii) $\{p \in \mathbb{P} : \xi \in F_p\}$, for some fixed $\xi < \omega_1$.

Proof. For (i), fix $k \in \mathbb{N}$ and let $p = (F_p, n_p, (f_\xi^p : \xi \in F_p)) \in \mathbb{P}$. If $n_p < k$, define $q = (F_q, n_q, (f_\xi^q : \xi \in F_q))$ by putting $F_q = F_p$, $n_q = k$ and for each $\xi \in F_q = F_p$, f_ξ^q is any function satisfying condition 3 of the definition of the forcing such that $f_\xi^q(t) = f_\xi^p(t|n_p)$, if $t|n_p \in 2^{n_p} \setminus \{x_\xi|n_p\}$; for example, let

$$f_\xi^q(t) = \begin{cases} f_\xi^p(t|n_p) & \text{if } t|n_p \in 2^{n_p} \setminus \{x_\xi|n_p\}, \\ (\varphi, \xi) & \text{otherwise,} \end{cases}$$

where φ is the constant function equal to 1. It is easy to see that $q \in \mathbb{P}$ and $q \leq p$.

For (ii), fix $\xi < \omega_1$ and let $p = (F_p, n_p, (f_\eta^p : \eta \in F_p)) \in \mathbb{P}$. By (i), we may assume that n_p is such that $x_\eta|n_p \neq x_\xi|n_p$ for all $\eta \in F_p$. Define $q = (F_q, n_q, (f_\eta^q : \eta \in F_q))$ by putting $F_q = F_p \cup \{\xi\}$, $n_q = n_p$, $f_\eta^q = f_\eta^p$ for each $\eta \in F_p$, and f_ξ^q is any function satisfying condition 3 of the definition of the forcing; for example, let $f_\xi^q(t) = (\varphi, \xi)$, where φ is the constant function equal to 1. It is easy to see that $q \in \mathbb{P}$ and $q \leq p$. \square

Definition 3.3. *Given a \mathbb{P} -generic filter G over a model V , we define the family $\{A_{\xi,j} : \xi \in \omega_1, j \in [2n]\}$ as follows: for each $\xi \in \omega_1$ and each $j \in [2n]$, let*

$$A_{\xi,j} = \{(x_\xi, j)\} \cup \bigcup \{V_s : \exists p \in G, f_\xi^p(s) = (\varphi, \xi) \text{ and } \varphi \text{ is the constant function equal to } j\} \\ \cup \bigcup \{V_s \cap A_{\eta,i} : \exists p \in G, f_\xi^p(s) = (\varphi, \eta), \text{ for some } \eta \neq \xi \text{ and } \varphi(i) = j\}.$$

The following lemma follows directly from the above definition.

Lemma 3.4. *Given $p \in G$, $\xi \in F_p$ and $s \in 2^{n_p} \setminus \{x_\xi|n_p\}$, we have that:*

- (a) if $f_\xi^p(s) = (\varphi, \xi)$, then $V_s \subseteq A_{\xi,j}$ for $j = \varphi(1)$;
- (b) if $f_\xi^p(s) = (\varphi, \eta)$ for some $\eta < \xi$, then $\forall i \in [2n]$, $V_s \cap A_{\eta,i} \subseteq A_{\xi,\varphi(i)}$. \square

Notice that in case $f_\xi^p(s) = (\varphi, \xi)$, φ is the constant function equal to j , so that we could have taken $j = \varphi(i)$ for any $i \in [2n]$.

Let us now check that the family $\{A_{\xi,j} : \xi \in \omega_1, j \in [2n]\}$ has the desired properties.

Theorem 3.5. *The family $\{A_{\xi,j} : \xi \in \omega_1, j \in [2n]\}$ is a balanced $2n$ -splitting family.*

Proof. Let us prove that the family satisfies conditions 2.1.(1) - (5) and 2.7.(6).

(1). It follows directly from the definition of $A_{\xi,j}$.

(2). We prove it by induction on ξ . First notice that by the definition of the forcing \mathbb{P} ,

$$\forall p \in \mathbb{P} \forall \xi \in F_p \forall s \in \text{dom} f_\xi^p \quad R_\xi \cap V_s = \emptyset,$$

since $x_\xi|n_p \notin \text{dom} f_\xi^p$. Then, $(x_\xi, j_1) \in A_{\xi,j_2}$ iff $j_1 = j_2$.

Now, fix $\xi < \omega_1$ and suppose we have that $A_{\eta,i}$ are pairwise disjoint for each fixed $\eta < \xi$. Suppose there is $x \in A_{\xi,j_1} \cap A_{\xi,j_2}$ for some distinct $j_1, j_2 \in [2n]$. By the above observation, $x \neq (x_\xi, j)$ for any $j \in [2n]$.

By the definition of A_{ξ,j_k} , for each $k \in \{1, 2\}$ there are $p_k \in G$ and $s_k \in \text{dom} f_\xi^{p_k}$ such that $x \in V_{s_k}$ and

$$\text{either } f_\xi^{p_k}(s_k) = (\varphi_k, \xi) \text{ and } \varphi_k \text{ is the constant function equal to } j_k$$

$$\text{or } f_\xi^{p_k}(s_k) = (\varphi_k, \eta_k) \text{ for some } \eta_k < \xi \text{ and } x \in A_{\eta_k, i} \text{ for some } i \in \varphi_k^{-1}(j_k).$$

Let $p \in G$ be such that $p \leq p_1, p_2$ and let $t \in 2^{n_p} \setminus \{x_\xi|n_p\}$ be such that $x \in V_t$. Then, $t \supseteq s_k$ since $x \in V_{s_k}$ and hence, by the definition of extension in \mathbb{P} , $f_\xi^{p_1}(s_1) = f_\xi^p(t) = f_\xi^{p_2}(s_2)$, so that $\varphi_1 = \varphi_2$.

Now, if $f_\xi^p(t) = (\varphi, \xi)$, this would mean that φ_1 and φ_2 are both constant equal to j_1 and j_2 , contradicting the hypothesis that $j_1 \neq j_2$. Otherwise, if $f_\xi^p(t) = (\varphi, \eta)$, for some $\eta < \xi$, we would get that $x \in A_{\eta, i_k}$ for some $i_k \in \varphi^{-1}(j_k)$. By the inductive hypothesis we get that $i_1 = i_2 \in \varphi^{-1}(j_1) \cap \varphi^{-1}(j_2)$, which implies that $j_1 = j_2$, contradicting again the hypothesis that those are distinct.

This concludes the proof that the family satisfies condition (2) of Definition 2.1.

(3). Again we prove it by induction on ξ . So, let $\xi < \omega_1$, suppose $K = A_{\eta,1} \cup \dots \cup A_{\eta,2n}$ for any $\eta < \xi$ and let $x \in K$.

If $x = (x_\xi, i)$ for some $i \in [2n]$, then $x \in A_{\xi,i}$ by definition.

By Lemma 3.2, let $p \in G$ be such that $x \in V_s$ for some $s \in 2^{n_p} \setminus \{x_\xi|n_p\}$.

If $f_\xi^p(s) = (\varphi, \xi)$, by Lemma 3.4.(a) we get that $V_s \subseteq A_{\xi, \varphi(1)}$, which guarantees that $x \in A_{\xi, \varphi(1)}$.

Otherwise, if $f_\xi^p(s) = (\varphi, \eta)$, for some $\eta < \xi$, by the inductive hypothesis, let $i \in [2n]$ be such that $x \in A_{\eta, i}$. Then, by Lemma 3.4.(b), $V_s \cap A_{\eta, i} \subseteq A_{\xi, \varphi(i)}$, which implies that $x \in A_{\xi, \varphi(i)}$ and concludes the proof of condition (3) of Definition 2.1.

(4). Fix $\eta < \xi < \omega_1$ and $i \in [2n]$. By Lemma 3.2, let $p \in G$ be such that $\xi, \eta \in F_p$ and $x_\eta|n_p \neq x_\xi|n_p$.

If $f_\xi^p(x_\eta|n_p) = (\varphi, \xi)$, by Lemma 3.4.(a) we get that $V_{x_\eta|n_p} \subseteq A_{\xi, \varphi(1)}$ (and in particular $V_{x_\eta|n_p} \cap A_{\eta, i} \subseteq V_{x_\eta|n_p} \cap A_{\xi, \varphi(1)}$).

If $f_\xi^p(x_\eta|n_p) = (\varphi, \eta)$, for some $\eta < \xi$, then, by Lemma 3.4.(b), $V_{x_\eta|n_p} \cap A_{\eta, i} \subseteq A_{\xi, \varphi(i)}$ (and in particular $V_{x_\eta|n_p} \cap A_{\eta, i} \subseteq V_{x_\eta|n_p} \cap A_{\xi, \varphi(i)}$) and we're done with condition (4) of Definition 2.1.

(5). Let us prove this by induction on $\xi < \omega_1$. Let $\xi < \omega_1$ and $x \in 2^\omega \setminus \{x_\eta : \eta \leq \xi\}$.

If $x = x_\eta$ for some $\eta > \xi$, by Lemma 3.2 there is $p \in G$ be such that $\xi, \eta \in F_p$. Otherwise, if $x \in 2^\omega \setminus \{x_\eta : \eta < \omega_1\}$, by Lemma 3.2 there is $p \in G$ be such that $\xi \in F_p$ and $x|n_p \neq x_\xi|n_p$. In both cases, put $s = x|n_p$.

If $f_\xi^p(s) = (\varphi, \xi)$, then, by Lemma 3.4.(a), $V_s \subseteq A_{\xi, \varphi(1)}$.

If $f_\xi^p(s) = (\varphi, \eta')$ for some $\eta' \in F_p \cap \xi$, by the inductive hypothesis, there is $k \in \mathbb{N}$ and $i \in [2n]$ such that $V_{x|k} \subseteq A_{\eta', i}$. By Lemma 3.2, let $q \in G$ be such that $q \leq p$ and $n_q \geq k$. Putting $t = x|n_q$, we get that $V_t \subseteq V_{x|k} \subseteq A_{\eta', i}$ and $f_\xi^q(t) = f_\xi^p(s) = (\varphi, \eta')$, since $t \supseteq s$. This implies by Lemma 3.4.(b) that $V_t = V_t \cap A_{\eta', i} \subseteq A_{\xi, \varphi(i)}$, which concludes the proof of condition (5) of Definition 2.1.

Hence, the family formed by $A_{\xi, i}$'s is a $2n$ -splitting family.

(6). Let us prove this by induction on $\xi < \omega_1$. So, fix $\xi < \omega_1$ and suppose we know that for all $\zeta < \xi$, all $\eta \neq \zeta$ and all $j \in [2n]$,

$$|\{i \in \{1, 3, \dots, 2n-1\} : (x_\eta, i) \in A_{\zeta, j}\}| = |\{i \in \{2, 4, \dots, 2n\} : (x_\eta, i) \in A_{\zeta, j}\}|.$$

Now, fix $\eta \neq \xi$. Let $p \in G$ be such that $\xi, \eta \in F_p$, so that $x_\eta|n_p \in \text{dom} f_\xi^p$.

If $f_\xi^p(x_\eta|n_p) = (\varphi, \xi)$, then, by Lemma 3.4.(a), $V_{x_\eta|n_p} \subseteq A_{\xi, \varphi(1)}$, which implies that $(x_\eta, i) \in A_{\xi, \varphi(1)}$ for all $i \in [2n]$. By the disjointness of the $A_{\xi, i}$'s (3) and condition (6) of Definition 2.7 holds both for $A_{\xi, \varphi(1)}$ (which contains all (x_η, i)) and for $A_{\xi, j}$, $j \neq \varphi(1)$ (which contain no (x_η, i)).

If $f_\xi^p(x_\eta|n_p) = (\varphi, \zeta)$ for some $\zeta < \xi$ in F_p , then for all $i \in [2n]$, $V_{x_\eta|n_p} \cap A_{\zeta, i} \subseteq A_{\xi, \varphi(i)}$. This means that each $A_{\xi, j}$ contains exactly those (x_η, k) which are in $A_{\zeta, i}$ for some $i \in \varphi^{-1}(j)$. In particular, we have that

$$\begin{aligned} & \{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\xi, j}\} \\ &= \{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\zeta, i} \text{ for some } i \in \varphi^{-1}(j)\} \\ &= \bigcup_{i \in \varphi^{-1}(j)} \{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\zeta, i}\} \end{aligned}$$

and

$$\begin{aligned} & \{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\xi, j}\} \\ &= \{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\zeta, i} \text{ for some } i \in \varphi^{-1}(j)\} \\ &= \bigcup_{i \in \varphi^{-1}(j)} \{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\zeta, i}\}. \end{aligned}$$

Let us now consider two cases:

If $\eta = \zeta$, since $(x_\eta, k) \in A_{\eta, k}$, we get that

$$\{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\xi, j}\} = \{k \in \{1, 3, \dots, 2n-1\} : k \in \varphi^{-1}(j)\}$$

and

$$\{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\xi, j}\} = \{k \in \{2, 4, \dots, 2n\} : k \in \varphi^{-1}(j)\}.$$

By property 3.b) of the definition of the partial ordering, the sets on the right-hand side of these two equalities have same size, which guarantees that

$$|\{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\xi, j}\}| = |\{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\xi, j}\}|,$$

concluding the proof in this case.

If $\eta \neq \zeta$, by the inductive hypothesis we know that for all $i \in [2n]$,

$$|\{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\zeta, i}\}| = |\{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\zeta, i}\}|.$$

Hence,

$$\begin{aligned} & |\{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\xi, j}\}| \\ &= \left| \bigcup_{i \in \varphi^{-1}(j)} \{k \in \{1, 3, \dots, 2n-1\} : (x_\eta, k) \in A_{\zeta, i}\} \right| \end{aligned}$$

$$\begin{aligned}
 &= \left| \bigcup_{i \in \varphi^{-1}(j)} \{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\zeta, i}\} \right| \\
 &= |\{k \in \{2, 4, \dots, 2n\} : (x_\eta, k) \in A_{\xi, j}\}|,
 \end{aligned}$$

which concludes the proof of condition (6) of Definition 2.7, that is, the family of $A_{\xi, i}$'s is a balanced $2n$ -splitting family. \square

Proposition 3.6. *Let $p_1 = (F_1, n_1, (f_\xi^1 : \xi \in F_1))$ and $p_2 = (F_2, n_2, (f_\xi^2 : \xi \in F_2))$ be conditions of \mathbb{P} such that:*

- $F_1 \cap F_2 < F_1 \setminus F_2 < F_2 \setminus F_1$;
- $n_1 = n_2 = n$;
- *there is an order-preserving bijection $e : F_1 \rightarrow F_2$ such that*
 - *for all $\xi \in F_1$, $x_\xi|n = x_{e(\xi)}|n$;*
 - *for all $\xi \in F_1$ and all $s \in 2^{n_1} \setminus \{x_\xi|n_1\} (= 2^{n_2} \setminus \{x_{e(\xi)}|n_2\})$,*

$$f_{e(\xi)}^2(s) = (\varphi, e(\eta)) \quad \text{where } f_\xi^1(s) = (\varphi, \eta).$$

Then, given $(\epsilon_\xi : \xi \in F_1 \setminus F_2) \subseteq [2n]^{[2n]}$ such that for all $\xi \in F_1 \setminus F_2$

$$\forall j \in [2n] \quad |\epsilon_\xi^{-1}(j) \cap \{1, 3, 5, \dots, 2n-1\}| = |\epsilon_\xi^{-1}(j) \cap \{2, 4, 6, \dots, 2n\}|$$

and given constant functions $(\delta_\xi : \xi \in F_1 \setminus F_2) \subseteq [2n]^{[2n]}$, there is $q \leq p_1, p_2$, $q \in \mathbb{P}$ such that

$$(3.1) \quad \forall \xi \in F_1 \setminus F_2 \quad f_\xi^q(x_{e(\xi)}|n_q) = (\delta_\xi, \xi) \quad \text{and} \quad f_{e(\xi)}^q(x_\xi|n_q) = (\epsilon_\xi, \xi).$$

Proof. Let $q = (F_q, n_q, (f_\xi^q : \xi \in F_q))$ be defined as follows: let $F_q = F_1 \cup F_2$; let $n_q \in \mathbb{N}$ be such $n_p \leq n_q$ and for all $\xi < \eta \in F_q$, $x_\xi|n_q \neq x_\eta|n_q$; for each $\xi \in F_q$ and $t \in 2^{n_q} \setminus \{x_\xi|n_q\}$, let

$$f_\xi^q(t) = \begin{cases} f_\xi^1(t|n) & \text{if } \xi \in F_1 \text{ and } t|n \neq x_\xi|n & \text{(Case 1.)} \\ f_\xi^2(t|n) & \text{if } \xi \in F_2 \text{ and } t|n \neq x_\xi|n & \text{(Case 2.)} \\ (\delta_\xi, \xi) & \text{if } \xi \in F_1 \text{ and } t|n = x_\xi|n & \text{(Case 3.)} \\ (\epsilon_{e^{-1}(\xi)}, e^{-1}(\xi)) & \text{if } \xi \in F_2 \setminus F_1 \text{ and } t|n = x_\xi|n & \text{(Case 4.)} \end{cases}$$

f_ξ^q is well-defined since $e(\xi) = \xi$ whenever $\xi \in F_1 \cap F_2$, so that $f_\xi^1(s) = f_{e(\xi)}^2(s) = f_\xi^2(s)$ for $s \in 2^n \setminus \{x_\xi|n\}$.

Let us now prove that $q \in \mathbb{P}$. Conditions 1 and 2 follow directly from the definition of F_q and n_q .

To prove that q satisfies condition 3, fix $\xi \in F_q$ and $t \in 2^{n_q} \setminus \{x_\xi|n_q\}$. In Case 1 (resp. Case 2), both conditions 3.a) and 3.b) follow from the fact that p_1 (resp. p_2) is in \mathbb{P} .

In Case 3, we only have to check condition 3.a), which is guaranteed by the fact that $(\delta_\xi : \xi \in F_1 \setminus F_2) \subseteq [2n]^{[2n]}$ are assumed to be constant.

Similarly, in Case 4, we only have to check condition 3.b), which is guaranteed by the fact that $(\epsilon_\xi : \xi \in F_1 \setminus F_2) \subseteq [2n]^{[2n]}$ are assumed to be as needed.

Let us now prove that $q \leq p_1, p_2$. Trivially, $F_1, F_2 \subseteq F_q$ and $n_1, n_2 \leq n_q$.

Given $\xi \in F_q$, $s \in 2^n \setminus \{x_\xi|n\}$ and $t \in 2^{n_q} \setminus \{x_\xi|n_q\}$ such that $s \subseteq t$, let $k \in \{1, 2\}$ be such that $\xi \in F_k$ and notice that we are in cases 1 or 2, since $t|n = s$. Therefore, $f_\xi^q(t) = f_\xi^k(t|n) = f_\xi^k(s)$, which concludes that $q \leq p_1, p_2$.

Finally, notice that the definition of $f_\xi^q(t)$ in cases 1 or 2 imply (3.1). \square

Theorem 3.7. \mathbb{P} is c.c.c.

Proof. For each $\alpha < \omega_1$, let $p_\alpha = (F_\alpha, n_\alpha, (f_\eta^\alpha)_{\eta \in F_\alpha}) \in \mathbb{P}$.

By the Δ -system Lemma, we can assume that $(F_\alpha)_{\alpha < \omega_1}$ forms a Δ -system with root Δ such that for every $\alpha < \beta < \omega_1$,

- $\Delta < F_\alpha \setminus \Delta < F_\beta \setminus \Delta$ and $|F_\alpha| = |F_\beta|$.

Since each $n_\alpha \in \mathbb{N}$, we can suppose that for every $\alpha < \beta < \omega_1$,

- $n_\alpha = n_\beta = n$.

Also, we may assume that if $e_{\alpha\beta} : F_\alpha \rightarrow F_\beta$ is the order-preserving bijective function, then

- for all $\xi \in F_\alpha$, $x_\xi | n = x_{e_{\alpha\beta}(\xi)} | n$ (since both belong to 2^n);
- for all $\xi \in F_\alpha$ and all $s \in 2^n \setminus \{x_\xi | n\}$,

$$f_{e_{\alpha\beta}(\xi)}^\beta(s) = (\varphi, e_{\alpha\beta}(\eta)), \text{ where } f_\xi^\alpha(s) = (\varphi, \eta).$$

Now, fix $\alpha < \beta < \omega_1$. Note that p_α and p_β satisfy the hypothesis of Proposition 3.6. Let, for $\xi \in F_\beta \setminus \Delta$, ϵ_ξ be any function satisfying the condition 3 of the definition of the forcing (for example, ϵ_ξ constant equal to 1); and for $\xi \in F_\alpha \setminus \Delta$, $\delta_\xi \in [2n]^{[2n]}$ be any constant function. Then, by Proposition 3.6, there is $q \leq p_\alpha, p_\beta$ in \mathbb{P} , which concludes the proof. \square

Theorem 3.8. *Let $n \geq 1$ be a natural number. It is consistent that there is a compact Hausdorff totally disconnected space K which is an unordered $2n$ -split Cantor set corresponding to a balanced $2n$ -splitting family $(A_{\xi,i} : \xi < \omega_1, i \in [2n])$ such that given any collection of pairwise disjoint sets $E_\alpha = \{\xi_\alpha^1, \dots, \xi_\alpha^k\} \subseteq \omega_1$ for $\alpha < \omega_1$, given $\epsilon : [k] \times [2n] \rightarrow [2n]$ such that*

$$|\{l \in \{1, 3, 5, \dots, 2n-1\} : \epsilon(i, l) = j\}| = |\{l \in \{2, 4, 6, \dots, 2n\} : \epsilon(i, l) = j\}|$$

and given $\delta : [k] \rightarrow [n]$, there are $\alpha < \beta$ such that for all $1 \leq i \leq k$,

$$R_{\xi_\beta^i} \subseteq A_{\xi_\alpha^i, \delta(i)}$$

and

$$(x_{\xi_\alpha^i}, l) \in A_{\xi_\beta^i, \epsilon(i, l)}.$$

Proof. By Theorem 3.5, \mathbb{P} forces that $(A_{\xi,i} : \xi < \omega_1, i \in [2n])$ as in Definition 3.3 is a balanced $2n$ -splitting family. By Proposition 2.3, we get that the corresponding unordered $2n$ -split Cantor set is a compact, Hausdorff, totally disconnected space. Let us now prove the remaining desired property.

In V , suppose $(\dot{E}_\alpha)_{\alpha < \omega_1}$ and $(\dot{\xi}_\alpha^i)_{\alpha < \omega_1, 1 \leq i \leq k}$ are sequences of names such that \mathbb{P} forces that $\dot{E}_\alpha = \{\dot{\xi}_\alpha^1 < \dots < \dot{\xi}_\alpha^k\}$ and $(\dot{E}_\alpha)_{\alpha < \omega_1}$ is pairwise disjoint.

For each $\alpha < \omega_1$, let $p_\alpha = (F_\alpha, n_\alpha, (f_\eta^\alpha)_{\eta \in F_\alpha}) \in \mathbb{P}$, $\xi_\alpha^1, \dots, \xi_\alpha^k \in \omega_1$ and $E_\alpha, \dots, E_\alpha \subseteq \omega_1$ be finite such that

$$p_\alpha \Vdash \forall 1 \leq i \leq k \quad \dot{\xi}_\alpha^i = \check{\xi}_\alpha^i \text{ and } \dot{E}_\alpha = \check{E}_\alpha.$$

By Lemma 3.2, we can assume without loss of generality that for all $\alpha < \omega_1$, $E_i^\alpha \subseteq F_\alpha$.

By the Δ -system Lemma, we can assume as well that $(F_\alpha)_{\alpha < \omega_1}$ forms a Δ -system with root Δ such that for every $\alpha < \beta < \omega_1$,

- $\Delta < F_\alpha \setminus \Delta < F_\beta \setminus \Delta$ and $|F_\alpha| = |F_\beta|$.

Since each $n_\alpha \in \mathbb{N}$, we can suppose that for every $\alpha < \beta < \omega_1$,

- $n_\alpha = n_\beta = n$.

Also, we may assume that if $e_{\alpha\beta} : F_\alpha \rightarrow F_\beta$ is the order-preserving bijective function, then

- for all $\xi \in F_\alpha$, $x_\xi|n = x_{e_{\alpha\beta}(\xi)}|n$ (since both belong to 2^n);
- for all $\xi \in F_\alpha$ and all $s \in 2^n \setminus \{x_\xi|n\}$,

$$f_{e_{\alpha\beta}(\xi)}^\beta(s) = (\varphi, e_{\alpha\beta}(\eta)), \text{ where } f_\xi^\alpha(s) = (\varphi, \eta).$$

- for all $1 \leq i \leq k$, $e_{\alpha\beta}(\xi_\alpha^i) = \xi_\beta^i$.

Finally, we may assume that for all $1 \leq i \leq k$ we have: either $\xi_\alpha^i = \xi_\beta^i$ for all $\alpha < \beta < \omega_1$; or $\xi_\alpha^i \notin \Delta$ for all $\alpha < \omega_1$ and actually the second case holds by the assumption that E_α 's are pairwise disjoint.

Now, fix $\alpha < \beta < \omega_1$. Note that p_α and p_β satisfy the hypothesis of Proposition 3.6. Taking $\epsilon_{\xi_\beta^i} = \epsilon(i, \cdot)$ and $\delta_{\xi_\alpha^i} = \delta(i)$ (and for $\xi \in F_\beta \setminus (\Delta \cup E_\beta)$, any function ϵ_ξ satisfying the condition 3 of the definition of the forcing; and for $\xi \in F_\alpha \setminus (\Delta \cup E_\alpha)$, any constant function $\delta_\xi \in [2n]^{[2n]}$), by the Proposition 3.6, there is $q \leq p_\alpha, p_\beta$ in \mathbb{P} such that

$$\forall \xi \in F_\alpha \setminus \Delta \quad f_\xi^q(x_{e_{\alpha\beta}(\xi)}|n_q) = (\delta_\xi, \xi) \quad \text{and} \quad f_{e_{\alpha\beta}(\xi)}^q(x_\xi|n_q) = (\epsilon_{e_{\alpha\beta}(\xi)}, \xi).$$

In particular, for all $1 \leq i \leq k$,

$$f_{\xi_\alpha^i}^q(x_{\xi_\beta^i}|n_q) = (\delta(i), \xi_\alpha^i) \quad \text{and} \quad f_{\xi_\beta^i}^q(x_{\xi_\alpha^i}|n_q) = (\epsilon(i, \cdot), \xi_\alpha^i).$$

By the definition of $A_{\xi, j}$, we get that for all $1 \leq i \leq k$,

$$R_{\xi_\beta^i} \subseteq A_{\xi_\alpha^i, \delta(i)} \quad \text{and} \quad (x_{\xi_\alpha^i}, l) \in A_{\xi_\beta^i, \epsilon(i, l)},$$

which concludes the proof. \square

The fact that $2n$ is even is exploited in the above proof. It turns out that there cannot be an analogue of an unordered N -split Cantor set for $N = 3$ which behaves as in Theorem 3.8, as we have the following:

Lemma 3.9. *Let $N \geq 3$ be a natural number. Suppose that K is an unordered N -split Cantor set corresponding to an N -splitting family $(A_{\xi, i} : \xi < \omega_1, i \in [N])$ such that given any sequence of distinct ordinals $(\xi_\alpha : \alpha < \omega_1)$ and $j \in [N]$, there are $\alpha < \beta$ such that*

$$R_{\xi_\beta} \subseteq A_{\xi_\alpha, j}.$$

Suppose that $(f_\alpha, \mu_\alpha)_{\alpha < \omega_1}$ is a biorthogonal system such that $f_\alpha = \chi_{A_\alpha}$ for some clopen subset $A_\alpha \subseteq K$ and $\mu_\alpha = r_\alpha \delta_{(x_{\eta_\alpha}, 1)} + s_\alpha \delta_{(x_{\eta_\alpha}, 2)} + t_\alpha \delta_{(x_{\eta_\alpha}, 3)}$ for all $\alpha < \omega_1$, for some reals $r_\alpha, s_\alpha, t_\alpha$ and some sequence $(\eta_\alpha : \alpha < \omega_1)$. Then there is an uncountable nice biorthogonal system in $C(K)$.

Proof. If there is a biorthogonal system of the form $(\chi_{A_\alpha}, r_\alpha \delta_{y_\alpha})$ for $\alpha < \omega_1$ and $y_\alpha \in K$, we have that $r_\alpha = 1$ for all $\alpha < \omega_1$ and that $y_\alpha \notin A_\beta$ for any $\beta \neq \alpha$ and $y_\alpha \in A_\alpha$. So $(\chi_{A_{\alpha+1}}, \delta_{y_{\alpha+1}} - \delta_{y_\alpha})$, say, for all limit ordinals α is a nice biorthogonal system.

If there is a biorthogonal system of the form $(\chi_{A_\alpha}, r_\alpha \delta_{y_\alpha} + s_\alpha \delta_{z_\alpha})$ for $\alpha < \omega_1$ and $y_\alpha, z_\alpha \in K$, and $r_\alpha, s_\alpha, r_\alpha + s_\alpha \neq 0$, then $r_\alpha, s_\alpha \notin A_\beta$ for any $\alpha \neq \beta$ and a similar argument as above gives a nice biorthogonal system. If $r_\alpha + s_\alpha = 0$ and $r_\alpha, s_\alpha \neq 0$, we may assume that $r_\alpha > 0$ and so $s_\alpha = -r_\alpha$. It follows from the fact that $(r_\alpha \delta_{y_\alpha} + s_\alpha \delta_{z_\alpha})(\chi_{A_\alpha}) = 1$ that $r_\alpha = 1$ and $s_\alpha = -1$, and so we have a nice biorthogonal system.

Hence, without loss of generality, we may assume that $r_\alpha, s_\alpha, t_\alpha \neq 0$ for all $\alpha < \omega_1$. First let us see that there is an uncountable $X \subseteq \omega_1$ such that $r_\alpha + s_\alpha + t_\alpha = 0$ for all $\alpha \in X$. If not, then there is an uncountable $X \subseteq \omega_1$ and an $\varepsilon > 0$ such that $|r_\alpha + s_\alpha + t_\alpha| > \varepsilon$ for each $\alpha \in X$.

Now note that as $\mu_\alpha(\chi_{A_\alpha}) = 1 \neq 0$, we have $j \in \{1, 2, 3\}$ such that $(x_{\eta_\alpha}, j) \in A_\alpha$. We may assume that it is the same j for all $\alpha \in X$. By the form of the basic neighbourhoods of points (x_{η_α}, j) we have $s \in 2^m$ for some $m \in \mathbb{N}$ such that $(x_{\eta_\alpha}, j) \in V_s \cap A_{\eta_\alpha, j} \subseteq A_\alpha$. We may assume that it is the same s for all $\alpha \in X$. It follows that for some $n \in \mathbb{N}$ we have $s = x_{\eta_\alpha} |n$ for all $\alpha \in X$ and so that $R_{\eta_\alpha} \subseteq V_s$ for all $\alpha \in X$. Apply the hypothesis of the lemma and obtain $\alpha < \beta$ both in X such that $R_{\eta_\beta} \subseteq A_{\eta_\alpha, j}$ and we get that $R_{\eta_\beta} \subseteq V_s \cap A_{\eta_\alpha, j} \subseteq A_\alpha$. This means that $0 = \mu_\beta(\chi_{A_\alpha}) = r_\beta + s_\beta + t_\beta$ contradicting the choice of $\beta \in X$. So we may assume that $r_\alpha + s_\alpha + t_\alpha = 0$ for all $\alpha < \omega_1$.

For three non-zero numbers whose sum is zero, there cannot be any subsum which is zero, this means, that for $\alpha \neq \beta$, as $\mu_\alpha(A_\beta) = 0$, we have that either $\{x_\alpha, y_\alpha, z_\alpha\} \cap A_\beta = \emptyset$ or $\{x_\alpha, y_\alpha, z_\alpha\} \subseteq A_\beta$. So, to make an uncountable nice biorthogonal system out of points $\{x_\alpha, y_\alpha, z_\alpha\}$ and functions χ_{A_α} , we need to find any fixed pair of them which is separated by A_α for uncountably many α 's.

But A_α must separate some pair as $\mu_\alpha(A_\alpha) = 1$, so choose an uncountable subset Y of ω_1 on which the same pair is separated, say $x_\alpha \in A_\alpha$ and $z_\alpha \notin A_\alpha$.

Define $\nu_\alpha = \delta_{x_\alpha} - \delta_{z_\alpha}$ and note that $(\chi_{A_\alpha}, \nu_\alpha)_{\alpha \in Y}$ is an uncountable nice biorthogonal system. \square

4. BIORTHOGONAL AND SEMIBIORTHOGONAL SYSTEMS IN $C(K_{2n})$ 'S

Lemma 4.1. *Suppose that $\theta > \rho > 0$, $n \in \mathbb{N}$, $n \geq 2$, r_1, \dots, r_{2n} are reals such that*

- (1) $|\sum_{1 \leq i \leq 2n} r_i| < \rho$,
- (2) *there is $1 \leq i_0 \leq 2n$ such that $r_{i_0} > \theta$,*
- (3) *there is $1 \leq i_1 \leq 2n$ such that $r_{i_1} = 0$.*

Then there are $1 \leq i, j \leq 2n$ such that $(-1)^{i+j} = -1$ and

$$r_i + r_j < \frac{2n\rho - \theta}{n(2n - 2)}.$$

Proof. By (1) and (2), since $\theta > \rho$, there must be an $i_2 \in \{1, \dots, 2n\} \setminus \{i_0, i_1\}$ such that

$$r_{i_2} < -\frac{\theta - \rho}{2n - 2} = \frac{\rho - \theta}{2n - 2} < \frac{2n\rho - \theta}{n(2n - 2)}.$$

So, if there is i_3 such that $(-1)^{i_2+i_3} = -1$, and $r_{i_3} \leq 0$, then we are done. Otherwise, there are at least n positive numbers r_i (at least all r_i 's for i of the other parity than i_2) and so, by (3), at most $n - 1$ negative numbers r_i . Let $k \geq 0$ be such that $n + k$ is the number of positive r_i 's, let r_{i_4} be the smallest positive number among r_i 's and let $r_{i_5} \leq -\frac{\theta - \rho}{2n - 2}$ be the smallest negative number (i.e., of the biggest absolute value) among r_i 's.

So we have

$$(n + k)r_{i_4} + (n - k - 1)r_{i_5} \leq \sum \{r_i : r_i > 0\} + \sum \{r_i : r_i < 0\} < \rho$$

So,

$$n(r_{i_4} + r_{i_5}) + k(r_{i_4} - r_{i_5}) < \rho + r_{i_5} \leq \rho - \frac{\theta - \rho}{2n - 2}$$

But $r_{i_4} - r_{i_5}$ is non-negative, so

$$r_{i_4} + r_{i_5} < (1/n)(\rho - \frac{\theta - \rho}{2n - 2}) = \frac{(2n - 1)\rho - \theta}{n(2n - 2)},$$

as required. \square

Lemma 4.2. *Let $n \geq 2$. Suppose that $(f_\alpha)_{\alpha < \omega_1}$ is a sequence of continuous rational simple functions on K_{2n} as in Theorem 3.8 and $(\mu_\alpha)_{\alpha < \omega_1}$ is a sequence of $(2n - 1)$ -supported atomic Radon measures on K_{2n} . Then either there are $\alpha < \beta < \omega_1$ such that*

$$a) \quad \left| \int f_\alpha d\mu_\beta \right| > 0.01/2n^2(2n - 2)$$

or there is $\alpha \in \omega_1$ such that

$$b) \quad \int f_\alpha d\mu_\alpha < 0.99$$

or there are $\alpha < \beta < \omega_1$ such that

$$c) \quad \int f_\beta d\mu_\alpha < -0.89/2n^2(2n - 2).$$

Proof. By the separability of $C_0 \equiv C(2^\omega)$ (see Definition 2.4), Lemma 2.6 and thinning out the sequence, we may assume that for all $\alpha < \omega_1$ we have

$$f_\alpha = g + \sum_{1 \leq i \leq k} \sum_{1 \leq l \leq 2n-1} q_{i,l} \chi_{A_{\xi_\alpha^i, l} \cap V_{s_i}}$$

for some simple rational function $g \in C_0$, $F_\alpha = \{\xi_\alpha^1, \dots, \xi_\alpha^k\} \subseteq \omega_1$, some $s_i \in 2^{m_i}$, $m_i \in N$ and some rationals $q_{i,l}$, $1 \leq i \leq k$ and $1 \leq l \leq 2n$ such that $s_i = r_{\xi_\alpha^i} | m_i$ and such that

$$\sum_{1 \leq i \leq k} \left(\max_{1 \leq l \leq 2n} |q_{i,l}| \right) |\mu_\alpha|(V_{s_i} \setminus R_{\xi_\alpha^i}) \leq 0.01/2n^2(2n - 2).$$

By thinning out the sequence (applying the Δ -system lemma, see [Ku]) and moving some identical parts to g we may assume that F_α 's are pairwise disjoint and g (no longer in C_0) is fixed. So, we will be allowed to use the following decompositions:

Claim 0. *For each $\alpha, \beta < \omega_1$ we have*

$$\begin{aligned} \int f_\alpha d\mu_\beta &= \int g d\mu_\beta + \sum_{1 \leq i \leq k} \sum_{1 \leq l \leq 2n-1} q_{i,l} \mu_\beta(A_{\xi_\alpha^i, l} \cap R_{\xi_\beta^i} \cap V_{s_i}) + \\ &+ \sum_{1 \leq i \leq k} \sum_{1 \leq l \leq 2n-1} q_{i,l} \mu_\beta(A_{\xi_\alpha^i, l} \cap V_{s_i} \setminus R_{\xi_\beta^i}) \end{aligned}$$

Here, the last term is small by the above application of Lemma 2.6, the first term will be shown small by the claim below and so the value of the integral will depend on the relation of the points from $R_{\xi_\beta^i, l}$ with the sets $A_{\xi_\alpha^i, l}$ which is "as we wish" on any uncountable set by Theorem 3.8.

Claim 1. *Either a) holds or for all but countably many α 's in ω_1 we have*

$$\left| \int g d\mu_\alpha \right| \leq 0.02/2n^2(2n - 2).$$

Proof of the claim. If the inequality does not hold for uncountably many α s, then by Theorem 3.8 we can find among them $\alpha < \beta < \omega_1$ such that $R_{\xi_\beta^i} \subseteq A_{\xi_\alpha^i, 2n}$ for all $1 \leq i \leq k$. By Claim 0 we get that

$$\begin{aligned} & \left| \int f_\alpha d\mu_\beta \right| \geq \left| \int g d\mu_\beta \right| - \sum_{1 \leq i \leq k} \sum_{1 \leq l \leq 2n-1} |q_{i,l}| |\mu_\beta(A_{\xi_\alpha^i, l} \cap V_{s_i})| \geq \\ & \geq \left| \int g d\mu_\beta \right| - \sum_{1 \leq i \leq k} \left(\max_{1 \leq l \leq m} |q_{i,l}| \right) |\mu_\beta|(V_{s_i} \setminus R_{\xi_\beta^i}) > \frac{0.02 - 0.01}{2n^2(2n-2)} = \frac{0.01}{2n^2(2n-2)} \end{aligned}$$

obtaining a) of the lemma and concluding Claim 1.

Claim 2. *Either a) holds or for all but countably many α 's in ω_1 we have for each $1 \leq l_0 \leq 2n-1$*

$$\left| \sum_{1 \leq i \leq k} q_{i, l_0} \mu_\alpha(R_{\xi_\alpha^i}) \right| \leq 0.04/2n^2(2n-2)$$

Proof of the claim. Without loss of generality we may assume that the condition from Claim 1 holds for all $\alpha < \beta < \omega_1$. Fix l_0 as above. Suppose that the condition above does not hold for uncountably many α s, then by Theorem 3.8 we obtain among them $\alpha < \beta$ such that for all $1 \leq i \leq k$

$$R_{\xi_\beta^i} \subseteq A_{\xi_\alpha^i, l_0}.$$

So by Claim 0 we have

$$\begin{aligned} & \left| \int f_\alpha d\mu_\beta \right| \geq \left| \sum_{1 \leq i \leq k} \sum_{1 \leq l \leq 2n-1} q_{i,l} \mu_\beta(R_{\xi_\beta^i} \cap A_{\xi_\alpha^i, l} \cap V_{s_i}) \right| - \left| \int g d\mu_\beta \right| - \\ & \quad - \sum_{1 \leq i \leq k} \left(\max_{1 \leq l \leq 2n-1} |q_{i,l}| \right) |\mu_\beta|(V_{s_i} \setminus R_{\xi_\beta^i}) \geq \\ & \geq \left| \sum_{1 \leq i \leq k} q_{i, l_0} \mu_\beta(R_{\xi_\beta^i}) \right| - \left| \int g d\mu_\beta \right| - \sum_{1 \leq i \leq k} \left(\max_{1 \leq l \leq 2n-1} |q_{i,l}| \right) |\mu_\beta|(V_{s_i} \setminus R_{\xi_\beta^i}) > \\ & > \frac{0.04 - 0.02 - 0.01}{2n^2(2n-2)} = 0.01/2n^2(2n-2), \end{aligned}$$

obtaining a) and concluding Claim 2.

Claim 3. *Either a) or b) holds or there is $l_0 \in \{1, \dots, 2n\}$ such that for uncountably many α 's in ω_1 we have*

$$\sum_{1 \leq i \leq k} q_{i, l_0} \mu_\alpha(\{(x_{\xi_\alpha^i}, l_0)\}) > 0.96/(2n)$$

Proof of the claim. Assume that a) does not hold, i.e., the conditions from Claim 1 and Claim 2 hold for all $\alpha < \omega_1$. Now, suppose also that the condition from Claim 3 does not hold for any $l_0 \in \{1, \dots, 2n\}$. By Claim 0 for $\alpha = \beta$ we have

$$\int f_\alpha d\mu_\alpha \leq \sum_{1 \leq i \leq k} \sum_{1 \leq l \leq 2n-1} q_{i,l} \mu_\alpha(\{(x_{\xi_\alpha^i}, l)\}) +$$

$$-0.02/2n^2(2n-2) - 0.01/2n^2(2n-2) \leq (2n-1)0.96/(2n) - 0.03 < 0.99$$

that is we obtain b), which concludes the proof of Claim 3.

To finish the proof of the lemma, we need to assume that a) and b) fail, i.e., the conditions of all the above claims hold, and we need to get c). Fix $\alpha < \omega_1$, we will apply Lemma 4.1 for

$$r_{l,\alpha} = \sum_{1 \leq i \leq k} q_{i,l_0} \mu_\alpha(\{(x_{\xi_\alpha^i}, l)\})$$

and $l \in \{1, \dots, 2n\}$. By the fact that the supports of the measures μ_α have at most $2n - 1$ elements, one of $r_{l,\alpha}$'s must be zero. By Claim 3 we have that $r_{l_0,\alpha} > \theta = 0.96/(2n)$ and by Claim 2 that $\sum_{1 \leq l \leq 2n} r_{l,\alpha} < \rho = 0.04/(2n)^2$. So by Lemma 4.1 we find $1 \leq l_{1,\alpha}, l_{2,\alpha} \leq 2n$ of different parities such that

$$\begin{aligned} \sum_{1 \leq i \leq k} q_{i,l_0} \mu_\alpha(\{(x_{\xi_\alpha^i}, l_{1,\alpha}), (x_{\xi_\alpha^i}, l_{2,\alpha})\}) &< \frac{2n\rho - \theta}{n(2n-2)} = \\ &= \frac{2n(0.04/(2n)^2) - 0.96/(2n)}{n(2n-2)} = -\frac{0.92}{2n^2(2n-2)}. \end{aligned}$$

We may assume that $l_{1,\alpha} = l_1$ and $l_{2,\alpha} = l_2$ for all $\alpha < \omega_1$. Note that by Theorem 3.8 we can find $\alpha < \beta < \omega_1$ such that

$$\{(x_{\xi_\alpha^i}, l_1), (x_{\xi_\alpha^i}, l_2)\} \subseteq A_{\xi_\beta^i, l_0}$$

and

$$R_{\xi_\alpha^i} \setminus \{(x_{\xi_\alpha^i}, l_1), (x_{\xi_\alpha^i}, l_2)\} \subseteq A_{\xi_\beta^i, 2n}$$

for all $1 \leq i \leq k$. This and Claim 0 with α and β switched implies that

$$\begin{aligned} \int f_\beta d\mu_\alpha &\leq \sum_{1 \leq i \leq k} q_{i,l_0} \mu_\alpha(\{(x_{\xi_\alpha^i}, l_1), (x_{\xi_\alpha^i}, l_2)\}) + \\ &+ \sum_{1 \leq i \leq k} \max_{1 \leq l \leq 2n-1} (|q_{i,l}|) |\mu_\alpha|(V_{s_i} \setminus R_{\xi_\alpha^i}) + \left| \int g d\mu_\alpha \right| \leq \\ &\leq \frac{-0.92 + 0 + 0.01 + 0.02}{(2n)^2(2n-2)} = -0.89/(2n)^2(2n-2) \end{aligned}$$

which completes the proof of the lemma. \square

Theorem 4.3. *Let $n \geq 2$. If K_{2n} is an unordered $2n$ -split Cantor set as in Theorem 3.8, then there are no uncountable semibiorthogonal sequences in $C(K_{2n})$ whose functionals are $(2n-1)$ -supported but there is an uncountable biorthogonal system whose functionals are $2n$ -supported.*

Proof. Suppose $(f_\alpha, \mu_\alpha)_{\alpha < \omega_1} \subseteq C(K_{2n}) \times M(K_{2n})$ is a sequence whose functionals are $2n-1$ -supported and that $\int f_\alpha d\mu_\beta = 0$ for all $\alpha < \beta < \omega_1$ as well as $\int f_\alpha d\mu_\alpha = 1$ for all $\alpha < \omega_1$.

We may assume without loss of generality that $\|\mu_\alpha\| \leq M$ for some positive M . By the Stone Weierstrass theorem we can choose $f'_\alpha \in C(K)$ which is a rational simple function and

$$\|f'_\alpha - f_\alpha\| < 0.01/2Mn^2(2n-2).$$

This means that a) and b) of Lemma 14 do not hold, for f'_α 's instead of f_α 's i.e., c) holds, which implies that $(f_\alpha, \mu_\alpha)_{\alpha < \omega_1}$ is not semibiorthogonal. \square

Theorem 4.4. *If K_{2n} is an unordered $2n$ -split Cantor set as in Theorem 3.8, then $hd(K_{2n}^n) = \omega$.*

Proof. We will be using the well-known fact that a regular space is hereditarily separable if and only if it has no uncountable left-separated sequence (see Theorem 3.1 of [16]).

Suppose $(y_\alpha)_{\alpha < \omega_1}$ is a left-separated sequence in K_{2n}^n of cardinality \aleph_1 . Hence, for each $\alpha < \omega_1$, $y_\alpha = (y_\alpha^1, \dots, y_\alpha^n)$, where each $y_\alpha^m \in K_{2n}$ and, by the definition of a left-separated sequence, for each $\alpha < \omega_1$ and each $m \in [n]$, there is an open basic neighbourhood U_α^m of y_α^m such that

$$\forall \alpha < \omega_1 \quad \forall m \in [n] \quad y_\alpha^m \in U_\alpha^m$$

and that

$$\forall \alpha < \beta < \omega_1 \quad \exists m \in [n] \quad y_\alpha^m \notin U_\beta^m.$$

We may assume without loss of generality that

$$\{m \in [n] : y_\alpha^m \in 2^\omega \setminus \{x_\xi : \xi < \omega_1\}\} = \{m \in [n] : y_\beta^m \in 2^\omega \setminus \{x_\xi : \xi < \omega_1\}\}$$

for every $\alpha < \beta < \omega_1$ and let us call this set I .

For each $m \in [n] \setminus I$, let ξ_α^m be a countable ordinal and j_α^m be an element of $[n]$ such that $y_\alpha^m = (x_{\xi_\alpha^m}, j_\alpha^m)$.

Now, for each $m \in [n]$, let $s_\alpha^m \in 2^{<\omega}$ such that

$$U_\alpha^m = \begin{cases} V_{s_\alpha^m} & \text{if } m \in I \\ V_{s_\alpha^m} \cap A_{\xi_\alpha^m, j_\alpha^m} & \text{if } m \notin I \end{cases}$$

Put $E_\alpha = \{\xi_\alpha^m : m \in [n] \setminus I\}$.

Without loss of generality, we may assume that:

- there is $j_m \in [n]$ such that $j_\alpha^m = j_m$ for all $\alpha < \omega_1$;
- there is $s_m \in 2^{<\omega}$ such that $s_\alpha^m = s_m$ for all $\alpha < \omega_1$ (this already guarantees that each $y_\alpha^m \in V_{s_m}$);
- for all $m \in [n] \setminus I$, either

$$\forall \alpha < \beta < \omega_1 \quad \xi_\alpha^m = \xi_\beta^m,$$

or

$$\forall \alpha < \beta < \omega_1 \quad \xi_\alpha^m < \xi_\beta^m.$$

- $(E_\alpha)_{\alpha < \omega_1}$ is a Δ -system with root Δ such that for every $\alpha < \beta < \omega_1$, $\Delta < E_\alpha \setminus \Delta < E_\beta \setminus \Delta$ and $|E_\alpha| = |E_\beta|$.

If $E_\alpha \setminus \Delta = \emptyset$, the left-separated sequence in K_{2n}^n would lead to a left-separated sequence in a finite power of 2^ω , which is not possible since 2^ω is hereditarily separable in all finite powers. Therefore, each $E_\alpha \setminus \Delta \neq \emptyset$ and they are pairwise disjoint.

For each $\alpha < \omega_1$, enumerate $E_\alpha \setminus \Delta = \{\eta_\alpha^1 < \dots < \eta_\alpha^k\}$. We may assume that $\xi_\alpha^m = \eta_\alpha^i$ if and only if $\xi_\beta^m = \eta_\beta^i$.

Claim. For each $1 \leq i \leq k$, it is possible to find $I_i \subseteq [2n]$ of cardinality N and a bijection $\sigma_i : I_i \rightarrow [2n] \setminus I_i$ such that $\sigma_i(l)$ and l have opposite parity and

$$\{j \in [2n] : \exists m \in [n] \text{ such that } j = j_m \text{ and } \xi_\alpha^m = \eta_\alpha^i\} \subseteq I_i.$$

Proof of the claim. It follows simply from the fact that the set

$$\{j \in [2n] : \exists m \in [n] \text{ such that } j = j_m \text{ and } \xi_\alpha^m = \eta_\alpha^i\}$$

has cardinality at most n so that we can find I_i containing it and that whenever we have a partition of $[2n]$ into two sets A and B , both of size n , A has as many odds as B has evens, and vice-versa. This concludes the proof of the claim.

Now, let $\epsilon : [k] \times [2n] \rightarrow [2n]$ be defined by

$$\epsilon(i, l) = \begin{cases} l & \text{if } l \in I_i \\ \sigma_i^{-1}(l) & \text{if } l \in [2n] \setminus I_i \end{cases}$$

Notice that for each $i \in [k]$, $l \in I_i$ and $j \in [2n]$, $\epsilon(i, l) = j$ if and only if $\epsilon(i, \sigma(l)) = j$. Since $\sigma(l)$ and l have opposite parities, we get that ϵ has the desired property, that is,

$$|\{l \in \{1, 3, 5, \dots, 2n-1\} : \epsilon(i, l) = j\}| = |\{l \in \{2, 4, 6, \dots, 2n\} : \epsilon(i, l) = j\}|.$$

By Theorem 3.8, there are $\alpha < \beta$ such that for all $i \in [k]$,

$$(x_{\eta_\alpha^i}, l) \in A_{\eta_\beta^i, \epsilon(i, l)}.$$

Fix $m \in [n]$ and let us prove that $y_\alpha^m \in U_\beta^m$, contradicting the assumption. If $m \notin I$, then $y_\alpha^m \in V_{s_m} = U_\beta^m$. If $m \in I$ and $\xi_\alpha^m \in \Delta$, then $\xi_\alpha^m = \xi_\beta^m \in U_\beta^m$. Finally, if $m \in I$ and $\xi_\alpha^m \notin \Delta$, then there is $i \in [k]$ such that $\xi_\alpha^m = \eta_\alpha^i$ and $\xi_\beta^m = \eta_\beta^i$. In this case we have that $j_m \in I_i$ and so, $\epsilon(i, j_m) = j_m$, which guarantees that

$$y_\alpha^m = (x_{\xi_\alpha^m}, j_m) = (x_{\eta_\alpha^i}, j_m) \in A_{\eta_\beta^i, j_m} = A_{\xi_\beta^m, j_m}.$$

Since also $y_\alpha^m \in V_{s_m}$, we get that $y_\alpha^m \in U_\beta^m$, which concludes the proof. \square

REFERENCES

1. M. Bell, J. Ginsburg, and S. Todorćević, *Countable spread of $\exp Y$ and λY* , *Topology Appl.* **14** (1982), no. 1, 1–12.
2. J. M. Borwein and J. D. Vanderwerff, *Banach spaces that admit support sets*, *Proc. Amer. Math. Soc.* **124** (1996), no. 3, 751–755.
3. C. Brech and P. Koszmider, *Thin-very tall compact scattered spaces which are hereditarily separable*, accepted to *Trans. Amer. Math. Soc.* (2009).
4. M. Džamonja, I. Juhász; *CH, a problem of Rolewicz and bidiscrete systems*, to appear in *Top. App.*
5. R. Engelking, *General topology*, second ed., Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989, Translated from the Polish by the author.
6. M. Fabian, P. Habala, P. Hájek, V. Montesinos Santalucía, J. Pelant, and V. Zizler, *Functional analysis and infinite-dimensional geometry*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 8, Springer-Verlag, New York, 2001.
7. C. Finet and G. Godefroy, *Biorthogonal systems and big quotient spaces*, *Banach space theory* (Iowa City, IA, 1987), *Contemp. Math.*, vol. 85, Amer. Math. Soc., Providence, RI, 1989, pp. 87–110.
8. A. S. Granero, M. Jiménez Sevilla, and J. P. Moreno, *Convex sets in Banach spaces and a problem of Rolewicz*, *Studia Math.* **129** (1998), no. 1, 19–29.
9. P. Hájek, V. Montesinos Santalucía, J. Vanderwerff, and V. Zizler, *Biorthogonal systems in Banach spaces*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 26, Springer, New York, 2008.
10. M. Katětov, *Complete normality of Cartesian products*, *Fund. Math.* **35** (1948), 271–274.
11. P. Koszmider, *On a problem of Rolewicz about Banach spaces that admit support sets*, *J. Funct. Anal.* **257** (2009), no. 9, 2723–2741.
12. K. Kunen, *Set theory. an introduction to independence proofs*, *Studies in Logic and the Foundations of Mathematics*, vol. 102, North-Holland Publishing Co., Amsterdam, 1980.
13. A. J. Lazar, *Points of support for closed convex sets*, *Illinois J. Math.* **25** (1981), no. 2, 302–305.
14. J. D. Monk, *Cardinal functions on Boolean algebras*, *Lectures in Mathematics ETH Zürich*, Birkhäuser Verlag, Basel, 1990.
15. S. Negrepontis, *The Stone space of the saturated Boolean algebras*, *Trans. Amer. Math. Soc.* **141** (1969), 515–527.

16. J. Roitman, *Basic S and L* , Handbook of set-theoretic topology, North-Holland, Amsterdam, 1984, pp. 295–326.
17. S. Rolewicz, *On convex sets containing only points of support*, Comment. Math. Special Issue **1** (1978), 279–281, Special issue dedicated to Władysław Orlicz on the occasion of his seventy-fifth birthday.
18. S. Shelah, *On uncountable Boolean algebras with no uncountable pairwise comparable or incomparable sets of elements*, Notre Dame J. Formal Logic **22** (1981), no. 4, 301–308.
19. ———, *Uncountable constructions for $B.A.$, $e.c.$ groups and Banach spaces*, Israel J. Math. **51** (1985), no. 4, 273–297.
20. I. Singer, *Bases in Banach spaces. I*, Springer-Verlag, New York, 1970, Die Grundlehren der mathematischen Wissenschaften, Band 154.
21. ———, *Bases in Banach spaces. II*, Editura Academiei Republicii Socialiste România, Bucharest, 1981.
22. Z. Szentmiklóssy, *S -spaces and L -spaces under Martin's axiom*, Topology, Vol. II (Proc. Fourth Colloq., Budapest, 1978), Colloq. Math. Soc. János Bolyai, vol. 23, North-Holland, Amsterdam, 1980, pp. 1139–1145.
23. S. Todorčević, *Irredundant sets in Boolean algebras*, Trans. Amer. Math. Soc. **339** (1993), no. 1, 35–44.
24. ———, *Biorthogonal systems and quotient spaces via Baire category methods*, Math. Ann. **335** (2006), no. 3, 687–715.

INSTITUTO DE MATEMÁTICA, ESTATÍSTICA E COMPUTAÇÃO CIENTÍFICA, UNIVERSIDADE ESTADUAL DE CAMPINAS, RUA SÉRGIO BUARQUE DE HOLANDA, 651 - 13083-859, CAMPINAS, BRAZIL

Current address: Departamento de Matemática, Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão, 1010 - 05508-090, São Paulo, Brazil

E-mail address: christina.brech@gmail.com

INSTYTUT MATEMATYKI POLITECHNIKI ŁÓDZKIEJ, UL. WÓLCZAŃSKA 215, 90-924 ŁÓDŹ, POLAND

E-mail address: pkoszmider.politechnika@gmail.com