

# On constrained optimization with nonconvex regularization\*

E. G. Birgin<sup>†</sup>      J. M. Martínez<sup>‡</sup>      A. Ramos<sup>§</sup>

September 13, 2019

## Abstract

In many engineering applications it is necessary to minimize smooth functions plus penalty (or regularization) terms that violate smoothness and convexity. Specific algorithms for this type of problems are available in recent literature. Here a smooth reformulation is proposed and equivalence with the original problem is proved both from the point of view of global and local optimization. Moreover, for the cases in which the objective function is much more expensive than the constraints, model-intensive algorithms, accompanied by their convergence and complexity theories, are introduced. Finally, numerical experiments are presented.

**Key words:** Constrained non-Lipschitz nonsmooth optimization, complexity analysis, optimality conditions.

## 1 Introduction

Given  $\mathcal{C} \subset \mathbb{R}^n$ , continuously differentiable functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $g : \mathbb{R}^n \rightarrow \mathbb{R}^d$ , a penalty function  $\pi : \mathbb{R} \rightarrow \mathbb{R}_+$  such that  $\pi(t) = 0$  if  $t \leq 0$ , and a regularization parameter  $\sigma > 0$ , we consider the constrained optimization problem given by

$$\begin{aligned} \text{Minimize} \quad & f(x) + \sigma \sum_{j=1}^d \pi(g_j(x)) \\ \text{subject to} \quad & x \in \mathcal{C}. \end{aligned} \tag{1}$$

Problem (1) may arise in the process of solving problem

$$\text{Minimize } f(x) \text{ subject to } g(x) \leq 0, x \in \mathcal{C} \tag{2}$$

---

\*This work was supported by FAPESP (grants 2013/07375-0, 2016/01860-1, and 2018/24293-0) and CNPq (grants 309517/2014-1, 303750/2014-6, and 438185/2018-8).

<sup>†</sup>Department of Computer Science, Institute of Mathematics and Statistics, University of São Paulo, Rua do Matão, 1010, Cidade Universitária, 05508-090, São Paulo, SP, Brazil. e-mail: egbirgin@ime.usp.br

<sup>‡</sup>Department of Applied Mathematics, Institute of Mathematics, Statistics and Scientific Computing, State University of Campinas, R. Sérgio Buarque de Holanda, 651, Cidade Universitária, 13083-859, Campinas, SP, Brazil. e-mail: martinez@ime.unicamp.br

<sup>§</sup>Department of Mathematics, Federal University of Paraná, Centro Politécnico, Jd. das Américas CP 19081, 81531-980, Curitiba, PR, Brazil. e-mail: albertoramos@ufpr.br

by means of penalization with respect to the constraints  $g(x) \leq 0$ . In this case, we expect that a solution to (2) should emerge when  $\sigma$  is sufficiently large or, at least, when  $\sigma \rightarrow \infty$  (see, for example, [25]). Many times, the objective function of (1) is smooth and  $\pi'(0) = 0$ . However, we are especially interested in the case in which  $\pi'(0) > 0$  and, above all, in the case in which  $\pi'(0)$  does not exist and  $\lim_{t \rightarrow 0^+} \pi'(t) = \infty$ .

For several reasons, solutions to (1) for different values of  $\sigma$  may be meaningful independently of its connection with (2). In fact, many times, the solution to (1) when  $\sigma \rightarrow \infty$  is not relevant and the fulfillment of all the constraints  $g_j(x) \leq 0$  is not desirable or, perhaps, impossible. Instead, the ideal asymptotic problem that one wishes to solve is, roughly speaking,

$$\text{Minimize } f(x) \text{ subject to } x \in \mathcal{C} \quad (3)$$

with the compromise that the number of indices  $j$  such that  $g_j(x) > 0$  should be small. For example, a typical portfolio problem could require the minimization of the expected loss  $f(x)$  subject to standard constraints  $x \in \mathcal{C}$  and the additional requirement that the number of scenarios under which shortfall occurs should be as small as possible [10]. We expect to achieve this objective employing penalty functions that are concave for  $t \geq 0$  and such that  $\lim_{t \rightarrow 0^+} \pi'(t) > 0$ .

The case in which  $\mathcal{C}$  is polyhedral and  $\pi(t) = t^q$  if  $t \geq 0$ , with  $q \in (0, 1)$ , has been considered in [5, 22]. The problem

$$\text{Minimize } f(x) + \sigma \sum_{j=1}^n |x_j|^q \text{ subject to } x \in \mathcal{C}, \quad (4)$$

called  $L_q$ -regularized optimization problem, has been object of many studies starting from Tikhonov [26] and the enormous literature on inverse problems. The case  $q = 2$  corresponds to the most classical regularization, intended to handle problems in which the mere minimization of  $f(x)$  onto  $\mathcal{C}$  is very ill-conditioned and, so, its exact solution is, frequently, meaningless. The case  $q = 1$  preserves the possible convexity of  $f(x)$  and has been shown to induce sparse regularized minimizers of  $f(x)$  (see [11]). With  $q < 1$  possible convexity of the objective function is lost but the tendency to find sparse solutions  $x$  is even more emphatic. Writing  $g_j(x) = x_j$  and  $g_{n+j}(x) = -x_j$  for  $j = 1, \dots, n$  and defining  $\pi(t) = t^q$  for all  $t \geq 0$ , problem (4) takes the form (1); so (4) is a particular case of the problem studied in this paper. Practical applications of problems of the form (1) include machine learning [16], information theory [18], image restoration [5], signal processing [4], variable selection [17] and others. Interesting examples were given in [22, Appx.1].

The rest of this paper is organized as follows. In Section 2, we introduce a smooth reformulation of (1) and we prove its equivalence with the original problem. In Section 3, we derive optimality conditions using the reformulation. In Section 4, we introduce a model-intensive algorithm for solving the problem and we prove its convergence and complexity. In Section 5, we introduce a specific algorithm for the case in which the constraints are linear. In Section 6, we present numerical experiments. Finally, in Section 7, we state conclusions and lines for future research.

**Notation.**  $\|\cdot\|$  denotes an arbitrary norm.  $\mathbb{R}_+$  denotes the set of nonnegative elements of  $\mathbb{R}$ . The  $i$ -th component of a vector  $v$  is denoted  $v_i$  or  $[v]_i$ . If  $v \in \mathbb{R}^n$  is a vector with components  $v_i$ ,  $v_+$  is the vector with components  $\max\{0, v_i\}$ ,  $i = 1, \dots, n$ .

## 2 Reformulation

A smooth reformulation of problem (1) will be introduced in this section. The penalty functions employed in this work will be required to satisfy the assumptions below.

**Assumption A1.**  $\pi$  is continuous, concave, strictly increasing for  $t \geq 0$ . Moreover,  $\pi'(t)$  exists and is continuous for all  $t > 0$ .

**Assumption A2.** The restriction of  $\pi$  to  $\mathbb{R}_+$  admits an inverse  $\pi^{-1} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ , the derivative  $(\pi^{-1})'(t)$  exists for all  $t > 0$ , and the right-derivative  $(\pi^{-1})'(0)_+$  also exists.

Assumptions A1 and A2 are satisfied by penalty functions in practical applications such as ridge regression [19], smoothly clipped absolute deviation penalization (SCAD) [17], minimax concave penalties (MCP) [27], fraction penalty optimization [24], log-penalty minimization [11], and many others. Note that Assumptions A1 and A2 do not require  $\lim_{t \rightarrow 0^+} \pi'(t) = \infty$ , although this property will be interesting for the purpose of minimizing the number of indices  $j$  such that  $g_j(x) > 0$ .

The possible lack of smoothness of (1) is due to the non-smoothness of  $\pi(t)$  when  $t = 0$ . Therefore, standard methods based on gradient information for solving (1) may be inappropriate. In order to avoid this inconvenient, several approaches were proposed. Among them, we can mention smoothing approximation methods [5, 14, 22], iterative reweighted algorithms [23, 21, 15], and interior point methods [6, 20]. Our proposal, based on a smooth reformulation has the advantage that standard well established algorithms for constrained optimization based on first-order or higher-order derivative information on the derivatives may be used. This is possible thanks to the fact that the function  $\pi^{-1} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is continuously differentiable.

Problem (1) is obviously equivalent to

$$\text{Minimize } f(x) + \sigma \sum_{j=1}^d y_j \text{ subject to } y_j = \pi(g_j(x)) \text{ for } j = 1, \dots, d \text{ and } x \in \mathcal{C}. \quad (5)$$

If  $y_j > \pi(g_j(x))$ , taking  $\bar{y}_j = \pi(g_j(x))$ , we obtain that  $f(x) + \sigma \sum_{j=1}^d \bar{y}_j \leq f(x) + \sigma \sum_{j=1}^d y_j$ . Therefore (5) is equivalent to

$$\text{Minimize } f(x) + \sigma \sum_{j=1}^d y_j \text{ subject to } y_j \geq \pi(g_j(x)) \text{ for } j = 1, \dots, d \text{ and } x \in \mathcal{C}. \quad (6)$$

As  $\pi(t) = 0$  if  $t < 0$ , (6) is equivalent to

$$\text{Minimize } f(x) + \sigma \sum_{j=1}^d y_j \text{ subject to } y_j \geq 0, y_j \geq \pi(g_j(x)) \text{ for } j = 1, \dots, d \text{ and } x \in \mathcal{C}. \quad (7)$$

If  $y_j \geq 0$  and  $y_j \geq \pi(g_j(x))$ , since both  $y_j$  and  $\pi(g_j(x))$  belong to  $\mathbb{R}_+$  and the inverse of  $\pi$  onto  $\mathbb{R}_+$  is increasing, we obtain that  $\pi^{-1}(y_j) \geq g_j(x)$ . Reciprocally, if  $\pi^{-1}(y_j) \geq g_j(x)$  and  $y_j \geq 0$

we obtain that  $y_j \geq \pi(g_j(x))$  or  $g_j(x) \leq 0$  in which case  $y_j \geq \pi(g_j(x))$  also holds. Therefore, (7) is equivalent to

$$\begin{aligned} & \text{Minimize} && \Phi(x, y) := f(x) + \sigma \sum_{j=1}^d y_j \\ & \text{subject to} && x \in \mathcal{C}, \\ & && g_j(x) - \pi^{-1}(y_j) \leq 0 \text{ for } j = 1, \dots, d, \\ & && y \geq 0. \end{aligned} \tag{8}$$

The theorem below shows that a solution to the original problem (1) can be found by solving its smooth reformulation (8). (The reciprocal is also true, as shown in the following theorem.)

**Theorem 2.1.** *Suppose that  $(x^*, y^*)$  is a local (global) minimizer of (8). Then,  $y_j^* = \pi(g_j(x^*))$  for all  $j = 1, \dots, d$  and  $x^*$  is a local (global) minimizer of (1). Reciprocally, if  $x^*$  is a local (global) minimizer of (1) and  $y_j^* = \pi(g_j(x^*))$  for all  $j = 1, \dots, d$ , then  $(x^*, y^*)$  is a local (global) minimizer of (8).*

*Proof.* Suppose that  $(x^*, y^*)$  is a local minimizer of (8). If  $y_j^* > \pi(g_j(x^*))$  and  $y_j^* > y_j \geq \pi(g_j(x^*))$ , we have that  $\pi^{-1}(y_j^*) > \pi^{-1}(y_j) \geq g_j(x^*)$ . Therefore, if  $y_i = y_i^*$  for all  $i \neq j$ , we have that  $(x^*, y)$  is a feasible point of (8) and  $\Phi(x^*, y) < \Phi(x^*, y^*)$ . Therefore  $(x^*, y^*)$  would not be a local minimizer of (8). This implies that, necessarily,  $y_j^* = \pi(g_j(x^*))$  for all  $j = 1, \dots, d$ .

Let  $\varepsilon > 0$  be such that for all  $(x, y)$  such that  $x \in \mathcal{C}$ ,  $y \geq 0$ ,  $g_j(x) \leq \pi^{-1}(y_j)$ ,  $j = 1, \dots, d$ ,  $\|x - x^*\| \leq \varepsilon$  and  $\|y - y^*\|_\infty \leq \varepsilon$ , we have that  $f(x) + \sigma \sum_{j=1}^d y_j \geq f(x^*) + \sigma \sum_{j=1}^d y_j^*$ . By the first part of the proof,  $y_j^* = \pi(g_j(x^*))$ , therefore,  $f(x) + \sigma \sum_{j=1}^d y_j \geq f(x^*) + \sigma \sum_{j=1}^d \pi(g_j(x^*))$ . Let  $\varepsilon_1 \in (0, \varepsilon]$  be such that  $\|x - x^*\| \leq \varepsilon_1$  implies that  $|\pi(g_j(x)) - \pi(g_j(x^*))| \leq \varepsilon$  for all  $j = 1, \dots, d$ . Then, if  $\|x - x^*\| \leq \varepsilon_1$ ,  $f(x) + \sigma \sum_{j=1}^d \pi(g_j(x)) \geq f(x^*) + \sigma \sum_{j=1}^d \pi(g_j(x^*))$ . This implies that  $x^*$  is a local minimizer of (1).

Reciprocally, suppose that  $x^*$  is a local minimizer of (1). Therefore, there exists  $\varepsilon > 0$  such that for all  $x \in \mathcal{C}$  such that  $\|x - x^*\| \leq \varepsilon$ ,  $f(x^*) + \sigma \sum_{j=1}^d \pi(g_j(x^*)) \leq f(x) + \sigma \sum_{j=1}^d \pi(g_j(x))$ . Therefore,  $f(x^*) + \sigma \sum_{j=1}^d y_j^* \leq f(x) + \sigma \sum_{j=1}^d \pi(g_j(x))$  for all  $x \in \mathcal{C}$  such that  $\|x - x^*\| \leq \varepsilon$ . Thus,  $f(x^*) + \sigma \sum_{j=1}^d y_j^* \leq f(x) + \sigma \sum_{j=1}^d y_j$  for all  $x \in \mathcal{C}$  such that  $\|x - x^*\| \leq \varepsilon$  if  $y_j \geq 0$  and  $y_j \geq \pi(g_j(x))$  for all  $j = 1, \dots, d$ . But, due to the increasing property of  $\pi^{-1}$  onto  $\mathbb{R}_+$ ,  $y_j \geq 0$  and  $y_j \geq \pi(g_j(x))$  is equivalent to  $y_j \geq 0$  and  $\pi^{-1}(y_j) \geq g_j(x)$ . Therefore,  $f(x^*) + \sigma \sum_{j=1}^d y_j^*$  is less than or equal to  $f(x) + \sigma \sum_{j=1}^d y_j$  for all  $(x, y)$  satisfying  $x \in \mathcal{C}$ ,  $\|x - x^*\| \leq \varepsilon$ , and the constraints of (8).

The equivalence proof with respect global minimizers is similar.  $\square$

### 3 Optimality conditions

Assume that

$$\mathcal{C} = \{x \in \mathbb{R}^n \mid H(x) = 0 \text{ and } G(x) \leq 0\}, \tag{9}$$

where  $H : \mathbb{R}^n \rightarrow \mathbb{R}^{n_H}$  and  $G : \mathbb{R}^n \rightarrow \mathbb{R}^{n_G}$  are continuously differentiable. If a local minimizer  $(x^*, y^*)$  of (8) satisfies a constraint qualification then the KKT conditions hold at  $(x^*, y^*)$ . By Theorem 2.1, if  $x^*$  is a local minimizer of (1),  $y_j^* = \pi(g_j(x^*))$  for all  $j = 1, \dots, d$ , and a constraint qualification for (8) is fulfilled, the KKT conditions are satisfied too. The KKT conditions for (8)

obviously involve both  $x^*$  and  $y^*$ . Since  $y$  is merely an auxiliary variable, it is interesting to derive KKT-like conditions in which only the primal variables  $x^*$  are involved. This is the objective of Theorem 3.1 below, which, in turn, generalizes [13, Corollary 2.2] and [22] using straightforward optimization arguments.

**Theorem 3.1.** *Let  $x^*$  be a local minimizer of (1) and assume that  $\pi$  satisfies Assumptions A1 and A2. Define*

$$I(x^*) := \{j \mid g_j(x^*) < 0\}, J(x^*) := \{j \mid g_j(x^*) = 0\}, \text{ and } K(x^*) := \{j \mid g_j(x^*) > 0\}. \quad (10)$$

Then,  $x^*$  is a local minimizer of the problem

$$\text{Minimize } f(x) + \sigma \sum_{j \in K(x^*)} \pi(g_j(x)) \text{ subject to } x \in \mathcal{C} \text{ and } g_j(x) \leq 0 \forall j \in J(x^*). \quad (11)$$

Moreover, if  $\mathcal{C}$  is defined by  $H(x) = 0$  and  $G(x) \leq 0$ , where  $H : \mathbb{R}^n \rightarrow \mathbb{R}^{n_H}$  and  $G : \mathbb{R}^n \rightarrow \mathbb{R}^{n_G}$  are continuously differentiable, and the system of equalities and inequalities

$$H(x) = 0, G(x) \leq 0, g_j(x) \leq 0, \forall j \in J(x^*) \quad (12)$$

satisfies a constraint qualification at  $x^*$  then we have that  $x^*$  satisfies the KKT conditions of problem (11).

*Proof.* Let  $y^* \geq 0$  be such that  $(x^*, y^*)$  is a local minimizer of (8). By Theorem 2.1,  $y_j^* = \pi(g_j(x^*))$  for all  $j = 1, \dots, d$ . If  $j \in I(x^*)$ , we have that  $g_j(x) < 0$  for all  $x$  in a neighborhood of  $x^*$ . Moreover,  $y_j^* = 0$  in this case. Therefore,  $(x^*, y^*)$  is a local minimizer of

$$\text{Minimize } f(x) + \sigma \sum_{j \notin I(x^*)} y_j \text{ subject to } x \in \mathcal{C}, y_j \geq \pi(g_j(x)) \text{ and } y_j \geq 0 \forall j \notin I(x^*).$$

So,  $(x^*, y^*)$  is a local minimizer of

$$\text{Minimize } f(x) + \sigma \sum_{j \notin I(x^*)} y_j \text{ subject to } x \in \mathcal{C}, y_j = \pi(g_j(x)) \text{ and } y_j \geq 0 \forall j \notin I(x^*).$$

If  $j \in K(x^*)$ , we have that  $g_j(x) > 0$  in a neighborhood of  $x^*$  and  $\pi(g_j(x))$  is continuous and differentiable on that neighborhood. Therefore,  $x^*$  is a local minimizer of the locally smooth problem given by

$$\begin{aligned} & \text{Minimize } f(x) + \sigma \left[ \sum_{j \in J(x^*)} y_j + \sum_{j \in K(x^*)} \pi(g_j(x)) \right] \\ & \text{subject to } x \in \mathcal{C}, y_j = \pi(g_j(x)) \text{ and } y_j \geq 0 \forall j \in J(x^*). \end{aligned}$$

Since  $g_j(x^*) \leq 0$  for all  $j \in J(x^*)$ ,  $x^*$  is also a local minimizer of

$$\text{Minimize } f(x) + \sigma \left[ \sum_{j \in J(x^*)} y_j + \sum_{j \in K(x^*)} \pi(g_j(x)) \right]$$

subject to  $x \in \mathcal{C}$ ,  $g_j(x) \leq 0$ ,  $y_j = \pi(g_j(x))$  and  $y_j \geq 0 \forall j \in J(x^*)$ .

Moreover, since  $y_j^* = 0$  for all  $j \in J(x^*)$ ,  $x^*$  is a local minimizer of

Minimize  $f(x) + \sigma \sum_{j \in K(x^*)} \pi(g_j(x))$  subject to  $x \in \mathcal{C}$ ,  $g_j(x) \leq 0$ ,  $y_j = \pi(g_j(x))$  and  $y_j \geq 0 \forall j \in J(x^*)$ .

Clearly, the variables  $y_j$  for  $j \in J(x^*)$  play no role in this problem, so  $x^*$  is a local minimizer of

Minimize  $f(x) + \sigma \sum_{j \in K(x^*)} \pi(g_j(x))$  subject to  $x \in \mathcal{C}$  and  $g_j(x) \leq 0 \forall j \in J(x^*)$ .

Then, if the constraints that define  $\mathcal{C}$  together with the constraints  $g_j(x) \leq 0$ ,  $j \in J(x^*)$ , satisfy a constraint qualification at  $x^*$ , the KKT conditions of (11) hold at  $x^*$ . This completes the proof.  $\square$

Observe that (11) is a constrained optimization problem of the form

$$\text{Minimize } f_{\text{obj}}(w) \text{ subject to } h_{\text{E}}(w) = 0 \text{ and } h_{\text{I}}(w) \leq 0, \quad (13)$$

where  $f_{\text{obj}} : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $h_{\text{E}} : \mathbb{R}^n \rightarrow \mathbb{R}^{n_{\text{E}}}$ , and  $h_{\text{I}} : \mathbb{R}^n \rightarrow \mathbb{R}^{n_{\text{I}}}$  are continuously differentiable. It is well known that, independently of constraint qualifications, every local minimizer  $w^*$  of (13) satisfies the Approximate KKT (AKKT) optimality condition [2, 7], which means that, given  $\varepsilon_{\text{sol}} > 0$ ,  $\varepsilon_{\text{kkt}} > 0$ ,  $\varepsilon_{\text{feas}} > 0$ , and  $\varepsilon_{\text{comp}} > 0$ , there exist  $w \in \mathbb{R}^n$ ,  $\lambda \in \mathbb{R}^{n_{\text{E}}}$ , and  $\mu \in \mathbb{R}_+^{n_{\text{I}}}$  such that  $\|w - w^*\| \leq \varepsilon_{\text{sol}}$ ,

$$\|h_{\text{E}}(w)\|_{\infty} \leq \varepsilon_{\text{feas}}, \quad \|h_{\text{I}}(w)_+\|_{\infty} \leq \varepsilon_{\text{feas}}, \quad (14)$$

$$\mu_j \leq \varepsilon_{\text{comp}} \text{ for all } j \text{ such that } [h_{\text{I}}(w)]_j < -\varepsilon_{\text{feas}}, \quad (15)$$

and

$$\|\nabla f_{\text{obj}}(w) + \nabla h_{\text{E}}(w)\lambda + \nabla h_{\text{I}}(w)\mu\| \leq \varepsilon_{\text{kkt}}. \quad (16)$$

If  $w \in \mathbb{R}^n$  is such that there exist  $\lambda \in \mathbb{R}^{n_{\text{E}}}$  and  $\mu \in \mathbb{R}_+^{n_{\text{I}}}$  fulfilling (14), (15), and (16), we say that  $w$  is an  $(\varepsilon_{\text{feas}}/\varepsilon_{\text{comp}}/\varepsilon_{\text{kkt}})$ -AKKT point of (13).

Applying the AKKT definition to (11) we obtain the following result, independently of constraint qualifications.

**Theorem 3.2.** *Let  $x^*$  be a local minimizer of (1) and assume that  $\pi$  satisfies Assumptions A1 and A2. Let  $\mathcal{C}$  be defined by  $H(x) = 0$  and  $G(x) \leq 0$ , where  $H : \mathbb{R}^n \rightarrow \mathbb{R}^{n_{\text{H}}}$  and  $G : \mathbb{R}^n \rightarrow \mathbb{R}^{n_{\text{G}}}$  are continuously differentiable. Assume that  $\varepsilon_{\text{sol}} > 0$ ,  $\varepsilon_{\text{kkt}} > 0$ ,  $\varepsilon_{\text{feas}} > 0$ , and  $\varepsilon_{\text{comp}} > 0$  are arbitrary. Then, there exist  $x \in \mathbb{R}^n$ ,  $\lambda_{\text{H}} \in \mathbb{R}^{n_{\text{H}}}$ ,  $\mu_{\text{G}} \in \mathbb{R}_+^{n_{\text{G}}}$ , and  $\mu_j \in \mathbb{R}_+$  for all  $j$  such that  $g_j(x^*) = 0$  such that*

$$\|x - x^*\| \leq \varepsilon_{\text{sol}},$$

$$\|H(x)\| \leq \varepsilon_{\text{feas}}, \quad \|G(x)_+\| \leq \varepsilon_{\text{feas}},$$

$$g_j(x) \leq \varepsilon_{\text{feas}} \text{ for all } j \text{ such that } g_j(x^*) = 0,$$

$$\left\| \nabla f(x) + \sigma \sum_{g_j(x^*) > 0} \pi'(g_j(x)) \nabla g_j(x) + \nabla H(x) \lambda_{\text{H}} + \nabla G(x) \mu_{\text{G}} + \sum_{g_j(x^*) = 0} \mu_j \nabla g_j(x) \right\| \leq \varepsilon_{\text{kkt}},$$

$$[\mu_G]_j \leq \varepsilon_{\text{comp}} \text{ for all } j \text{ such that } G_j(x) < -\varepsilon_{\text{feas}},$$

and

$$\mu_j \leq \varepsilon_{\text{comp}} \text{ for all } j \text{ such that } g_j(x) < -\varepsilon_{\text{feas}}.$$

*Proof.* By Theorem 3.1,  $x^*$  is a local minimizer of (11). Then, the thesis follows from (14), (15), and (16).  $\square$

## 4 Model-Intensive algorithms

As we mentioned in Section 2, problem (8) can be solved using standard methods for smooth constrained optimization. In many applications, the cost of evaluating  $f(x)$  exceeds, by far, the cost of evaluating the constraints. For these situations “model intensive” (MI) methods are recommendable. At each iteration of such an algorithm, a model of the objective function is built and this model is minimized using the original constraints (instead of their linear approximations as in SQP and other popular approaches). The idea is to exploit the model as much as possible so that, hopefully, it will not be necessary to evaluate the expensive function many times.

The most popular model for an arbitrary sufficiently smooth function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  comes from considering its  $p$ -th Taylor approximation

$$T_p(\bar{x}, x) := \sum_{j=1}^p \frac{1}{j!} \left( (x - \bar{x}_1) \frac{\partial}{\partial x_1} + \cdots + (x - \bar{x}_n) \frac{\partial}{\partial x_n} \right)^j f(\bar{x}), \quad (17)$$

so that

$$f(x) = f(\bar{x}) + T_p(\bar{x}, x) + o(\|x - \bar{x}\|^{p+1}). \quad (18)$$

Observe that  $T_1(\bar{x}, x) = \nabla f(\bar{x})^T(x - \bar{x})$  and  $T_2(\bar{x}, x) = \nabla f(\bar{x})^T(x - \bar{x}) + \frac{1}{2}(x - \bar{x})^T \nabla^2 f(\bar{x})(x - \bar{x})$ . Consequently, the model for  $f(x) + \sigma \sum_{j=1}^d y_j$  will be  $f(\bar{x}) + T_p(\bar{x}, x) + \sigma \sum_{j=1}^d y_j$ .

In this section  $M_{\bar{x}, \bar{y}}(x, y)$  will be a model of  $\Phi(x, y) - \Phi(\bar{x}, \bar{y})$  for all  $\bar{x}, x \in \mathbb{R}^n$  and  $\bar{y}, y \in \mathbb{R}^d$ . Therefore,  $\Phi(x, y) \approx \Phi(\bar{x}, \bar{y}) + M_{\bar{x}, \bar{y}}(x, y)$ . Let us define, for all  $\delta \geq 0$ ,

$$\mathcal{C}_\delta = \{x \in \mathbb{R}^n \mid \|H(x)\|_\infty \leq \delta \text{ and } \|G(x)_+\|_\infty \leq \delta\}, \quad (19)$$

$$D_\delta = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^d \mid x \in \mathcal{C}_\delta, y \geq 0, \text{ and } g_j(x) - \pi^{-1}(y_j) \leq \delta, j = 1, \dots, d\}, \quad (20)$$

and  $D = D_0$ . The following algorithm is similar to Algorithm 3.1 of [9].

**Algorithm 4.1.** Assume that  $p \in \{1, 2, 3, \dots\}$ ,  $\eta > 0$ ,  $\alpha > 0$ ,  $\rho_{\min} > 0$ ,  $\tau_2 \geq \tau_1 > 1$ ,  $\theta > 0$ , and  $(x^0, y^0) \in D_{\eta/4}$  are given with  $y_j^0 = \pi(g_j(x^0))$ ,  $j = 1, \dots, d$ . Initialize  $k \leftarrow 0$ .

**Step 1.** Set  $\rho \leftarrow 0$ .

**Step 2.** Compute  $x^{\text{trial}} \in \mathbb{R}^n$  and  $y^{\text{trial}} \in \mathbb{R}^d$  such that

$$(x^{\text{trial}}, y^{\text{trial}}) \in D_{\eta_k} \text{ with } \eta_k = \eta \left( \frac{k+1}{k+2} \right) \quad (21)$$

and

$$M_{x^k, y^k}(x^{\text{trial}}, y^{\text{trial}}) + \rho \|(x^{\text{trial}} - x^k, y^{\text{trial}} - y^k)\|^{p+1} \leq 0. \quad (22)$$

**Step 3.** Test the condition

$$\Phi(x^{\text{trial}}, y^{\text{trial}}) \leq \Phi(x^k, y^k) - \alpha \|(x^{\text{trial}} - x^k, y^{\text{trial}} - y^k)\|^{p+1}. \quad (23)$$

If (23) holds, accept the trial point  $(x^{\text{trial}}, y^{\text{trial}})$ , define  $\rho_k = \rho$ ,  $x^{k+1} = x^{\text{trial}}$ ,  $y^{k+1} = y^{\text{trial}}$ , set  $k \leftarrow k + 1$ , and go to Step 1. Otherwise, update  $\rho \leftarrow \max\{\rho_{\min}, \tau\rho\}$  with  $\tau \in [\tau_1, \tau_2]$ , and go to Step 2.

Under the model-intensive strategy, conditions (21) and (22) may be obtained by solving the subproblem

$$\text{Minimize } M_{x^k, y^k}(x, y) + \rho \|(x - x^k, y - y^k)\|^{p+1} \text{ subject to } (x, y) \in D, \quad (24)$$

with rather high precision, since we assume that the cost of evaluating the model is negligible with respect to the cost of evaluating  $f$ . Note that the feasibility required at  $x^{\text{trial}}$  is looser than the precision with which  $x^k$  satisfies those constraints. Therefore, it is plausible to require the fulfillment of (21) and (22) by means of a standard constrained optimization algorithm. This means that, although conditions (22) and (23) are trivially satisfied by  $x^{\text{trial}} = x^k$  and  $y^{\text{trial}} = y^k$ , it is reasonable to expect that a nontrivial solution with decrease of  $\Phi$  could be obtained with non-null increments.

Complexity results for Algorithm 4.1 are presented below. Before that, the standard assumptions (see, for example, [8, 12]) are given.

**Assumption A3.** For all  $(x^k, y^k)$  generated by Algorithm 4.1, we have that

$$\nabla\Phi(x^k, y^k) = \nabla M_{x^k, y^k}(x, y)|_{(x, y) = (x^k, y^k)}.$$

**Assumption A4.** There exists  $L_1 > 0$  such that, for every  $(x^k, y^k)$  calculated by Algorithm 4.1 and for every  $(x^{\text{trial}}, y^{\text{trial}})$  satisfying (22), we have that

$$\Phi(x^{\text{trial}}, y^{\text{trial}}) \leq \Phi(x^k, y^k) + M_{x^k, y^k}(x^{\text{trial}}, y^{\text{trial}}) + L_1 \|(x - x^{\text{trial}}, y - y^{\text{trial}})\|^{p+1}.$$

Assumption A4 is valid if  $M$  is obtained using the  $p$ -th Taylor approximation and the derivatives of order  $p$  of  $f$  are Lipschitz continuous. See [3, 8, 12].

**Assumption A5.** There exists  $L_2 > 0$  such that, for every  $x^k, y^k, x^{k+1},$  and  $y^{k+1}$  obtained by Algorithm 4.1, we have that

$$\|\nabla_{x, y}\Phi(x^{k+1}, y^{k+1}) - \nabla_{x, y}M_{x^k, y^k}(x^{k+1}, y^{k+1})\| \leq L_2 \|(x^{k+1} - x^k, y^{k+1} - y^k)\|^p.$$

As in the case of Assumption A4, Assumption A5 also holds if we use Taylor  $p$ -th approximations for computing  $M$  and the  $p$ -th order derivatives of  $f$  are Lipschitz-continuous.

Assumption A6 below says that (24) must be approximately solved. Before stating this assumption let us recall that, assuming that  $\|(x - x^k, y - y^k)\|^{p+1}$  is continuously differentiable with respect to  $(x, y)$ , (24) is a constrained optimization problem of the form (13).



**Assumption A6.** We say that this assumption holds at iteration  $k$  of Algorithm 4.1 if the function  $\|(x - x^k, y - y^k)\|^{p+1}$  is continuously differentiable with respect to  $(x, y)$  and  $(x^{\text{trial}}, y^{\text{trial}})$  is an  $(\varepsilon_{\text{feas}}/\varepsilon_{\text{comp}}/\varepsilon_{\text{kkt}})$ -AKKT point of (24) with  $\varepsilon_{\text{feas}} = \eta_k$ ,  $\varepsilon_{\text{comp}} = \eta$ , and  $\varepsilon_{\text{kkt}} = \theta\|(x^{\text{trial}} - x^k, y^{\text{trial}} - y^k)\|^p$ .

**Theorem 4.1.** Assume that the sequence  $\{(x^k, y^k)\}$  is generated by Algorithm 4.1,  $\Phi_{\text{target}} \in \mathbb{R}$ ,  $\varepsilon > 0$ , and  $L = \max\{L_1, L_2\}$ . Define

$$K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta) = \left\lfloor (\Phi(x^0, y^0) - \Phi_{\text{target}}) \left( \frac{\alpha^{p/(p+1)}\varepsilon}{L + \tau_2(L + \alpha)(p + 1) + \theta} \right)^{-(p+1)/p} \right\rfloor. \quad (25)$$

Suppose that Assumptions A3, A4, A5, and A6 hold for all

$$k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta).$$

Then, the number of iterations  $k$  such that

$$\Phi(x^{k+1}, y^{k+1}) > \Phi_{\text{target}}$$

and  $(x^{k+1}, y^{k+1})$  is not an  $(\eta_k/\eta/\varepsilon)$ -AKKT point of (8) is bounded above by  $K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$ . Moreover, the number of functional evaluations per iteration is bounded above by

$$\left\lceil \log_{\tau_1} \left( \frac{\tau_2(L + \alpha)}{\rho_{\min}} \right) \right\rceil + 1.$$

*Proof.* The desired result follows as in [9, Thm.3.1].  $\square$

**Theorem 4.2.** Suppose that Assumptions A3, A4, and A5 hold, and the function  $\|(x - x^k, y - y^k)\|^{p+1}$  is continuously differentiable with respect to  $(x, y)$ . Assume, moreover, that for all  $k = 0, 1, 2, \dots$  and  $\varepsilon_{\text{feas}} > 0$ ,  $\varepsilon_{\text{comp}} > 0$ , and  $\varepsilon_{\text{kkt}} > 0$ , we are able to compute, using a suitable algorithm, an  $(\varepsilon_{\text{feas}}/\varepsilon_{\text{comp}}/\varepsilon_{\text{kkt}})$ -AKKT point of (24) such that (22) holds. Then, there are two possibilities:

1. Assumption A6 holds for all  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$ .
2. There exists  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$  such that  $(x^k, y^k)$  is an AKKT point of (8).

*Proof.* Assume that  $(x^{k,\ell}, y^{k,\ell})$  is a sequence generated by an algorithm that, when applied to the problem

$$\text{Minimize } M_{x^k, y^k}(x, y) + \rho\|(x - x^k, y - y^k)\|^{p+1} \text{ subject to } (x, y) \in D_{\eta_{k-1}}, \quad (26)$$

satisfies the hypotheses of the theorem. By construction,  $(x^k, y^k) \in D_{\eta_{k-1}}$ . Therefore the minimum of  $M_{x^k, y^k}(x, y) + \rho\|(x - x^k, y - y^k)\|^{p+1}$  onto  $D_{\eta_{k-1}}$  is non-positive, as well as the minimum of  $M_{x^k, y^k}(x, y) + \rho\|(x - x^k, y - y^k)\|^{p+1}$  onto  $D_{\eta_k}$ .

By the fulfillment of the AKKT optimality conditions, we have two possibilities:

1. There exists an iterate  $(x^{k,\ell}, y^{k,\ell})$  which is an  $(\varepsilon_{\text{feas}}/\varepsilon_{\text{comp}}/\varepsilon_{\text{kkt}})$ -AKKT point of (24) with  $\varepsilon_{\text{feas}} = \eta_k$ ,  $\varepsilon_{\text{comp}} = \eta$ , and  $\varepsilon_{\text{kkt}} = \theta\|(x^{k,\ell} - x^k, y^{k,\ell} - y^k)\|^p$ .
2. There exist infinitely many iterates  $(x^{k,\ell}, y^{k,\ell})$  that are  $(\varepsilon_{\text{feas}}/\varepsilon_{\text{comp}}/\varepsilon_{\text{kkt}})$ -AKKT points of (24) with  $\varepsilon_{\text{feas}} = 1/\ell < \eta_k$ ,  $\varepsilon_{\text{comp}} = 1/\ell < \eta$ , and  $\varepsilon_{\text{kkt}} = 1/\ell$  but

$$\theta\|(x^{k,\ell} - x^k, y^{k,\ell} - y^k)\| < 1/\ell. \quad (27)$$

In the first case we can choose  $x^{\text{trial}} = x^{k,\ell}$  and  $y^{\text{trial}} = y^{k,\ell}$ , satisfying Assumption A6 and (22). In the second case, (27) implies that

$$\lim_{\ell \rightarrow \infty} (x^{k,\ell}, y^{k,\ell}) = (x^k, y^k).$$

Therefore  $(x^k, y^k)$  satisfies the AKKT optimality condition for (26). Since the first derivatives of  $\Phi(x, y)$  at  $(x^k, y^k)$  coincide with the first derivatives of the objective function of (26), it turns out that  $(x^k, y^k)$  was an AKKT point for (26). In other words, Assumption A6 did not hold because  $(x^k, y^k)$  already was an approximate solution of the original problem. This completes the proof.  $\square$

Theorems 4.1 and 4.2 say that after at most the number of iterations given by (25) one finds an iterate that satisfies the constraints of (8) with tolerance  $\eta$  such that the objective function value is smaller than or equal to  $\Phi_{\text{target}}$ ; or, alternatively, we find an iterate that satisfies KKT conditions with tolerance  $\eta$  for feasibility, tolerance  $\eta$  for complementarity, and tolerance  $\varepsilon$  for optimality. The conclusion of these theorems is that, ultimately, Assumption A6 is not necessary since, if it holds for all  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$ , it guarantees that an approximate solution is found for some  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$ . But, if Assumption A6 does not hold for some  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$ , the iterate  $x^k$  is an approximate solution. This is stated in the following Corollary.

**Corollary 4.3.** *Suppose that the hypotheses of Theorem 4.2 hold. Then, one of the following statements is true:*

1. *There exists  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$ , such that*

$$\Phi(x^{k+1}, y^{k+1}) \leq \Phi_{\text{target}}$$

*and  $(x^{k+1}, y^{k+1})$  is  $(\eta_k/\eta/\varepsilon)$ -AKKT point of (8).*

2. *There exists  $k \leq K(x^0, y^0, \Phi_{\text{target}}, \alpha, p, L, \tau_2, \theta)$  such that  $(x^k, y^k)$  is an AKKT point of (8).*

## 5 Linearly-constrained problems

In this section, we consider the case in which  $G(x)$ ,  $H(x)$ , and  $g_j(x)$ ,  $j = 1, \dots, d$ , are affine functions. Thus, their first derivatives are constant. Consequently, we denote

$$A_G = G'(x), \quad A_H = H'(x), \quad a_j^T = g_j'(x), \quad j = 1, \dots, d,$$

for all  $x \in \mathbb{R}^n$ . Therefore, for all  $\bar{x} \in \mathbb{R}^n$ ,

$$G(x) = A_G(x - \bar{x}) + G(\bar{x}), \quad H(x) = A_H(x - \bar{x}) + H(\bar{x}), \quad g_j(x) = a_j^T(x - \bar{x}) + g_j(\bar{x}), \quad j = 1, \dots, d.$$

The constraints  $H(x) = 0$  and  $G(x) \leq 0$  are now linear but the constraints  $g_j(x) - \pi^{-1}(y_j) \leq 0$  of (8) are not, due to the nonlinearity of  $\pi^{-1}$ . Therefore, the set  $\mathcal{C}$  is a polytope but the set  $D_0$ , defined by (19) and (20) is not.

Of course, we may use Algorithm 4.1 for solving (8) in this case, but the presence of linearity suggests that linear constraints could be satisfied exactly at the solution of each subproblem. If this is possible and if the model is also linear, each subproblem would be a linear programming problem and, if the model is quadratic, we would have a quadratic programming subproblem.

For achieving this improvement, the nonlinear constraints  $g_j(x) - \pi^{-1}(y_j) \leq 0$  must be linearized. This amounts to replace, at each subproblem, each constraint  $g_j(x) - \pi^{-1}(y_j) \leq 0$  with

$$g_j(x) - [\pi^{-1}(y_j^k) + (\pi^{-1})'(y_j^k)(y_j - y_j^k)] \leq 0. \quad (28)$$

By the convexity of the function  $\pi^{-1}$ , the fulfillment of (28) implies that  $g_j(x) - \pi^{-1}(y_j) \leq 0$ . Therefore, the feasible set defined by  $H(x) = 0$ ,  $G(x) \leq 0$ , and (28) for  $j = 1, \dots, d$  is contained in  $D_0$ .

**Algorithm 5.1.** This algorithm is identical to Algorithm 4.1 except that condition (21) is replaced by

$$(x^{\text{trial}}, y^{\text{trial}}) \in \mathcal{C} \text{ and } g_j(x) - [\pi^{-1}(y_j^k) + (\pi^{-1})'(y_j^k)(y_j - y_j^k)] \leq 0, \quad j = 1, \dots, d. \quad (29)$$

Of course, the parameter  $\eta$  is not necessary anymore since we assume that linear constraints can be satisfied exactly using well-established constrained optimization methods.

Assumptions A3, A4, and A5 stand exactly in the same way as in Section 4. However, Assumption A6 needs to be replaced in order to take into account that now the subproblem has only linear constraints and that  $d$  constraints are linear approximations of the true ones.

**Assumption A7.** *At each iteration  $k$  of Algorithm 5.1, the function  $\|(x - x^k, y - y^k)\|^{p+1}$  is continuously differentiable with respect to  $(x, y)$  and  $(x^{k+1}, y^{k+1})$  satisfies the KKT conditions for the linearly constrained problem that consists of minimizing  $M_{x^k, y^k}(x, y) + \rho\|(x - x^k, y - y^k)\|^{p+1}$  subject to  $H(x) = 0$ ,  $G(x) \leq 0$ , and  $g_j(x) - [\pi^{-1}(y_j^k) + (\pi^{-1})'(y_j^k)(y_j - y_j^k)] \leq 0$ ,  $j = 1, \dots, d$ .*

Assumption A7 is plausible because every minimizer of linearly constrained optimization problems satisfy KKT conditions.

According to Assumption A7 the increment  $(x^{k+1}, y^{k+1})$  at each iteration of Algorithm 5.1 must satisfy the following conditions:

$$\nabla_x \left[ M_{x^k, y^k}(x^{k+1}, y^{k+1}) + \rho\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^{p+1} \right] + A_H^T \lambda + A_G^T \mu + \sum_{j=1}^d a_j \beta_j = 0, \quad (30)$$

$$\rho \frac{\partial}{\partial y_j} \|(x^{k+1} - x^k, y^{k+1} - y^k)\|^{p+1} + \sigma_j - (\pi^{-1})'(y_j^{k+1}) \beta_j - \gamma_j = 0, \quad j = 1, \dots, d, \quad (31)$$

$$H(x^{k+1}) = 0, \quad (32)$$

$$G(x^{k+1}) \leq 0, \quad (33)$$

$$g_j(x^{k+1}) - \left[ \pi^{-1}(y_j^k) + (\pi^{-1})'(y^k)(y_j^{k+1} - y_j^k) \right] \leq 0, \quad j = 1, \dots, d, \quad (34)$$

$$\beta_j = 0 \text{ whenever } g_j(x^{k+1}) - \left[ \pi^{-1}(y_j^k) + (\pi^{-1})'(y^k)(y_j^{k+1} - y_j^k) \right] < 0, \quad (35)$$

$$y^{k+1} \geq 0, \quad (36)$$

$$\mu_j = 0 \text{ for all } j \text{ such that } G(x^{k+1})_j < 0, \quad (37)$$

$$\gamma_j = 0 \text{ for all } j \text{ such that } y_j^{k+1} > 0, \quad (38)$$

$$\mu, \beta, \gamma \geq 0. \quad (39)$$

Recall that  $(x, y)$  satisfies the KKT conditions of (8) if conditions (30)–(39) hold with  $(x, y)$  replacing  $(x^{k+1}, y^{k+1})$ , except that (30) and (31) must be replaced by

$$\nabla f(x) + A_{\text{H}}^T \lambda + A_{\text{G}}^T \mu + \sum_{j=1}^d a_j \beta_j = 0 \quad (40)$$

and

$$\sigma_j - (\pi^{-1})'(y_j) \beta_j - \gamma_j = 0, \quad j = 1, \dots, d, \quad (41)$$

whereas (34) and (35) must be replaced by

$$g_j(x) - \pi^{-1}(y_j) \leq 0 \quad (42)$$

and

$$\beta_j = 0 \text{ whenever } g_j(x) - \pi^{-1}(y_j) < 0. \quad (43)$$

Therefore, the question is whether (31)–(39) implies some relaxed version of (40)–(43) for  $x = x^{k+1}$  and  $y = y^{k+1}$ .

As in [9, Lemma 3.2], by Assumption A4, it follows that the values of  $\rho$  used at each subproblem of Algorithm 5.1 are bounded. By Assumption A5, (30) implies that

$$\|\nabla f(x^{k+1}) + A_{\text{H}}^T \lambda + A_{\text{G}}^T \mu + \sum_{j=1}^d a_j \beta_j\| \leq O(\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^p). \quad (44)$$

By the limitation of  $\rho$  and the fact that  $|\frac{\partial}{\partial y_j}| \|(x - x^k, y - y^k)\|^{p+1} \leq O(\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^p)$ , (31) implies that

$$|\sigma_j - (\pi^{-1})'(y_j^{k+1}) \beta_j - \gamma_j| \leq O(\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^p), \quad j = 1, \dots, d. \quad (45)$$

By (34) and the convexity of  $\pi^{-1}$ , we have that (42) holds with  $x = x^{k+1}$  and  $y = y^{k+1}$ . Thus,

$$g_j(x^{k+1}) - \pi^{-1}(y_j^{k+1}) \leq 0, \quad j = 1, \dots, d. \quad (46)$$

For proving approximate complementarity we need a new assumption regarding the replacement of  $\pi^{-1}$  with its linear approximation.

**Assumption A8.** *There exists  $c_\pi > 0$  such that, for all  $z, y \geq 0$ ,*

$$|\pi^{-1}(y) - [\pi^{-1}(z) + (\pi^{-1})'(z)(z - y)]| \leq c_\pi |y - z|^2.$$

By Assumption A8, we have that  $g_j(x^{k+1}) - \pi^{-1}(y_j^{k+1}) < -c_\pi |y_j^{k+1} - y_j^k|^2$  implies that  $g_j(x^{k+1}) - [\pi^{-1}(y_j^k) + (\pi^{-1})'(y_j^k)(y_j^{k+1} - y_j^k)] < 0$ ; so, by (35),  $\beta_j = 0$ .

**Theorem 5.1.** *Suppose that Assumptions A3, A4, A5, A7, and A8 hold,  $L = \max\{L_1, L_2\}$ , the sequence  $\{(x^k, y^k)\}$  is generated by Algorithm 5.1,  $\Phi_{\text{target}} \in \mathbb{R}$ , and  $\varepsilon > 0$ . Then, the number of iterations  $k$  such that*

$$\Phi(x^{k+1}, y^{k+1}) > \Phi_{\text{target}}$$

and  $(x^{k+1}, y^{k+1})$  is not an  $(\eta_k/\eta/\varepsilon)$ -AKKT point of (8) is not greater than

$$\left\lceil \left( (\Phi(x^0, y^0) - \Phi_{\text{target}}) \left( \frac{\alpha^{p/(p+1)} \varepsilon}{L + \tau_2(L + \alpha)(p+1) + \theta} \right)^{-(p+1)/p} \right) \right\rceil. \quad (47)$$

Moreover, the number of functional evaluations per iteration is bounded above by

$$\left\lceil \log_{\tau_1} \left( \frac{\tau_2(L + \alpha)}{\rho_{\min}} \right) \right\rceil + 1.$$

*Proof.* As in [9, Lemma 3.1], we obtain that

$$\lim_{k \rightarrow \infty} \|x^{k+1} - x^k\| = \lim_{k \rightarrow \infty} \|y^{k+1} - y^k\| = 0.$$

As in [9, Lemma 3.1], we prove that the sequence of penalty parameters is bounded.

By the arguments presented above, we have that, for all  $k$ , there exist  $\lambda = \lambda^k \in \mathbb{R}^{n_H}$ ,  $\mu = \mu^k \in \mathbb{R}_+^{n_G}$ ,  $\beta = \beta^k \in \mathbb{R}_+^d$ , and  $\gamma = \gamma^k \in \mathbb{R}_+^d$  such that

$$\|\nabla f(x^{k+1}, y^{k+1}) + A_H^T \lambda + A_G^T \mu + \sum_{j=1}^d a_j \beta_j\| \leq O(\|(x^{k+1}, y^{k+1}) - (x^k, y^k)\|^p), \quad (48)$$

$$\sigma_j - (\pi^{-1})'(y_j^{k+1}) \beta_j - \gamma_j = 0, \quad j = 1, \dots, d, \quad (49)$$

$$H(x^{k+1}) = 0, \quad (50)$$

$$G(x^{k+1}) \leq 0, \quad (51)$$

$$g_j(x^{k+1}) - \pi^{-1}(y_j^{k+1}) \leq 0, \quad j = 1, \dots, d, \quad (52)$$

$$\beta_j = 0 \text{ whenever } g_j(x^{k+1}) - \pi^{-1}(y_j^{k+1}) < -c_\pi |y_j^{k+1} - y_j^k|^2, \quad (53)$$

$$y^{k+1} \geq 0, \quad (54)$$

$$\mu_j = 0 \text{ for all } j \text{ such that } G(x^{k+1})_j < 0, \quad (55)$$

$$\gamma_j = 0 \text{ for all } j \text{ such that } y_j^{k+1} > 0, \quad (56)$$

$$\mu, \beta, \gamma \geq 0. \quad (57)$$

By the sufficient descent condition and (48), there exists  $c_{\nabla} > 0$  such that, for all  $k$ ,

$$\Phi(x^{k+1}, y^{k+1}) \leq \Phi(x^k, y^k) - c_{\nabla} \left\| \nabla f(x^{k+1}, y^{k+1}) + A_{\text{H}}^T \lambda + A_{\text{G}}^T \mu + \sum_{j=1}^d a_j \beta_j \right\|^{\frac{p+1}{p}}. \quad (58)$$

Therefore, given  $\varepsilon > 0$ , the number of iterations such that  $\Phi(x^{k+1}, y^{k+1}) > \Phi_{\text{target}}$  and

$$\left\| \nabla f(x^{k+1}, y^{k+1}) + A_{\text{H}}^T \lambda + A_{\text{G}}^T \mu + \sum_{j=1}^d a_j \beta_j \right\| > \varepsilon \quad (59)$$

is, at most,

$$(\Phi(x^0, y^0) - \Phi_{\text{target}}) \times O(\varepsilon^{\frac{p+1}{p}}).$$

Given  $\xi > 0$ , by the descent condition (23), the number of iterations such that  $\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^{p+1} > \xi$  cannot exceed the quantity  $(\Phi(x^0, y^0) - \Phi_{\text{target}})/(\alpha\xi)$ . Therefore, the number of iterations such that  $\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^2 > \xi^{2/(p+1)}$  cannot exceed the quantity  $(\Phi(x^0, y^0) - \Phi_{\text{target}})/(\alpha\xi)$ . Therefore, the number of iterations such that  $\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^2 > \xi$  cannot exceed the quantity  $(\Phi(x^0, y^0) - \Phi_{\text{target}})/(\alpha\xi^{\frac{p+1}{2}})$ . Therefore, the number of iterations such that  $c_{\pi}\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^2 > c_{\pi}\xi$  cannot exceed the quantity  $(\Phi(x^0, y^0) - \Phi_{\text{target}})/(\alpha\xi^{\frac{p+1}{2}})$ . Therefore, the number of iterations such that  $c_{\pi}\|(x^{k+1} - x^k, y^{k+1} - y^k)\|^2 > \xi$  cannot exceed the quantity  $(\Phi(x^0, y^0) - \Phi_{\text{target}})/(\alpha(\xi/c_{\pi})^{\frac{p+1}{2}})$ . Therefore, after at most

$$\frac{(\Phi(x^0, y^0) - \Phi_{\text{target}})}{\alpha(\xi/c_{\pi})^{\frac{p+1}{2}}}$$

iterations, we have that all the iterates satisfy the approximate complementarity condition

$$\beta_j = 0 \text{ whenever } g_j(x^{k+1}) - \pi^{-1}(y_j^{k+1}) < -\xi$$

for all  $j = 1, \dots, d$ . This completes the proof.  $\square$

## 6 Numerical illustration

In this section, we present numerical experiment that illustrate in which way the use of the penalty function  $\pi(\cdot) = \sqrt{\cdot}$  promotes sparsity. Three toy problems are reformulated and solved with the general use nonlinear programming solver Algencon [7, 1] that applies to smooth problems.

### Problem 1. Regularized gradient minimization

The continuous version of this problem is to find a function  $u : [0, 1] \rightarrow \mathbb{R}$  that solves the problem

$$\text{Minimize } \|\nabla u\|^2 \text{ subject to } \int_0^1 u(t) dt = \int_0^1 \bar{u}(t),$$

where  $\bar{u}(t) = 400$  if  $t \in [0, 0.5]$  and  $\bar{u}(t) = 0$ , otherwise. In addition, we aim  $u$  to coincide with  $\bar{u}$  in “as much as possible”. After discretization, the problem becomes

$$\text{Minimize } \frac{1}{2} \sum_{i=2}^{99} (u_{i-1} - u_{i+1})^2 \text{ subject to } \sum_{i=1}^{100} u_i = \sum_{i=1}^{100} \bar{u}_i,$$

where  $\bar{u}_i = 400$  for  $i = 1, \dots, 50$  and  $\bar{u}_i = 0$  for  $i = 51, \dots, 100$ , with the additional constraint of having  $u_i = \bar{u}_i$ , for  $i = 1, \dots, 100$ , as many times as possible.

Solutions to the discretized version of the problem can be found by solving the reformulated problem given by

$$\begin{aligned} &\text{Minimize } \frac{1}{2} \sum_{i=2}^{99} (u_{i-1} - u_{i+1})^2 + \sigma \sum_{i=1}^{100} y_i \\ &\text{subject to } \quad \quad \quad \sum_{i=1}^{100} u_i = \sum_{i=1}^{100} \bar{u}_i \\ &\quad \quad \quad -y_i^2 \leq u_i - \bar{u}_i \leq y_i^2, \quad i = 1, \dots, 100 \end{aligned}$$

for different values of  $\sigma > 0$ . Note that the reformulation corresponds to penalizing  $\pi(|u_i - \bar{u}_i|)$ ,  $i = 1, \dots, 100$ , with  $\pi(\cdot) = \sqrt{\cdot}$ . Figure 1 shows a graphical representation of solutions to problems with  $\sigma \in \{0, 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, 100\}$ . Note that, the larger  $\sigma$  is, the larger the number of constraints of the form  $u_i = \bar{u}_i$  being satisfied.

## Problem 2. Two-dimensional dam

This problem is a two dimensional version of Problem 1. The discretized version of the problem is given by

$$\text{Minimize } \frac{1}{2} \sum_{i=2}^{29} \sum_{j=2}^{19} (4u_{ij} - u_{i,j-1} - u_{i,j+1} - u_{i-1,j} - u_{i+1,j})^2 \text{ subject to } \sum_{i=1}^{30} \sum_{j=1}^{20} u_{ij} = \sum_{i=1}^{30} \sum_{j=1}^{20} \bar{u}_{ij},$$

where

$$u_{ij} = \begin{cases} 400, & \text{if } j \leq 30 \left[ \frac{1}{2} + \frac{1}{10} \sin \left( 2\pi \left( \frac{i-1}{30-1} \right) \right) \right] \\ 0, & \text{otherwise,} \end{cases}$$

with the additional constraint of having  $u_{ij} = \bar{u}_{ij}$ , for  $i = 1, \dots, 30$  and  $j = 1, \dots, 20$ , as many times as possible. Solutions to the problem can be found by solving the reformulated problem given by

$$\begin{aligned} &\text{Minimize } \frac{1}{2} \sum_{i=2}^{99} (u_{i-1} - u_{i+1})^2 + \sigma \sum_{i=1}^{100} y_i \\ &\text{subject to } \quad \quad \quad \sum_{i=1}^{100} u_i = \sum_{i=1}^{30} \sum_{j=1}^{20} \bar{u}_{ij} \\ &\quad \quad \quad -y_{ij}^2 \leq u_{ij} - \bar{u}_{ij} \leq y_{ij}^2, \quad i = 1, \dots, 30, \quad j = 1, \dots, 20, \end{aligned}$$




Figure 1: Graphical representation of solutions to Problem 1 for different values of  $\sigma$ .

for different values of  $\sigma > 0$ . Note that the reformulation corresponds to penalizing  $\pi(|u_{ij} - \bar{u}_{ij}|)$ ,  $i = 1, \dots, 30$ ,  $j = 1, \dots, 20$ , with  $\pi(\cdot) = \sqrt{\cdot}$ . Figure 2 shows a graphical representation of solutions to problems with  $\sigma \in \{10^8, 10^7, \dots, 1, 10^{-1}, \dots, 10^{-8}\}$ . Once again, the larger  $\sigma$  is, the larger the number of constraints of the form  $u_{ij} = \bar{u}_{ij}$  being satisfied.

### Problem 3. Mass transportation

Assume that in a rectangle of  $n_x \times n_y$  pixels, to each pixel  $(i, j)$  there are associated functions  $P(i, j) \geq 0$  and  $Q(i, j) \geq 0$  such that

$$\sum_{i=1}^{n_x} \sum_{j=1}^{n_y} P(i, j) = \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} Q(i, j).$$

$P(i, j)$  represents white mass and  $Q(i, j)$  represents red mass. We goal is to find a function  $u((i, j), (k, \ell)) \geq 0$  that represents the transportation of white mass from pixel  $(i, j)$  to pixel  $(k, \ell)$ . There is no transportation of red mass. The function of transported mass  $u$  must be such that, at every pixel, the amount of white mass be equal to the amount of red mass. Therefore, for every pixel  $(i, j) \in V$ , we must have that

$$Q(i, j) = P(i, j) + \sum_{(k, \ell) \in V, (k, \ell) \neq (i, j)} u((k, \ell), (i, j)) - u((i, j), (k, \ell)),$$

where  $V = \{\{1, 2, \dots, n_x\} \times \{1, 2, \dots, n_y\}\}$ . With these constraints, the objective is to minimize the number of non-null transports, i.e. to minimize the number of  $u((i, j), (k, \ell))$ , for  $(i, j) \neq (k, \ell) \in V$ , that are positive.






Figure 2: Graphical representation of solutions to Problem 2 for different values of  $\sigma$ .

The reformulated problem is given by

$$\begin{aligned}
 &\text{Minimize} && \sigma \sum_{(i,j) \neq (k,\ell) \in V} y_{ijkl} \\
 &\text{subject to} && Q(i,j) = P(i,j) + \sum_{(k,\ell) \in V, (k,\ell) \neq (i,j)} u((k,\ell), (i,j)) - u((i,j), (k,\ell)), \forall (i,j) \in V \\
 &&& 0 \leq u((i,j), (k,\ell)) \leq y_{ijkl}^2, \forall (i,j) \neq (k,\ell) \in V.
 \end{aligned}$$

It corresponds to penalize  $\sqrt{\sum_{(i,j) \neq (k,\ell) \in V} u((i,j), (k,\ell))}$ , that represents the cost of the transportation process; its concavity representing economy of scale (decreasing marginal cost of transportation). Figure 3 illustrates solutions to a small instance of Problem 3 with  $\sigma = 0$  and  $\sigma > 0$ .




Figure 3: Graphical representation of solutions to a small instance of Problem 3.

## 7 Final remarks

In this work, we have introduced a smooth reformulation for constrained smooth problems with nonsmooth regularizations. The reformulation is entirely equivalent to the original problem both from the global as for the local point of view. Moreover, we were able to prove optimality conditions in which auxiliary variables do not appear at all.

Our main interest relies now in the application of these techniques to real problems in which the evaluation of the objective function is overwhelmingly more expensive than the evaluation of the constraints. This type of functions appear when PDE calculations are involved in the objective function evaluation and when the objective function is related to some phenomenon that takes place in real time. In this cases, model-intensive algorithms as the ones introduced in this paper may be useful. So, it is interesting to show that, from the theoretical point of view (convergence and complexity), these algorithms are well supported. Of course, smooth reformulations are valuable in these cases because one may take advantage of well-established constrained optimization software.

## References

- [1] R. Andreani, E. G. Birgin, J. M. Martínez and M. L. Schuverdt, On Augmented Lagrangian methods with general lower-level constraints, *SIAM Journal on Optimization* 18, pp. 1286–1309, 2008.
- [2] R. Andreani, G. Haeser, and J. M. Martínez, On sequential optimality conditions for smooth constrained optimization, *Optimization* 60, pp. 627–641, 2011.
- [3] D. P. Bertsekas, *Nonlinear Programming*, Athenas Scientific, 1999.
- [4] W. Bian and X. Chen, From sparse solutions of systems of equations to sparse modeling of signals and image, *SIAM Review* 51, pp. 34–81, 2009.
- [5] W. Bian and X. Chen, Linearly constrained non Lipschitz optimization for image restoration, *SIAM Journal on Imaging Sciences* 8, pp. 2294–2322, 2015.
- [6] W. Bian, X. Chen, and Y. Ye, Complexity analysis of interior point algorithms for non-Lipschitz and nonconvex minimization, *Mathematical Programming* 149, pp. 301–327, 2015.
- [7] E. G. Birgin and J. M. Martínez, *Practical Augmented Lagrangian Methods for Constrained Optimization*, Society for Industrial and Applied Mathematics, Philadelphia, 2014.
- [8] E. G. Birgin, J. L. Gardenghi, J. M. Martínez, S. A. Santos, and Ph. L. Toint, Worst-case evaluation complexity for unconstrained nonlinear optimization using higher order regularized models, *Mathematical Programming* 163, pp. 359–368, 2017.
- [9] E. G. Birgin and J. M. Martínez, On regularization and active-set methods with complexity for constrained optimization, *SIAM Journal on Optimization* 28, pp. 1367–1395, 2018.
- [10] S. Browne, The risk and rewards of minimizing shortfall probability, *Journal of Portfolio Management* 25, pp. 76–85, 1999.
- [11] E. Candes, M. Wakin, and S. Boyd, Enhancing sparsity by reweighted  $L_1$  minimization, *Journal of Fourier Analysis and Applications* 14, pp. 877–905, 2008.
- [12] C. Cartis, N. I. M. Gould, and Ph. L. Toint, Second-order optimality and beyond: characterization and evaluation complexity in convexly-constrained nonlinear optimization, *Foundations of Computational Mathematics* 18, pp. 1073–1107, 2018.
- [13] X. Chen, L. Guo, Z. Lu, and J. Ye, An augmented Lagrangian method for non-Lipschitz nonconvex programming, *SIAM Journal on Numerical Analysis* 55, pp. 168–193, 2017.
- [14] X. Chen, L. Niu, and Y. Yuan, Optimality conditions and a smoothing trust region Newton method for non-Lipschitz optimization, *SIAM Journal on Optimization* 23, pp. 1528–1552, 2013.
- [15] X. Chen and W. Zhou, Convergence of the reweighted  $L_1$  minimization algorithm for  $L_2$ - $L_p$  minimization, *Computational Optimization and Applications* 59, pp. 47–61, 2014.

- [16] C. Cortes and V. Vapnik, Support vector networks, *Machine Learning* 20, pp. 273–297, 1995.
- [17] J. Fan and R. Li, Variable selection via nonconcave penalized likelihood and its oracle properties, *Journal of American Statistical Association* 96, pp. 1348–1360, 2001.
- [18] J. Fang and H. Peng, Nonconcave penalized likelihood with a diverging number of parameters, *Annals of Statistics* 32, pp. 928–961, 2004.
- [19] L. E. Frank and J. H. Friedman, A statistical view of some chemometrics regression tools, *Technometrics* 35, pp. 109–135, 1993.
- [20] G. Haeser, H. Liu, and Y. Ye, Optimality condition and complexity analysis for linearly-constrained optimization without differentiability on the boundary, *Mathematical Programming*, to appear. (DOI: 10.1007/s10107-018-1290-4)
- [21] M. Lai and J. Wang, An unconstrained  $L_q$  minimization with  $0 < q \leq 1$  for sparse solution of underdetermined linear systems, *SIAM Journal on Optimization* 21, pp. 82–101, 2010.
- [22] Y. F. Liu, S. Ma, Y. H. Dai, and S. Zhang, A smoothing SQP framework for a class of composite  $L_q$  minimization over polyhedron, *Mathematical Programming* 158, pp. 467–500, 2016.
- [23] Z. Lu, Iterative reweighted minimization methods for  $L_p$  regularized unconstrained nonlinear programming, *Mathematical Programming* 147, pp. 277–307, 2014.
- [24] M. Nikolova, M. K. Ng, S. Zhang, and W. K. Ching, Efficient reconstruction of piecewise constant images using nonsmooth nonconvex minimization, *SIAM Journal on Imaging Sciences* 1, pp. 2–25, 2008.
- [25] J. Nocedal and S. J. Wright, *Numerical Optimization*, Springer-Verlag, New York, 2006.
- [26] C. R. Vogel, *Computational Methods for Inverse Problems*, Society for Industrial and Applied Mathematics, Philadelphia, 2002.
- [27] C. H. Zhang, Nearly unbiased variable selection under minimax nonconcave penalty, *The Annals of Statistics* 38, pp. 894–942, 2010.