

There is no largest proper operator ideal

Valentin Ferenczi, Universidade de São Paulo

IWOTA, August 16, 2021

Supported by Fapesp 2016/25574-8 and CNPq 303731/2019-2

- 1 Background
- 2 A new class of proper ideals

Operator ideals in the sense of Pietsch

All notions from the 1979 Pietsch's book "Operator ideals"

An operator ideal \mathcal{U} is a collection of subspaces $\mathcal{U}(X, Y)$ of $L(X, Y)$ for all X, Y Banach spaces, such that

- $\mathcal{U}(X, Y)$ contains $F(X, Y) = \{\text{finite rank operators}\}$
- (for all spaces Z, W and all appropriate operators T, V)
 $S \in \mathcal{U}(X, Y) \Rightarrow TSV \in \mathcal{U}(Z, W)$

In particular the class $\mathcal{U}(X) := \mathcal{U}(X, X)$ is a two-sided ideal of $L(X)$ in the usual sense.

Some classical operator ideals

- F =ideal of finite rank operators
- K =ideal of compact operators
- S =ideal of strictly singular
- In =ideal of *inessential* operators

$$F \subsetneq F^{\text{closure}} \subsetneq K \subsetneq S \subsetneq In$$

Inessential operators were defined by Kleinecke (63) when $X = Y$, by Pietsch (78) in general.

Definition

$S : X \rightarrow Y$ is *inessential* iff $\forall T : Y \rightarrow X$, $Id_X - TS$ is Fredholm

When there exist Fredholm operators between X and Y , then more intuitively, S is inessential iff $\forall T : X \rightarrow Y$

T Fredholm $\Leftrightarrow T + S$ Fredholm.

Definition (Pietsch)

A class \mathcal{U} is *proper* if for any Banach space X , Id_X belongs to $\mathcal{U} \Leftrightarrow X$ is finite dimensional

Therefore an ideal \mathcal{U} is proper iff for any *infinite-dimensional* space X , $\mathcal{U}(X)$ is a proper ideal of $L(X)$.

Fact

The ideals \mathcal{F} , \mathcal{K} , \mathcal{S} , \mathcal{I}_n are proper.

Two questions of Pietsch (79)

Question 1

Is I_n the largest proper operator ideal?

Question 2

Does there exist a largest proper operator ideal?

Theorem

The answer is no to both. Actually:

- *I_n is not even a maximal proper ideal, i.e. there exists V proper ideal such that $I_n \subsetneq V$*
- *moreover, there exist two proper ideals $I_n \subsetneq V_1, V_{-1}$ with $V_1 + V_{-1}$ not proper.*

We rely heavily on previous work by Aiena-González (00) and Gowers-Maurey (97) + notes by Maurey (96), and thank M. González and A. Martínez-Abejón for useful conversations.

Aiena-González's approach: improjective operators

Definition (Tarafdar 72)

An operator $T : X \rightarrow Y$ is *projective* if it induces an isomorphism between infinite dimensional *complemented* subspaces of X and Y respectively; and *improjective* otherwise.

Facts

- Fredholm operators between ∞ -dim spaces are projective.
- Imp is proper.
- We have the inclusion $\text{In} \subseteq \text{Imp}$; actually for an ideal U , U is proper $\Leftrightarrow U \subseteq \text{Imp}$.

Aiena-González (00) investigated whether $\text{In} = \text{Imp}$ or at least whether Imp is an ideal. In which case Imp would be the largest proper ideal.

Aiena-González's results: $\text{In} \subsetneq \text{Imp}$

- If X is an HI space then

$$L(X) = \text{Fredholm}(X) \cup S(X)$$

So $S(X) = \text{In}(X) = \text{Imp}(X)$ is the largest proper ideal of $L(X)$. This is useless for their purpose.

- Aiena-González note that if X is an indecomposable space, then

$$L(X) = \text{Fredholm}(X) \cup \text{Imp}(X)$$

So Aiena-González need an indecomposable, non HI space.

Theorem (Gowers-Maurey 97)

There exists a Banach space \mathbb{X} such that for any ∞ -dim subspace Y of \mathbb{X} , the following are equivalent

- *Y is finite codimensional*
- *Y is complemented in \mathbb{X}*
- *Y is isomorphic to \mathbb{X}*

The space \mathbb{X} is indecomposable and the isomorphism between \mathbb{X} and its hyperplanes is provided by the Right Shift operator R on the basis (e_n) of \mathbb{X} .

Theorem (Aiena-González, last line of p. 477)

We have $\text{In}(\mathbb{X}) \subsetneq \text{Imp}(\mathbb{X})$

Proof.

1 is in the essential spectrum of R , therefore $S_1 := Id_{\mathbb{X}} - R$ is not Fredholm. Therefore

- (a) it is improper
- (b) but it is essential. Indeed $2Id_{\mathbb{X}} - S_1 = Id_{\mathbb{X}} + R$ is not Fredholm, since -1 is also in the essential spectrum of R

Note: for future use and wlog, replace S_1 by some compact perturbation T_1 taking value in some Y with $\dim \mathbb{X}/Y = \infty$. □

Theorem (Aiena-González, last line of p. 477)

We have $\text{In}(\mathbb{X}) \subsetneq \text{Imp}(\mathbb{X})$

Proof.

1 is in the essential spectrum of R , therefore $S_1 := Id_{\mathbb{X}} - R$ is not Fredholm. Therefore

- (a) it is improjective
- (b) but it is essential. Indeed $2Id_{\mathbb{X}} - S_1 = Id_{\mathbb{X}} + R$ is not Fredholm, since -1 is also in the essential spectrum of R

Note: for future use and wlog, replace S_1 by some compact perturbation T_1 taking value in some Y with $\dim \mathbb{X}/Y = \infty$. \square

If Imp were an ideal, then it would be the largest proper ideal. But:

Proposition (Aiena-González, second line of p. 478)

Imp is not an ideal!

Proposition (Aiena-González)

Imp is not an ideal (actually $\text{Imp}(\mathbb{X})$ is not a linear subspace of $L(\mathbb{X})$).

But we can use their work by **shrinking** Imp to an ideal:

Proposition

There exist a proper ideal $U_1 \subseteq \text{Imp}$ such that $T_1 \in U_1$.

Since $T_1 \notin \text{In}$, let us denote $V_1 := \text{In} + U_1$:

Theorem

*The ideal In is not the largest proper ideal.
More precisely, $\text{In} \subsetneq V_1$ and V_1 is a proper ideal.*

But one can actually deduce more...

Indeed Aiena-González observe the same as above holds for $Id_{\mathbb{X}} - \lambda R$, $|\lambda| = 1$ (and not only for $\lambda = 1$) and associated compact perturbation T_{λ} . I.e. T_{λ} is improper $\forall \lambda$.
However $T_1 + T_{-1} = 2Id_{\mathbb{X}} + compact$ which is projective.
This is how they deduce that $\text{Imp}(\mathbb{X})$ is not a subspace of $L(\mathbb{X})$.
The case of T_{-1} is “the same as” T_1 so summing up:

Proposition

There exist two proper ideals U_1 and U_{-1} such that $T_i \in U_i$, for $i = -1, 1$.

Therefore $Id_{\mathbb{X}} \in U_1(\mathbb{X}) + U_{-1}(\mathbb{X})$ and it follows

Theorem

There exist two proper ideals whose sum is not proper. In particular there is no largest proper ideal.

A new kind of proper ideal

Summing up everything boils down to proving:

Proposition

For $i = -1, 1$, there exist a proper ideal U_i such that $T_i \in U_i$.

Definition

If X is an (∞ -dim) Banach space, $O_p(X)$ is the class of operators factorizing through X .

This is an ideal as soon as, e.g., $X \simeq X^2$.

(if $T = AB$ and $T' = A'B'$ then $T + T' = \begin{pmatrix} A & A' \end{pmatrix} \begin{pmatrix} B \\ B' \end{pmatrix}$)

However this is never proper if X is infinite dimensional...

Lemma

The class $\text{Op}(X) \cap \text{Op}(X')$ is proper iff no ∞ -dimensional complemented subspace of X is isomorphic to a complemented subspace of X'

(proof: $Id_Z \in \text{Op}(X)$ if and only if $Id_Z = AB$ where $A \in L(X, Z), B \in L(Z, X)$; this is equivalent to saying that Z embeds (by B) as a subspace of X which is complemented (by the projection BA). So $Id_Z \in \text{Op}(X) \cap \text{Op}(X')$ iff Z embeds complementably into both X and X' .)

Lemma

The class $\text{Op}(X) \cap \text{Op}(X')$ is proper iff no ∞ -dimensional complemented subspace of X is isomorphic to a complemented subspace of X'

(proof: $Id_Z \in \text{Op}(X)$ if and only if $Id_Z = AB$ where $A \in L(X, Z), B \in L(Z, X)$; this is equivalent to saying that Z embeds (by B) as a subspace of X which is complemented (by the projection BA). So $Id_Z \in \text{Op}(X) \cap \text{Op}(X')$ iff Z embeds complementably into both X and X' .)

So the properties of Gowers-Maurey's \mathbb{X} mean that:

Proposition

The class $\text{Op}(\mathbb{X}) \cap \text{Op}(Y_i)$ is proper.

Furthermore recall that T_i is defined on \mathbb{X} and takes values in ∞ -codimensional Y_i ; so $T_i \in \text{Op}(\mathbb{X}) \cap \text{Op}(Y_i)$.

The problem is....

The problem is.... that $\mathbb{X} \not\cong \mathbb{X}^2$ so we do not know whether $\text{Op}(\mathbb{X})$ and $\text{Op}(Y_i)$ are ideals!

The problem is.... that $\mathbb{X} \not\cong \mathbb{X}^2$ so we do not know whether $\text{Op}(\mathbb{X})$ and $\text{Op}(Y_i)$ are ideals!

So we **enlarge** $\text{Op}(X)$ to:

Definition

$\text{Op}^{<\omega}(X) := \bigcup_{n \in \mathbb{N}} \text{Op}(X^n)$ is the *ideal* of operators factorizing through *some power of X*,

and we enlarge $\text{Op}(\mathbb{X}) \cap \text{Op}(Y)$ to $\text{Op}^{<\omega}(\mathbb{X}) \cap \text{Op}^{<\omega}(Y)$ hoping it is still proper. And indeed:

Proposition

If Y is infinite codimensional in \mathbb{X} then the ideal $\text{Op}^{<\omega}(\mathbb{X}) \cap \text{Op}^{<\omega}(Y)$ is proper.

This is enough: let $U_i := \text{Op}^{<\omega}(\mathbb{X}) \cap \text{Op}^{<\omega}(\overline{\text{Im}T_i})$... then $T_i \in U_i$.

So our main result follows from the technical result:

Theorem

If Y is infinite codimensional in \mathbb{X} , and $m, n \in \mathbb{N}$, then no infinite dimensional complemented subspace Z of \mathbb{X}^m embeds complementably into Y^n .

(which involves extending the techniques of Gowers-Maurey to the multidimensional setting)

Question






An abstract **space ideal** \mathbb{A} (Pietsch) is a class of Banach spaces such that

- $F \subseteq \mathbb{A}$
- $E_1, E_2 \in \mathbb{A} \Rightarrow E_1 \oplus E_2 \in \mathbb{A}$
- $F \in \mathbb{A}$ and E embeds complementably in $F \Rightarrow E \in \mathbb{A}$.

Examples:

- \mathbb{F} , {separable spaces}, {hilbertian spaces},
- $\text{Space}(U) := \{X : \text{Id}_X \in U\}$, for any ideal U

Pietsch proves that a space ideal \mathbb{A} is always $\text{Space}(U)$ for some ideal U , and asks (Problem 2.2.8) whether there is always a largest U such that $\mathbb{A} = \text{Space}(U)$. We just proved that the answer is no for $\mathbb{A} = \mathbb{F}$ but seems open for other cases...

-  P. Aiena, M. González, *Examples of improper operators*, Math. Z. 233 (2000), 471–479.
-  B. Blackadar, *K-theory for operator algebras*, MSRI Publications 5, Springer Verlag 1986.
-  V. Ferenczi, *There is no largest proper operator ideal*, arXiv:2005.07672 or my webpage.
-  W.T. Gowers, B. Maurey, *Banach spaces with small spaces of operators*. Math Ann. (307) (1997), 543–568.
-  B. Maurey, *Operator theory and exotic Banach spaces*. 1996. Notes available on the author's webpage.