

ON QUADRUPLES OF PROJECTORS CONNECTED BY A LINEAR RELATION

K. A. Yusenko

UDC 517.98

We describe the set of $\gamma \in \mathbb{R}$ for which there exist quadruples of projectors P_i for a fixed collection of numbers $\alpha_i \in \mathbb{R}_+$, $i = \overline{1, 4}$, such that $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 + \alpha_4 P_4 = \gamma I$.

1. Introduction

Let $M_i = \{0 = \alpha_0^{(i)} < \alpha_1^{(i)} < \dots < \alpha_{m_i}^{(i)}\}$, $i = \overline{1, n}$, be given sets in \mathbb{R}_+ . Collections of self-adjoint operators $A_i = A_i^*$ with spectra $\sigma(A_i) \subset M_i$ and the sum divisible by a scalar operator were studied in many papers (see, e.g., the references in [1]). Considering these operators as representations of the generators of an involutive algebra, we obtain the equivalent problem of the description of irreducible $*$ -representations of the algebra

$$\mathcal{A}_{M_1, M_2, \dots, M_n; \gamma} = \mathbb{C} \langle a_1 \dots a_n \mid a_i = a_i^*, R_i(a_i) = 0, a_1 + a_2 + \dots + a_n = \gamma e \rangle,$$

where R_i is an annihilating polynomial of the corresponding generator a_i . This algebra is isomorphic to the algebra generated by the collection of projectors

$$\mathcal{P}_{M_1, M_2, \dots, M_n; \gamma} = \mathbb{C} \left\langle p_1^{(1)}, \dots, p_{m_1}^{(1)}, \dots, p_1^{(n)}, \dots, p_{m_n}^{(n)} \mid p_i^{(k)} = p_i^{(k)2} = p_i^{(k)*}, \sum_{i=1}^n \sum_{k=1}^{m_i} \alpha_k^{(i)} p_k^{(i)} = \gamma e, p_j^{(i)} p_k^{(i)} = 0 \right\rangle.$$

We can associate every algebra of this type with a graph Γ that has n branches meeting at a single vertex (the root of the graph). Each i th branch of the graph contains m_i vertices marked by numbers $\alpha_i^{(k)}$, $k = \overline{1, m_i}$. We associate the root of the graph with the number γ (for more details on the relationship between the problem considered and representations of graphs, see [2]). The vector $\chi = (\alpha_1^{(1)}, \dots, \alpha_{m_1}^{(1)}; \dots; \alpha_n^{(1)}, \dots, \alpha_{m_n}^{(n)})$ is called the character of the algebra $\mathcal{P}_{M_1, M_2, \dots, M_n; \gamma}$. The algebra $\mathcal{P}_{M_1, M_2, \dots, M_n; \gamma}$ is uniquely defined by its graph, character χ , and number γ . In what follows, we denote it by $\mathcal{P}_{\Gamma, \chi, \gamma}$. It was shown in [3] that, independently of χ and γ , the following assertions are true:

- (i) the algebra $\mathcal{P}_{\Gamma, \chi, \gamma}$ is finite-dimensional if Γ is a Dynkin diagram of the type D_n , E_6 , E_7 , or E_8 ;
- (ii) $\mathcal{P}_{\Gamma, \chi, \gamma}$ is an infinite-dimensional algebra of polynomial growth if Γ is an extended Dynkin diagram of the type \tilde{D}_4 , \tilde{E}_6 , \tilde{E}_7 , or \tilde{E}_8 ;
- (iii) for the other graphs, the algebra $\mathcal{P}_{\Gamma, \chi, \gamma}$ contains a free algebra with two self-adjoint generators.

Institute of Mathematics, Ukrainian Academy of Sciences, Kyiv.

Translated from Ukrains'kyi Matematychnyi Zhurnal, Vol. 58, No. 9, pp. 1289–1295, September, 2006. Original article submitted May 30, 2005.

In the investigation of $*$ -representations of these algebras, the following problems naturally arise:

- 1a. Describe the set of pairs $(\chi; \gamma)$ for which the algebra $\mathcal{P}_{\Gamma, \chi, \gamma}$ has a $*$ -representation; we denote this set by Σ_{Γ} .
- 1b. For every character χ , describe the set of γ for which the algebra $\mathcal{P}_{\Gamma, \chi, \gamma}$ has a $*$ -representation; we denote this set by $\Sigma_{\Gamma, \chi}$.
2. For every pair $(\chi; \gamma) \in \Sigma_{\Gamma}$, describe all irreducible (up to unitary equivalence) $*$ -representations of the algebra $\mathcal{P}_{\Gamma, \chi, \gamma}$.

The structure of the sets Σ_{Γ} and $\Sigma_{\Gamma, \chi}$ and $*$ -representations of $\mathcal{P}_{\Gamma, \chi, \gamma}$ essentially depends on the graph. In [4], problems 1 and 2 were considered for ordinary Dynkin diagrams. In the case where Γ is an extended Dynkin diagram, several authors described the sets $\Sigma_{\Gamma, \chi}$ for special characters (see the references in [1]). A complete description of the set $\Sigma_{\tilde{D}_4}$ is given in [5]. However, despite the fact that the set $\Sigma_{\tilde{D}_4, \chi}$ is a subset of $\Sigma_{\tilde{D}_4}$, it is difficult to obtain its description from the results of [5].

In the present paper, we give a direct description of the set $\Sigma_{\tilde{D}_4, \chi}$. We investigate in what cases this set is infinite [a necessary and sufficient condition is given, namely, all components of the character $\chi = (\alpha_1; \alpha_2; \alpha_3; \alpha_4)$ must satisfy the inequality $\alpha_i < (\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)/2$ (Sec. 3)], which enables us, by analogy with [5], to investigate in what cases the algebra $\mathcal{P}_{\tilde{D}_4, \chi, \gamma}$ has a representation on the hyperplane $\gamma = (\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)/2$ (Sec. 3). In Sec. 5, we describe the structure of the set $\Sigma_{\tilde{D}_4, \chi}$ for the special character $\chi_{\delta} = (1, 1, \delta, \delta)$.

1. Auxiliary Statements

Recall that the set of possible values γ for which there exist triples of projectors P_1, P_2, P_3 such that $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 = \gamma I$ for a fixed collection $\alpha_i \in \mathbb{R}, i = 1, 2, 3$, arranged in ascending order is described by the relation (see [4])

$$\Sigma_{D_4, (\alpha_1, \alpha_2, \alpha_3)} = \{0\} \cup \left\{ \sum_{i \in J} \alpha_i, J \subset \{1, 2, 3\} \right\} \cup \{(\alpha_1 + \alpha_2 + \alpha_3)/2\}. \tag{1}$$

Assume that $\alpha_3 < \alpha_1 + \alpha_2$. Otherwise the set does not contain the point $(\alpha_1 + \alpha_2 + \alpha_3)/2$.

Proposition 1. *If a collection of projectors P_1, P_2, \dots, P_n satisfies the equality*

$$\alpha_1 P_1 + \alpha_2 P_2 + \dots + \alpha_{n-1} P_{n-1} + P_n = I,$$

then the projector P_n commutes with the other projectors, i.e., $[P_n, P_i] = 0, i = \overline{1, n-1}$.

Proof. Consider the operator $Q = I - P_n$. Multiplying the equality $\alpha_1 P_1 + \alpha_2 P_2 + \dots + \alpha_{n-1} P_{n-1} = Q$ by P_n from the right and from the left and taking into account that $QP_n = P_nQ = 0$, we get

$$\alpha_1 P_n P_1 P_n + \alpha_2 P_n P_2 P_n + \dots + \alpha_3 P_n P_{n-1} P_n = 0.$$

Since all operators $P_n P_i P_n$ are nonnegative, we have $P_n P_i P_n = 0$, $i = \overline{1, n-1}$. One can verify that $P_n P_i Q P_i P_n = 0$. Therefore, $P_n P_i Q = Q P_i P_n = 0$. Taking into account that $P_n P_i Q = P_n P_i (I - P_n) = P_n P_i$, we get $P_n P_i = 0$. By analogy, one can show that $P_i P_n = 0$. This means that $[P_n, P_i] = 0$, $i = \overline{1, n-1}$.

Corollary 1. *There exists a collection of projectors P_i , $i = \overline{1, 4}$, such that $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 + \alpha_4 P_4 = \alpha_4 I$.*

2. Quadruple of Projectors and Coxeter Functors

Let a collection of numbers $\alpha_i \in \mathbb{R}$, $i = \overline{1, 4}$, be given. Our aim is to describe the set of γ for which there exist quadruples of projectors P_i , $i = \overline{1, 4}$, such that $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 + \alpha_4 P_4 = \gamma I$.

We can associate these quadruples of projectors with the associative \mathbb{C} -algebra $\mathcal{P}_{\tilde{D}_4, \chi, \gamma}$ generated by the generators $\{p_i\}_{i=1}^4$ and the relations

$$p_i = p_i^2 = p_i^*,$$

$$\alpha_1 p_1 + \alpha_2 p_2 + \alpha_3 p_3 + \alpha_4 p_4 = \gamma e,$$

where $\chi = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ is the character of the algebra (we assume that $\alpha_1 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4$). Then the problem can be reformulated as follows: For every character, describe the set $\Sigma_{\tilde{D}_4, \chi}$ of γ for which the algebra $\mathcal{P}_{\tilde{D}_4, \chi, \gamma}$ has a $*$ -representation. Let χ_i denote the i th component of the character χ and let $\alpha = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$.

The set $\Sigma_{\tilde{D}_4, \chi}$ possesses the following properties (see [6]):

- (i) $\Sigma_{\tilde{D}_4, \chi} \subset [0, \alpha]$;
- (ii) $\Sigma_{\tilde{D}_4, \chi} \ni \sum_{i \in J} \alpha_i$, $J \subset \{0, 1, 2, 3, 4\}$;
- (iii) $\tau \in \Sigma_{\tilde{D}_4, \chi} \Leftrightarrow \alpha - \tau \in \Sigma_{\tilde{D}_4, \chi}$.

Since the set $\Sigma_{\tilde{D}_4, \chi}$ is symmetric with respect to $\alpha/2$ [property (iii)], we study the set $\Sigma_{\tilde{D}_4, \chi} \cap [0; \alpha/2)$.

To investigate the set $\Sigma_{\tilde{D}_4, \chi}$, we use the method of Coxeter functors (introduced in [6]), which establish an equivalence between the categories of $*$ -representations $\text{Rep } \mathcal{P}_{\tilde{D}_4, \chi, \gamma}$ for different values of the parameters χ and γ . The linear T and hyperbolic S functors were constructed in [6]. The action of these functors between the categories generates the action on the pair $(\chi; \gamma)$:

$$S: (\chi; \gamma) \mapsto (\gamma - \alpha_1, \gamma - \alpha_2, \gamma - \alpha_3, \gamma - \alpha_4; \gamma),$$

$$T: (\chi; \gamma) \mapsto (\alpha_1, \alpha_2, \alpha_3, \alpha_4; \alpha - \gamma).$$

Let $(\chi^{(n)}; \gamma^{(n)}) = (ST)^n(\chi; \gamma)$ and $\lambda = \alpha/2 - \gamma$. Then the following statement is true:

Proposition 2. *The components of the character $\chi^{(n)}$ and the number $\gamma^{(n)}$ are determined by the formulas*

$$\chi_i^{(2n-1)} = \frac{\alpha}{2} - \alpha_i - (2n - 1)\lambda, \quad \chi_i^{(2n)} = \alpha_i - 2n\lambda, \quad i = \overline{1, 4}, \tag{2}$$

$$\gamma^{(n)} = \frac{\alpha}{2} - (2n + 1)\lambda, \quad n \in \mathbb{N}. \tag{3}$$

To prove this proposition, it is necessary to write the action of the functor ST on the pair $(\chi; \gamma)$ and to use the method of mathematical induction.

Corollary 2. *For any $\gamma \in [0, \alpha/2)$, there exists $n \in \mathbb{N}$ such that either one of the components of the character $\chi^{(n)}$ or the number $\gamma^{(n)}$ is less than or equal to zero.*

Proof. Since $\lambda > 0$ for any $\gamma \in [0, \alpha/2)$, it follows from relations (2) and (3) that the sequences $\{\chi_i^{(2n)}\}_{n=1}^\infty$, $\{\chi_i^{(2n-1)}\}_{n=1}^\infty$, and $\{\gamma_i^{(n)}\}_{n=1}^\infty$ are infinitely decreasing, and, hence, there exists n for which Proposition 2 is true.

Theorem 1. *The number $\gamma \in [0, \alpha/2)$ belongs to the set $\Sigma_{\tilde{D}_4, \chi}$ if and only if there exist $n \in \mathbb{Z}_+$ and $j \in \{1, 2, 3, 4\}$ such that the following two conditions are satisfied:*

$$\chi_j^{(n)} \leq 0, \quad \chi_i^{(k)} > 0, \quad \gamma^{(k)} \geq 0 \quad \forall k < n, \tag{4}$$

$$\gamma^{(n-1)} \in \Sigma_{D_4, \chi^*}, \tag{5}$$

where the character χ^* is defined by the triple of coefficients $\chi_i^{(n-1)}$, $i = \overline{1, 4}$, $i \neq j$.

Proof. The proof of the theorem follows from Corollary 2 and the functoriality of the mapping (ST) used for the construction of the corresponding sequences.

3. Infinite Sets $\Sigma_{\tilde{D}_4, \chi}$ and Representations on a Hyperplane

Theorem 2. *The set $\Sigma_{\tilde{D}_4, \chi}$ contains an infinite subset Σ_∞ with limit point $\alpha/2$ if and only if $\alpha_i < \alpha/2$, $i = \overline{1, 4}$. If this condition is satisfied, then the following assertions are true:*

$$(i) \quad \Sigma_\infty = \left\{ \frac{\alpha}{2} - \frac{\alpha_1}{2n} \mid n \in \mathbb{N} \right\} \quad \text{if } \alpha_2 + \alpha_3 > \alpha_1 + \alpha_4;$$

$$(ii) \quad \Sigma_\infty = \left\{ \frac{\alpha}{2} - \frac{\alpha - 2\alpha_4}{2(2n-1)} \mid n \in \mathbb{N} \right\} \quad \text{if } \alpha_2 + \alpha_3 < \alpha_1 + \alpha_4;$$

$$(iii) \quad \Sigma_\infty = \left\{ \frac{\alpha}{2} - \frac{\alpha_1}{n} \mid n \in \mathbb{N} \right\} \quad \text{if } \alpha_2 + \alpha_3 = \alpha_1 + \alpha_4.$$

Proof. Necessity. If one of the coefficients satisfies the inequality $\alpha_i \geq \alpha/2$, then the corresponding projector P_i commutes with the other projectors and, hence, is equal to either 0 or I in an irreducible representation. In this case, the problem reduces to a triple (or, correspondingly, to a smaller number) of projectors. Therefore, by virtue of Theorem 1, the set $\Sigma_{\tilde{D}_4, \chi}$ is finite.

Sufficiency. Assume, e.g., that $\alpha_2 + \alpha_3 > \alpha_1 + \alpha_4$. We show that, for every $\gamma = \alpha/2 - \alpha_1/(2n)$, $n \in \mathbb{N}$, there exists a representation of the algebra $\mathcal{P}_{\tilde{D}_4, \chi, \gamma}$. For this γ , we have $\chi_1^{(2n)} = 0$, $\chi_i^{(2n-1)} > 0$, $i = \overline{1, 4}$, and $\chi_1^{(2n-1)} = \gamma^{(2n-1)}$ ($\chi_1^{(2n-1)}$ is equal to zero at the next step). According to Corollary 1, this algebra has a representation, and, hence, the initial algebra also has a representation. The case $\alpha_2 + \alpha_3 < \alpha_1 + \alpha_4$ is proved by analogy. If $\alpha_2 + \alpha_3 = \alpha_1 + \alpha_4$, then these two sets are infinite. With regard for the equalities

$$\frac{\alpha}{2} - \frac{\alpha - 2\alpha_4}{2(2n-1)} = \frac{\alpha}{2} - \frac{2\alpha_1 + 2\alpha_4 - 2\alpha_4}{2(2n-1)} = \frac{\alpha}{2} - \frac{\alpha_1}{2n-1},$$

we get

$$\Sigma_\infty = \left\{ \frac{\alpha}{2} - \frac{\alpha_1}{n} \mid n \in \mathbb{N} \right\}.$$

Remark 1. By analogy, we can show that, in the case where the condition $\alpha_i < \alpha/2$, $i = \overline{1, 4}$, is satisfied, parallel with the infinite set Σ_∞ , $\Sigma_{\tilde{D}_4, \chi}$ also contains a finite set Σ_0 defined by the following rule:

$$(i) \quad \Sigma_0 = \left\{ \frac{\alpha}{2} - \frac{\alpha - 2\alpha_4}{2(2n-1)} \mid n < \frac{\alpha_1}{\alpha_2 + \alpha_3 - \alpha_1 - \alpha_4}, n \in \mathbb{N} \right\} \quad \text{if } \alpha_2 + \alpha_3 > \alpha_1 + \alpha_4;$$

$$(ii) \quad \Sigma_0 = \left\{ \frac{\alpha}{2} - \frac{\alpha_1}{2n} \mid n < \frac{\alpha_1}{\alpha_1 + \alpha_4 - \alpha_2 - \alpha_3}, n \in \mathbb{N} \right\} \quad \text{if } \alpha_2 + \alpha_3 < \alpha_1 + \alpha_4;$$

$$(iii) \quad \Sigma_0 = \emptyset \quad \text{if } \alpha_2 + \alpha_3 = \alpha_1 + \alpha_4.$$

Theorem 3. Suppose that the numbers $\alpha_i \in \mathbb{R}$, $i = \overline{1, 4}$, are such that $\alpha_i < \alpha/2$. Then there exists a collection of projectors P_1, P_2, P_3 , and P_4 such that $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 + \alpha_4 P_4 = \alpha I/2$.

Proof. It is necessary to show that the set $\Sigma_{\tilde{D}_4, \chi}$ contains the point $\alpha/2$. According to the Shulman theorem [7], the set $\Sigma_{\tilde{D}_4, \chi}$ is closed. The character $\chi = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ satisfies the conditions of Theo-

rem 2. Consequently, the set $\Sigma_{\tilde{D}_4, \chi}$ contains the infinite subset Σ_∞ , and, hence, by virtue of closedness, it also contains the limit point of the series $\alpha/2$.

Note that this theorem was presented in a somewhat different form by Kirichenko (see, e.g., [5]).

4. Subsets in the Set $\Sigma_{\tilde{D}_4, \chi}$

As shown in the proof of Theorem 2, the problem reduces to the case of a smaller number of projectors if at least one component of the character satisfies the inequality $\chi_i \geq \alpha/2$. Therefore, in what follows, we assume without loss of generality that $\chi_i < \alpha/2, i = \overline{1, 4}$.

To describe other sets, we use Theorem 1. Let $\gamma \in \Sigma_{\tilde{D}_4, \chi}$ and let k be such that condition (4) is satisfied. Two cases are possible, namely, $k = 2n$ and $k = 2n - 1$.

- 1. *Case $k = 2n$.* Using relations (2) and (3), we can rewrite condition (4) in the form of the following system of inequalities:

$$\begin{aligned} \lambda &> \frac{\alpha_1}{2n}, \quad \lambda < \frac{\alpha - 2\alpha_4}{2(2n - 1)}, \\ \lambda &< \frac{\alpha_1}{2(n - 1)}, \quad \lambda \leq \frac{\alpha}{2(4n - 1)}, \end{aligned} \tag{6}$$

where, as above, $\lambda = \alpha/2 - \gamma$. By virtue of Theorem 1, condition (5) can be rewritten as follows:

$$\begin{aligned} 0 &= \frac{\alpha}{2} - (4n - 1)\lambda, \\ \frac{\alpha}{2} - \alpha_i - (2n - 1)\lambda &= \frac{\alpha}{2} - (4n - 1)\lambda, \\ \frac{\alpha}{2} - \alpha_i - (2n - 1)\lambda + \frac{\alpha}{2} - \alpha_j - (2n - 1)\lambda &= \frac{\alpha}{2} - (4n - 1)\lambda, \quad i, j = 2, 3, 4, \quad i \neq j, \\ \sum_{i=2}^4 \left(\frac{\alpha}{2} - \alpha_i - (2n - 1)\lambda \right) &= \frac{\alpha}{2} - (4n - 1)\lambda, \\ \sum_{i=2}^4 \left(\frac{\alpha}{2} - \alpha_i - (2n - 1)\lambda \right) &= 2 \left(\frac{\alpha}{2} - (4n - 1)\lambda \right). \end{aligned} \tag{7}$$

Solving the system of inequalities (6) for every λ that satisfies one of the equations in (7), we obtain the following subsets in $\Sigma_{\tilde{D}_4, \chi}$:

$$\Sigma_1 = \left\{ \frac{\alpha}{2} - \frac{\alpha}{2(4n - 1)} \mid n < \frac{\alpha_4}{4\alpha_4 - \alpha}, n < \frac{\alpha - \alpha_1}{\alpha - 4\alpha_1}, n \in \mathbb{N} \right\},$$

$$\Sigma_2^i = \left\{ \frac{\alpha}{2} - \frac{\alpha_i}{2n} \mid n < \frac{\alpha_i}{2\alpha_i + 2\alpha_4 - \alpha}, n < \frac{\alpha_i}{\alpha_i - \alpha_1}, n < \frac{\alpha_i}{4\alpha_i - \alpha}, n \in \mathbb{N} \right\},$$

$$\Sigma_3 = \left\{ \frac{\alpha}{2} - \frac{\alpha - 2\alpha_1}{2(2n+1)} \mid n < \frac{\alpha - \alpha_1}{\alpha - 4\alpha_1}, n < \frac{\alpha_2 + \alpha_3}{2(\alpha_4 - \alpha_1)}, n(4\alpha_i - \alpha) < \alpha_i, n \in \mathbb{N} \right\}, \quad i = 2, 3, 4.$$

2. Case $k = 2n + 1$. Reasoning as in the previous case, we obtain the system of inequalities

$$\lambda > \frac{\alpha - 2\alpha_4}{2(2n+1)}, \quad \lambda < \frac{\alpha_1}{2n}, \tag{8}$$

$$\lambda < \frac{\alpha - 2\alpha_4}{2(2n-1)}, \quad \lambda \leq \frac{\alpha}{2(4n+1)}$$

and the equations

$$0 = \frac{\alpha}{2} - (4n+1)\lambda,$$

$$\alpha_i - 2n\lambda = \frac{\alpha}{2} - (4n+1)\lambda,$$

$$\alpha_i - 2n\lambda + \alpha_j - 2n\lambda = \frac{\alpha}{2} - (4n+1)\lambda, \tag{9}$$

$$\sum_{i=1}^3 \alpha_i - 2n\lambda = \frac{\alpha}{2} - (4n+1)\lambda,$$

$$\sum_{i=1}^3 \alpha_i - 2n\lambda = 2\left(\frac{\alpha}{2} - (4n+1)\lambda\right), \quad i, j = 1, 2, 3, \quad i \neq j.$$

Solving the system of inequalities (8) for every λ that satisfies one of the equations (9), we obtain the following subsets in $\Sigma_{\tilde{D}_4, \chi}$:

$$\Sigma_4 = \left\{ \frac{\alpha}{2} - \frac{\alpha}{2(4n+1)} \mid n < \frac{\alpha - \alpha_4}{4\alpha_4 - \alpha}, n < \frac{\alpha_1}{\alpha - 4\alpha_1}, n \in \mathbb{N} \cup \{0\} \right\},$$

$$\Sigma_5^i = \left\{ \frac{\alpha}{2} - \frac{\alpha - 2\alpha_i}{2(2n+1)} \mid n < \frac{\alpha_1}{\alpha - 2\alpha_i - 2\alpha_1}, n < \frac{\alpha_i}{\alpha - 4\alpha_i}, n < \frac{\alpha - \alpha_4 - \alpha_i}{2(\alpha_4 - \alpha_i)}, n \in \mathbb{N} \cup \{0\} \right\}, \quad i = 1, 2, 3.$$

Thus, the structure of the set $\Sigma_{\tilde{D}_4, \chi}$ is completely described by the following theorem:

Theorem 4. *The set of γ for which the algebra $\mathcal{P}_{\tilde{D}_4, \chi, \gamma}$ has a representation is described by the following relation:*

$$\Sigma_{\tilde{D}_4, \chi} \cap [0; \alpha/2) = \Sigma_\infty \cup \Sigma_0 \cup \Sigma_1 \cup \Sigma_2^i \cup \Sigma_3 \cup \Sigma_4 \cup \Sigma_5^j, \quad i = 2, 3, 4, \quad j = 1, 2, 3.$$

The entire set $\Sigma_{\tilde{D}_4, \chi}$ is obtained by symmetric mapping with respect to $\alpha/2$ and adjunction of the point $\alpha/2$.

5. The Set $\Sigma_{\tilde{D}_4, \chi}$ for the Character $\chi_\delta = (1, 1, \delta, \delta)$

The structure of the set $\Sigma_{\tilde{D}_4, \chi}$ is considerably simplified if $\chi = \chi_1 = (1, 1, 1, 1)$ or $\chi_\delta = (1, 1, \delta, \delta)$ (see [5, 8]). We now show how to construct the set $\Sigma_{\tilde{D}_4, \chi}$ for the character $\chi_\delta = (1, 1, \delta, \delta)$ by using Theorems 2 and 4.

Using Proposition 2, we get

$$\Sigma_\infty = \left\{ 1 + \delta - \frac{1}{2n} \mid n \in \mathbb{N} \right\}, \quad \Sigma_0 = \emptyset.$$

The sets $\Sigma_2^i, \Sigma_5^j, i = 2, 3, 4, j = 1, 2, 3$, and Σ_3 do not exist, and the sets Σ_1 and Σ_4 take the form

$$\Sigma_1 = \left\{ 1 + \delta - \frac{1 + \delta}{4n - 1} \mid n < \frac{\delta}{2(\delta - 1)}, n \in \mathbb{N} \right\},$$

$$\Sigma_4 = \left\{ 1 + \delta - \frac{1 + \delta}{4n + 1} \mid n < \frac{1}{2(\delta - 1)}, n \in \mathbb{N} \cup \{0\} \right\}.$$

Thus,

$$\begin{aligned} \Sigma_{\tilde{D}_4, (1, 1, \delta, \delta)} \cap [0; \alpha/2) &= \left\{ 1 + \delta - \frac{1}{2n} \mid n \in \mathbb{N} \right\} \cup \left\{ 1 + \delta - \frac{1 + \delta}{4n - 1} \mid n < \frac{\delta}{2(\delta - 1)}, n \in \mathbb{N} \right\} \\ &\cup \left\{ 1 + \delta - \frac{1 + \delta}{4n + 1} \mid n < \frac{1}{2(\delta - 1)}, n \in \mathbb{N} \cup \{0\} \right\}. \end{aligned}$$

The author is deeply grateful to V. L. Ostrovs'kyi for the statement of the problem and to Yu. S. Samoilenko and Yu. P. Moskal'ova for helpful remarks.

This work was partially supported by the Ukrainian State Foundation for Fundamental Research (grant No. 01.07/071).

REFERENCES

1. M. V. Zavodovs'kyi and Yu. S. Samoilenko, "Theory of operators and involutive representations of algebras," *Ukr. Mat. Visn.*, **1**, No. 4, 532–547 (2004).
2. S. A. Kruglyak and A. V. Roiter, "Locally scalar representations of graphs in the category of Hilbert spaces," *Funkts. Anal. Prilozhen.*, **39**, No. 2, 13–30 (2005).
3. M. A. Vlasenko, A. S. Mellit, and Yu. S. Samoilenko, "On algebras generated by linearly connected generators with given spectrum," *Funkts. Anal. Prilozhen.*, **39**, No. 3, 14–27 (2005).
4. S. A. Kruglyak, S. V. Popovich, and Yu. S. Samoilenko, " $*$ -Representation of algebras associated with Dynkin graphs and Horn's problem," *Uchen. Zap. Tavrich. Univ., Ser. Mat. Mekh. Inform. Kiber.*, **16 (55)**, No. 2, 132–139 (2003).
5. A. A. Kirichenko, "On linear combinations of orthoprojectors," *Uchen. Zap. Tavrich. Univ., Ser. Mat. Mekh. Inform. Kiber.*, No. 2, 31–39 (2002).
6. S. A. Kruglyak, V. I. Rabanovich, and Yu. S. Samoilenko, "On sums of projectors," *Funkts. Anal. Prilozhen.*, **36**, No. 3, 20–35 (2002).
7. V. S. Shulman, "On representations of limit relations," *Meth. Funct. Anal. Top.*, No. 4, 85–86 (2001).
8. V. Ostrovskiy and Yu. Samoilenko, "Introduction to the theory of representations of finitely presented $*$ -algebras. I. Representations by bounded operators," *Revs. Math. Math Phys.*, **11**, Part 1 (1999).